We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Small Noncoding RNA Expression in Cancer

*Florian Guisier, Mateus Camargo Barros-Filho,
Leigha D. Rock, Flavia B. Constantino, Brenda C. Minatel,
Adam P. Sage, Erin A. Marshall, Victor D. Martinez
and Wan L. Lam*

Abstract

Despite an inability to encode proteins, small noncoding RNAs (sncRNAs) have critical functions in the regulation of gene expression. They have demonstrated roles in cancer development and progression and are frequently dysregulated. Here we review the biogenesis and mechanism of action, expression patterns, and detection methods of two types of sncRNAs frequently described in cancer: miRNAs and piRNAs. Both miRNAs and piRNAs have been observed to play both oncogenic and tumor-suppressive roles, with miRNAs acting to directly regulate the mRNA of key cancer-associated genes, while piRNAs play crucial roles in maintaining the integrity of the epigenetic landscape. Elucidating these important functions of sncRNAs in normal and cancer biology relies on numerous *in silico* workflows and tools to profile sncRNA expression. Thus, we also discuss the key detection methods for cancer-relevant sncRNAs, including the discovery of genes that have yet to be described.

Keywords: small noncoding RNAs, miRNAs, piRNAs, transcriptome, gene expression profiling, novel, cancer, neoplasms, computational biology

1. Introduction

The central dogma of molecular biology that has prevailed for many decades, states that genetic information flows from DNA to RNA to protein. Nevertheless, RNAs that do not encode proteins were discovered as early as the 1950s [1, 2]. While protein-coding genes represent less than 2% of the human genome, it has been established that ~90% of the genome can be transcribed [3].

Small noncoding (snc) RNAs refer to ncRNA species that are <200 nucleotides in length and can be further categorized by their shared molecular features and biological mechanisms of action (**Table 1**). SncRNAs have diverse structural and functional roles in the regulation of gene expression, RNA splicing, epigenetic processes and chromatin structure. Due to their broad roles, the deregulation of sncRNAs has been shown to be involved in human diseases, including cancer. MicroRNAs (miRNAs) and PIWI-interacting RNAs (piRNAs) are two of the most studied sncRNA species. Here we describe current knowledge in the biogenesis and mechanisms of action for these sncRNAs and their expression profiling in cancer.

1.1 Biogenesis

1.1.1 miRNA biogenesis

MiRNAs are transcribed by RNA polymerase II to produce primary miRNA (pri-miRNA) transcripts [4]. Pri-miRNAs are folded hairpin intermediary RNA structures that can harbor multiple mature miRNA sequences and even protein-coding exons [5]. After transcription, pri-miRNAs are then processed and cleaved into mature miRNAs through different pathways (**Figure 1a**). In the “canonical” pathway, pri-miRNAs go through two cleavage events: (i) in the nucleus, the RNaseIII enzyme Drosha cleaves the pri-miRNA hairpin at its base to generate a precursor miRNA (pre-miRNA, ~60 nt) [6] and (ii) the pre-miRNA is translocated to the cytoplasm by Exportin-5, where it is cleaved into two mature (~22 nt) miRNA

Types	Comments	Size (nt)	Ref
MicroRNAs (miRNAs)	Evolutionarily conserved, endogenous, single-stranded sncRNAs, derived from endogenous short hairpin transcripts	18–25	[17]
PIWI interacting RNAs (piRNAs)	Largest group; single-stranded ncRNAs; generated by a Dicer-independent mechanism; a uridine at the 5' end, 5' monophosphate, and 2'-O-methyl at the 3' end	21–36	[18]
Transfer RNAs and ribosomal RNAs	Often referred to as “housekeeping” RNAs; take part of the translation process in ribonucleoproteins		
Small nuclear RNAs (snRNAs)	Found within the splicing speckles and cajal bodies of the nucleus; role in processing pre-messenger RNA, regulation of transcription factors and maintaining telomeres	150	[19]
Small nucleolar RNAs (snoRNAs)	Regulators of rRNA stability and function; some snoRNAs regulate gene expression and silencing processes (i) <i>C/D box snoRNAs</i> (60–200 nt): catalyzing the 2'-O-ribose methylation of rRNA residues (ii) <i>H/ACA box snoRNAs</i> (120–250 nt): guiding pseudouridylation of rRNA (iii) <i>Small Cajal body specific RNAs</i> : functions as a Cajal-body localization signal	60–250	[20]
Small interfering RNAs (siRNAs)	Partially complementary passenger and guide RNA strands; involved in post-transcriptional gene silencing through the RISC-mediated degradation of mRNA targets	19–23	[21]
Transfer RNA Fragments (tRFs)	Generated by specific cleavage of tRNA transcripts; (i) <i>Stress induced tRFs</i> (31–40 nt): repress translation and modulate cellular stress-response; interact with AGO proteins to form complexes for RNA interference silencing (ii) <i>Smaller tRFs</i> (14–30 nt): biogenesis and function unclear; some interact with PIWI or AGO proteins	14–40	[22]
Y RNAs	Parts of the Ro ribonucleoprotein. Involved in DNA replication, RNA stability, and responses to stress	100	[23]
7SL RNAs	Component of the signal recognition particle (SRP) that mediates co-translational insertion of secretory proteins into the endoplasmic reticulum lumen		[24]
Small NF90 associated RNAs (SNaRs)	Interact with NF90's double-stranded RNA-binding motifs and act as transcriptional regulator	117	[25]
Vault RNAs (vtRNAs)	Associated in large ribonucleoprotein particles (Vaults); essential for intracellular trafficking	100	[26]

Table 1.
Classification of small noncoding RNAs.

Figure 1.
 Biogenesis of (A) miRNAs and (B) piRNAs.

molecules by Dicer (also an RNase III enzyme) [7]. The major alternative miRNA processing pathway is the Mirtron pathway [8]. Mirtrons are short hairpin introns with splice acceptor and donor sites. In this pathway, a splicing event takes place instead of cleavage by Drosha. Here, the Mirtron and canonical miRNA pathways converge. Thus, the Mirtron pathway is considered as Drosha-independent, but Dicer-dependent. Several other miRNA processing pathways have also been reported [9]. Co-transcribed miRNAs that share similar seed regions are considered as members of a miRNA family [10]. Mechanistically, either of the strands derived from a mature miRNA duplex can be loaded into the Argonaute (AGO) family of proteins (AGO1–4 in humans) in an ATP-dependent manner to form the RNA-induced silencing complex (RISC) [11]. Although one of the strands is usually preferentially incorporated, this varies according to context, and the sequence of the strand incorporated will determine the targets that will be recognized by RISC [12].

1.1.2 piRNA biogenesis

PiRNAs are typically transcribed from genomic regions called piRNA clusters, regions which are typically 50–100 kb long, contain mainly transposable DNA elements and their remnants, and are found in large pericentromeric or subtelomeric domains [13]. PiRNAs are generated by RNaseIII-independent pathways that do not involve double-stranded RNA precursors, through two main biogenesis pathways (**Figure 1b**). (i) *Primary processing pathway*: cleavage of long piRNA precursors, by PIWI proteins, preferentially at uridine residues [14]. The 3' ends of piRNAs harbor extra nucleotides, which are trimmed upon association with PIWI proteins [15]. Here, the lengths of mature primary piRNAs are determined and depend on the molecular size of PIWI proteins [16].

Upon maturation, the 3' ends of piRNAs are 2'-O-methylated by Hen1/Pimet, which is associated with PIWI proteins [27]. This modification maintains the stability of piRNAs *in vivo* and can be used as a distinguishing feature in piRNA studies [28]. (ii) *Ping-Pong cycle*: this pathway is initiated in the cytoplasm to produce “secondary” piRNAs. The PIWI protein-piRNA complex (loaded with primary piRNAs) together with AGO3 are responsible for cleaving both sense and antisense transposon transcripts. Secondary piRNAs result from these transposon fragments and are complementary to the first 10 nt of the loaded primary piRNA [29]. This complex shows a strong bias for uracil at the 5' end (1-U), and, accordingly,

Ago3-piRNAs tend to have adenosine at the 10th nucleotide from the 5' end (10-A). Thus, 1-U and 10-A are signature to piRNAs made via the Ping-Pong cycle [30]. The cleavage of transposons by the AGO3-piRISCs and Aub-piRISCs, and the generation of secondary piRNAs are the main mechanisms involved in the control of transcript levels and silencing of transposons [13, 31].

Each step of miRNA and piRNA biogenesis is subject to regulation [32]. Thus, examining the biogenesis pathways of these sncRNAs through high throughput sequencing techniques may uncover mechanisms of aberrant miRNA/piRNA expression and deregulation in many human diseases.

1.2 Mechanisms of action

1.2.1 miRNA-mediated mechanisms

Once assembled into RISC, the miRNA 5' seed region (between nucleotides 2–7) interacts with specific region(s) within the 3' untranslated region (3' UTR) of target messenger RNAs (mRNAs) [33]. A single miRNA can interact with multiple target mRNAs. Depending on the miRNA/mRNA complementarity, degradation or repression of the targeted mRNA(s) will be triggered [33]. Pairing with complete complementary target leads to cleavage of the target mRNA and subsequent miRNA and mRNA degradation [34]. However, pairing with imperfect complementarity can lead to AGO2-mediated RNA interference. The interference mechanisms include having: (i) the GW182 component of the RISC to recruit associated proteins that would deadenylate, decap and degrade the target mRNA [35], (ii) Eukaryotic Translation Initiation Factor 4A2 (eIF4A2) as a “roadblock” to inhibit the ribosome-scanning step of initiation [36], and (iii) translational activation through recruitment of AGO2 and FXR1 instead of GW182 [37]. Of note, the miRNA-RISC can shuttle between the cytoplasm and the nucleus through Importin-8 or Exportin-1, highlighting the ability of newly-transcribed miRNAs to act in different cellular compartments [38].

Beyond the regulation of their production, several processes modify miRNA function. MiRNAs have a functional role in transcriptional gene silencing through DNA modification [39], deposition of repressive histone marks [40], promoting a transcriptionally active chromatin state [41], and altering alternative splicing profiles [42]. Alternative splicing, alternative polyadenylation affecting 3' UTRs, and cell type-specific RNA binding proteins that affect target mRNA secondary structures can change the available pool of miRNA binding targets. Moreover, subcellular localization of a given miRNA-RISC modulates its ability to bind target mRNAs [43]. These cell-type and biological state-specific factors contribute to the specificity of miRNA. Lastly, miRNAs can be released into and detected in extracellular fluids, delivered to different cells, and so act as regulators in autocrine, paracrine and/or endocrine processes [44].

1.2.2 piRNA-mediated mechanisms

The most well-known function of piRNAs is the silencing of transposons in germline cells to ensure genome stability during gametogenesis [45]. Similar to the lesser-known function of miRNAs, piRNAs primarily act as guides for PIWI proteins and drive histone modifications promoting heterochromatin assembly and DNA methylation [46].

PIWI-proteins are mainly found in the nucleus and co-localize with Polycomb group protein, playing crucial roles as epigenetic modifiers [47]. Knockout of PIWI proteins decreases histone H3 lysine 9 methylation, a marker of repressed

gene expression [48]. The complementary sequence of the piRNAs is responsible for directing these proteins to the specific targets on the genome and recruiting epigenetic factors [49], supposedly participating in epigenetic control [47], cell metabolism [50] and genome stability [51]. Alterations in piRNA expression have significant implications to the biology of stem-cells and cancer [52].

2. MicroRNA expression profiling in cancer

2.1 MicroRNA detection

Various experimental approaches can be used for measuring miRNA expression levels. The most frequently used are quantitative PCR (qPCR), digital color-coded barcoding profiling, miRNA microarrays, and high-throughput RNA sequencing (RNA-seq) methods. Material considerations and experimental aims dictate which approach is optimal [53]. While qPCR is efficient in analyzing few miRNAs, array and sequencing based methods offer parallel analyses of multiple miRNAs. Experiments that aim to discover previously undescribed transcripts require RNA-seq approaches [54].

2.2 MicroRNA expression in cancer

RNA expression has been shown to be dysregulated in all stages of cancer and nearly every cancer type [55–57]. Genome-wide profiling has demonstrated that miRNA expression signatures are associated with tumor type, tumor grade and clinical outcomes; thus, miRNAs are potential candidates for diagnostic and prognostic biomarkers, as well as therapeutic targets [56, 58, 59]. In fact, miRNA expression signatures have been observed to be impacted by smoking status in lung adenocarcinoma patients [59]. Furthermore, the expression patterns of miRNAs may be able to supplement the diagnostic utility of mRNAs, particularly in key tumor features such as subtype identification [58, 60]. There are currently ~2400 human miRNA annotated in miRBase (http://www.mirbase.org/cgi-bin/mirna_summary.pl?org=hsa), and it is believed that they collectively regulate one third of the genome [61]. The development of high-throughput deep sequencing analysis platforms has enabled our ability to detect and characterize miRNAs, as well as to identify the impact of their deregulation [57]. A summary of miRNA databases and the tools available for gene expression profiling is provided in **Table 2**.

2.3 Identification of novel microRNA sequences

The annotated human miRNA transcriptome mainly contains abundant and conserved miRNA sequences. Therefore, cell lineage- and tissue-specific miRNAs, especially the less abundant species, may not necessarily be included in current miR-Base annotations [55]. Re-analyses of high-throughput sequencing data of human tissues, cancers and cell lines have resulted in large scale discoveries of previously unannotated miRNAs that are expressed in a tissue-specific manner [55, 62–64].

A wide range of stand-alone and web-based miRNA discovery bioinformatics tools have been designed to quantify miRNA expression and to predict miRNA candidates and their isoforms from small RNA sequencing data (**Table 2**). These tools align the small RNA sequences to reference genomes and predicts putative novel miRNAs precursors based on the molecular features of these sequences, such as their folding characteristics, the formation of hairpin structures and whether this precursor gives rise to the three products of miRNA processing by DICER: a 5'

Resource	Name	Description	Link
Gene expression databases	ArrayExpress	EMBL-EBI ArrayExpress functional genomics data	https://www.ebi.ac.uk/arrayexpress/
	GEO	NCBI Gene Expression Omnibus	https://www.ncbi.nlm.nih.gov/geo/
	Oncomine	Web applications for translational bioinformatics	https://www.oncomine.org/resource/login.html
	TCGA	NIH The Cancer Genome Atlas	https://portal.gdc.cancer.gov
miRNAs databases	miRBase	miRbase 22: the microRNA database	www.mirbase.org
	miRCancer	microRNA Cancer Association Database	http://mircancer.ecu.edu/
	SonamiR DB	Somatic mutations altering microRNA-ceRNA interactions	http://compbio.uthsc.edu/SomamiR/
	TransmiR	Transcription factor microRNA regulations	http://www.cuilab.cn/transmir
piRNAs databases	piRBase	piRNA annotation and function analyses	http://www.regulatoryrna.org/database/piRNA/
	piRNABank	Web analysis of mammalian and Drosophila piRNAs	http://pirnabank.ibab.ac.in/
	piRNA cluster database	Resource for genomic piRNAs clusters	http://www.smallrnagroup.uni-mainz.de/piRNAclusterDB.html
miRNA discovery tools	deepBase	Annotate and discover small, long and circular ncRNAs	http://rna.sysu.edu.cn/deepBase
	miRDeep	Identification of novel and known miRNAs in NGS data	https://www.mdc-berlin.de/n-rajewsky#t-data,software&resources
	miRMaster	miRNA analysis framework, novel miRNA detection, isoforms and variants search	https://ccb-compute.cs.uni-saarland.de/mirmaster/
	miRNAkey	Software for the analysis of miRNA sequencing data	http://ibis.tau.ac.il/miRNAkey/
	OASIS	Online small-RNA detection and prediction platform	http://oasis.dzne.de
	Tools4miRs	Curation of methods for miRNA analysis	https://tools4mirs.org/
miRNA target prediction tools	miRDB	miRNA target prediction and functional annotations	http://mirdb.org/
	miRTargetLink	Human microRNA-mRNA interaction networks	https://ccb-web.cs.uni-saarland.de/mirtargetlink/
	miRWalk	Online prediction of microRNA binding sites	http://mirwalk.umm.uni-heidelberg.de/
	pathDIP	Pathway enrichment analysis by online data integration	http://ophid.utoronto.mirDIP/
	Targetscan	Predict target sites of conserved miRNAs	http://www.targetscan.org/vert_72/

Table 2.
Resources for ncRNA profiling studies.

Figure 3. Pipeline for detection and characterization of known and novel miRNAs. A) An example of bioinformatic pipeline for the detection of miRNAs. B) Main features for assessment of the biological relevance of miRNAs.

The growing availability of high throughput next generation sequencing (NGS) data will not only lead to novel miRNA discovery but will allow us to further elucidate the role of miRNA expression in human biology and disease such as cancer.

3. PIWI-interacting RNA expression profiling in cancer

PiRNAs are known to act in an evolutionarily conserved innate protection mechanism against transposable elements in germ cell genomes [69]. Beyond the

piRNA functions described in germ cells, there is increasing evidence of multifaceted action not restricted to transposon silencing in somatic cells [70]. Although the function of piRNAs in somatic cells and their relationship with tumorigenesis and cancer progression are still unknown, many studies seek to evaluate PIWI proteins and piRNA expression in a variety of malignancies [71].

3.1 piRNA detection and resources

Since piRNAs resemble miRNAs in length and structure, the same expression profiling platforms are applicable, wherein small RNA sequencing, microarrays, and quantitative PCR are the most widely used. The identification of piRNAs is mainly performed by small RNA sequencing, through extracting the reads with the proper length (generally from 24 to 32 nucleotides) that present piRNA-like features [72]. As previously discussed, piRNAs are frequently identified by a uridine nucleotide in the first position, have an adenosine nucleotide at the 10th position, have a 2'-O-methylation at the 3' end, and are mapped in clusters in the genome [72]. Although the expression can be confirmed by *in situ* hybridization and Northern blotting [73], the co-immunoprecipitation assay is the gold standard technique [74]. This analysis allows the isolation and characterization of RNAs physically interacting with PIWI-proteins [74]. However, the lack of highly specific antibodies for human PIWI-proteins limits the discovery of relevant piRNAs [75]. Functional studies using knockdown or knockout experiments for newly discovered piRNAs are fundamental to elucidate the biological role of these sequences [73].

The increasing application of large-scale small RNA sequencing has enabled the discovery of a large amount of piRNAs. The most widely used piRNA compendiums are piRBase and piRNABank, which contain millions of annotated human piRNA sequences—8,438,265 and 11,147,151 annotated piRNAs to date, respectively (Table 2) [76, 77]. Despite the large number of annotated sequences in these databases and many studies describing piRNA expression in somatic and malignant tissues, this knowledge must be considered with caution. It has been demonstrated that different piRNA databases include some RNA fragments that have similar sizes and features to piRNAs, representing possible contaminants; yet, sncRNAs derived from tRNAs have been described to interact with PIWI2 and are deregulated in cancer [78].

PIWI-interacting RNAs regulate the expression of mRNAs by guiding PIWI-proteins [46]. Bioinformatics approaches have shown that approximately 28.5% of human mRNA sequences contain at least one retrotransposon sequence in their 3' UTRs, and those mRNAs can be post-transcriptionally regulated by piRNAs [79]. In addition, many piRNAs do not match transposon sequences, suggesting an even greater set of targets and functional roles for piRNAs [80]. In fact, cross-linking immunoprecipitation (CLIP) analyses unveils many nontransposon mRNAs engaged with PIWI proteins [81]. In *Caenorhabditis elegans*, it was previously demonstrated that piRNA action is analogous to miRNAs, in that seed sequences are required for mRNA targeting, but unlike miRNAs, piRNAs do not tolerate many mismatches out of this region [82]. Potential piRNA targets can be retrieved using algorithms initially designed for miRNAs, such as miRanda [83], where stringent alignment (≥ 170) and free-energy scores (≤ -20.0 kcal/mol) are required for piRNA analyses [84]. However, the identification of piRNA targets is very challenging, as the targeting rules are still unsolved [82].

3.2 piRNA profiles in cancer

PIWI proteins 1–4 and PIWI-related proteins (DDX4, HENMT1, MAEL and TDRD1) have been reported to be disrupted in tumor cell line and patient samples

[85, 86]. The sncRNA repertoire of cancer cell lines from the NCI-60 panel (59 cell lines from nine different tissues) was recently characterized, where piRNAs comprised the largest proportion of expressed transcripts, followed by miRNAs and snRNAs [62]. In lung cancer cell lines, it was previously described 555 differentially expressed piRNAs and piRNA-like sncRNAs (piRNA-Ls) compared with lung bronchial epithelial cell lines [87]. Among them, piR-L-163 was found to be down-regulated in cancer cell lines and interact with phosphorylated ERM, regulating cell proliferation, migration and invasion.

Interestingly, piRNA expression profiling studies in tumor tissues revealed that piRNAs can be influenced by etiologic factors, such as tobacco consumption and HPV infection in lung and head and neck cancer [88–90]. The piRNA transcriptome of 6260 samples (from 11 organs) from The Cancer Genome Atlas (TCGA) consortium was prior screened [56]. Tumor samples presented a higher number of expressed piRNAs (n = 522) compared to somatic non-neoplastic tissues (n = 273), suggesting their potential as biomarkers. RNA sequencing found piR-1245 to be overexpressed and demonstrate oncogenic roles in colorectal cancer, inducing proliferation, colony formation, invasion, and apoptosis resistance [91]. Several other piRNAs have been reported to be overexpressed in numerous human malignancies [92–95]. Alternatively, piRNAs have also been described to have anti-tumor effects. For example, piR-39980 was demonstrated through functional assays to decrease proliferation, migration, invasion, colony formation, and to induce apoptosis in fibrosarcoma cell lines upon piRNA-mimic transfection [96].

The role of piRNAs in the response to chemotherapy has also been addressed [97–99]. In PIWL2-knockout embryonic fibroblast mouse models, the commonly overexpressed gene *PIWL2* was demonstrated to facilitate chromatin acetylation and relaxation in response to cisplatin treatment, leading to enhanced DNA repair and highlighting its potential role in treatment resistance [97]. piR-FTH1 was reported to drive chemoresistance in breast tumor cell lines, where its repression could sensitize tumor cells to doxorubicin [98]. Similarly, inhibition of piR-L-138 can increase apoptosis in cisplatin-treated lung cancer cell lines and patient-derived xenografts [99].

4. Emerging roles of sncRNA as cancer biomarkers

Considering the tissue-specificity of miRNAs and piRNAs in cancerous and healthy samples [55, 56], several individual or sncRNA-sets have been proposed as diagnostic or prognostic markers [56]. A set of 24 miRNAs evaluated by qPCR has been shown to correctly discriminate malignant from benign thyroid nodules with high sensitivity and specificity, potentially avoiding unnecessary diagnostic thyroidectomies [100]. In gastric adenocarcinoma, a three-piRNA recurrence risk signature was reported, using the small RNA sequencing data from the TCGA database [101]. Similarly, a higher expression of piR-1245 was linked to a lower overall survival in three independent cohorts of colorectal cancer patients [91].

The use of sncRNAs as liquid biopsy cancer-markers is also under intense investigation [102, 103]. In fact, both miRNAs and piRNAs are detectable in human serum, as demonstrated by a recent study based on RNA sequencing analysis in 477 serum samples [104]. Moreover, sncRNAs are enriched in extracellular vesicles (miRNAs ~40%, piRNAs ~40%) [105], allowing for their export from the cell in which they were synthesized to affect cells at a distance [106]. Models based on miRNA and piRNA combinations were able to correctly classify colon and prostate cancer patients from healthy individuals [106]. A four-miRNA expression signature in the serum of triple negative breast cancer patients was also demonstrated to be

an optimal survival predictor [107]. Recently, a qPCR assay comprising two targets (piR-5937 and piR-28876) and one reference piRNA (piR-28131) was suggested to detect early colon cancer [108]. Despite the low piRNA levels in the serum of cancer patients, they presented better detection sensitivity than the currently used biomarkers such as CA19-9 and carcinoembryonic antigen (CEA).

Many studies are currently investigating the ability of miRNA/piRNA signatures to empower cancer screening through the prediction of cancer recurrence or progression, stratification of patients by prognosis, and prediction of tumor response to various treatments. However, more efforts are still needed to screen miRNA/piRNA biomarker candidates and further validate them in large cohorts.

5. Conclusions

Here, we summarized the roles of small noncoding RNAs in normal and disease molecular biology and highlighted the importance of developing high-throughput sncRNA-detection methods in genome analyses. Transcribed through a variety of mechanisms, these molecules act in the widespread and specific regulation of gene expression. However, before these results can be translated to the clinic many factors must still be considered, including the development of effective and specific delivery system for sncRNA-based therapeutics and the broad validation of these sequences in large external cohorts. As our ability to detect and validate these sequences develops, we will continue to uncover their biological functions and potential uses in the clinical management of many diseases, including cancer.

Acknowledgements

This work was supported by grants from the Canadian Institutes for Health Research (CIHR FDN-143345), and scholarships from CIHR, Vanier Canada, the BC Cancer Foundation, the Ligue nationale contre le cancer, the Fonds de Recherche en Santé Respiratoire (appel d'offres 2018 emis en commun avec la Fondation du Souffle), the Fondation Charles Nicolle, and the São Paulo Research Foundation (FAPESP 2015/17707-5 and 2018/06138-8). E.A.M. is a Vanier Canada Graduate Scholar.

Conflict of interest

The authors have no conflicts to declare.

IntechOpen

Author details

Florian Guisier^{1,2*†}, Mateus Camargo Barros-Filho^{1,3†}, Leigha D. Rock^{1,4,5†},
Flavia B. Constantino^{1,6}, Brenda C. Minatel¹, Adam P. Sage¹, Erin A. Marshall¹,
Victor D. Martinez¹ and Wan L. Lam¹

1 Department of Integrative Oncology, British Columbia Cancer Research Centre,
Vancouver, BC, Canada

2 Pneumology Department, Rouen University Hospital, Rouen, France

3 International Research Center, A.C. Camargo Cancer Center, Sao Paulo, SP, Brazil

4 Department of Cancer Control Research, British Columbia Cancer Research
Centre, Vancouver, BC, Canada

5 Department of Oral and Biological Medical Sciences, Faculty of Dentistry,
University of British Columbia, Vancouver, BC, Canada

6 Institute of Biosciences, São Paulo State University, Botucatu, SP, Brazil

*Address all correspondence to: fguisier@bccrc.ca

† These authors contributed equally to this work.

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Stark BC, Kole R, Bowman EJ, Altman S. Ribonuclease P: An enzyme with an essential RNA component. *Proceedings of the National Academy of Sciences of the United States of America*. 1978;**75**(8):3717-3721
- [2] Yang VW, Lerner MR, Steitz JA, Flint SJ. A small nuclear ribonucleoprotein is required for splicing of adenoviral early RNA sequences. *Proceedings of the National Academy of Sciences of the United States of America*. 1981;**78**(3):1371-1375
- [3] Carninci P, Kasukawa T, Katayama S, Gough J, Frith MC, Maeda N, et al. The transcriptional landscape of the mammalian genome. *Science*. 2005;**309**(5740):1559-1563
- [4] Macfarlane LA, Murphy PR. MicroRNA: Biogenesis, function and role in cancer. *Current Genomics*. 2010;**11**(7):537-561
- [5] Cech TR, Steitz JA. The noncoding RNA revolution-trashing old rules to forge new ones. *Cell*. 2014;**157**(1):77-94
- [6] Lee Y, Ahn C, Han J, Choi H, Kim J, Yim J, et al. The nuclear RNase III Drosha initiates microRNA processing. *Nature*. 2003;**425**(6956):415-419
- [7] Hammond SM, Bernstein E, Beach D, Hannon GJ. An RNA-directed nuclease mediates post-transcriptional gene silencing in *Drosophila* cells. *Nature*. 2000;**404**(6775):293-296
- [8] Ruby JG, Jan CH, Bartel DP. Intronic microRNA precursors that bypass Drosha processing. *Nature*. 2007;**448**(7149):83-86
- [9] Yang JS, Lai EC. Alternative miRNA biogenesis pathways and the interpretation of core miRNA pathway mutants. *Molecular Cell*. 2011;**43**(6):892-903
- [10] Tanzer A, Stadler PF. Molecular evolution of a microRNA cluster. *Journal of Molecular Biology*. 2004;**339**(2):327-335
- [11] Yoda M, Kawamata T, Paroo Z, Ye X, Iwasaki S, Liu Q, et al. ATP-dependent human RISC assembly pathways. *Nature Structural & Molecular Biology*. 2010;**17**(1):17-23
- [12] Meijer HA, Smith EM, Bushell M. Regulation of miRNA strand selection: Follow the leader? *Biochemical Society Transactions*. 2014;**42**(4):1135-1140
- [13] Brennecke J, Aravin AA, Stark A, Dus M, Kellis M, Sachidanandam R, et al. Discrete small RNA-generating loci as master regulators of transposon activity in *Drosophila*. *Cell*. 2007;**128**(6):1089-1103
- [14] Weick EM, Miska EA. piRNAs: From biogenesis to function. *Development*. 2014;**141**(18):3458-3471
- [15] Vourekas A, Zheng Q, Alexiou P, Maragkakis M, Kirino Y, Gregory BD, et al. Mili and Miwi target RNA repertoire reveals piRNA biogenesis and function of Miwi in spermiogenesis. *Nature Structural & Molecular Biology*. 2012;**19**(8):773-781
- [16] Ishizu H, Siomi H, Siomi MC. Biology of PIWI-interacting RNAs: New insights into biogenesis and function inside and outside of germlines. *Genes & Development*. 2012;**26**(21):2361-2373
- [17] Friedman RC, Farh KK, Burge CB, Bartel DP. Most mammalian mRNAs are conserved targets of microRNAs. *Genome Research*. 2009;**19**(1):92-105
- [18] Thomson T, Lin H. The biogenesis and function of PIWI proteins and piRNAs: Progress and prospect. *Annual*

Review of Cell and Developmental Biology. 2009;**25**:355-376

[19] Valadkhan S, Gunawardane LS. Role of small nuclear RNAs in eukaryotic gene expression. *Essays in Biochemistry*. 2013;**54**:79-90

[20] Brameier M, Herwig A, Reinhardt R, Walter L, Gruber J. Human box C/D snoRNAs with miRNA like functions: Expanding the range of regulatory RNAs. *Nucleic Acids Research*. 2011;**39**(2):675-686

[21] Watanabe T, Totoki Y, Toyoda A, Kaneda M, Kuramochi-Miyagawa S, Obata Y, et al. Endogenous siRNAs from naturally formed dsRNAs regulate transcripts in mouse oocytes. *Nature*. 2008;**453**(7194):539-543

[22] Ivanov P, Emara MM, Villen J, Gygi SP, Anderson P. Angiogenin-induced tRNA fragments inhibit translation initiation. *Molecular Cell*. 2011;**43**(4):613-623

[23] Wolin SL, Steitz JA. Genes for two small cytoplasmic Ro RNAs are adjacent and appear to be single-copy in the human genome. *Cell*. 1983;**32**(3):735-744

[24] Ullu E, Weiner AM. Human genes and pseudogenes for the 7SL RNA component of signal recognition particle. *The EMBO Journal*. 1984;**3**(13):3303-3310

[25] Parrott AM, Mathews MB. Novel rapidly evolving hominid RNAs bind nuclear factor 90 and display tissue-restricted distribution. *Nucleic Acids Research*. 2007;**35**(18):6249-6258

[26] Stadler PF, Chen JJ, Hackermuller J, Hoffmann S, Horn F, Khaitovich P, et al. Evolution of vault RNAs. *Molecular Biology and Evolution*. 2009;**26**(9):1975-1991

[27] Horwich MD, Li C, Matranga C, Vagin V, Farley G, Wang P, et al. The

Drosophila RNA methyltransferase, DmHen1, modifies germline piRNAs and single-stranded siRNAs in RISC. *Current Biology*. 2007;**17**(14):1265-1272

[28] Lim SL, Qu ZP, Kortschak RD, Lawrence DM, Geoghegan J, Hempfling AL, et al. HENMT1 and piRNA stability are required for adult male germ cell transposon repression and to define the spermatogenic program in the mouse. *PLoS Genetics*. 2015;**11**(10):e1005620

[29] Czech B, Hannon GJ. One loop to rule them all: The ping-pong cycle and piRNA-guided silencing. *Trends in Biochemical Sciences*. 2016;**41**(4):324-337

[30] Grimson A, Srivastava M, Fahey B, Woodcroft BJ, Chiang HR, King N, et al. Early origins and evolution of microRNAs and Piwi-interacting RNAs in animals. *Nature*. 2008;**455**(7217):1193-1197

[31] Gunawardane LS, Saito K, Nishida KM, Miyoshi K, Kawamura Y, Nagami T, et al. A slicer-mediated mechanism for repeat-associated siRNA 5' end formation in *Drosophila*. *Science*. 2007;**315**(5818):1587-1590

[32] Kim VN, Han J, Siomi MC. Biogenesis of small RNAs in animals. *Nature Reviews. Molecular Cell Biology*. 2009;**10**(2):126-139

[33] Bartel DP. MicroRNAs: Target recognition and regulatory functions. *Cell*. 2009;**136**(2):215-233

[34] Hutvagner G, Simard MJ. Argonaute proteins: Key players in RNA silencing. *Nature Reviews. Molecular Cell Biology*. 2008;**9**(1):22-32

[35] Behm-Ansmant I, Rehwinkel J, Doerks T, Stark A, Bork P, Izaurralde E. mRNA degradation by miRNAs and GW182 requires both CCR4:NOT deadenylase and DCP1:DCP2 decapping

complexes. *Genes & Development*. 2006;**20**(14):1885-1898

[36] Meijer HA, Kong YW, Lu WT, Wilczynska A, Spriggs RV, Robinson SW, et al. Translational repression and eIF4A2 activity are critical for microRNA-mediated gene regulation. *Science*. 2013;**340**(6128):82-85

[37] Truesdell SS, Mortensen RD, Seo M, Schroeder JC, Lee JH, LeTonqueze O, et al. MicroRNA-mediated mRNA translation activation in quiescent cells and oocytes involves recruitment of a nuclear microRNP. *Scientific Reports*. 2012;**2**:842

[38] Nishi K, Nishi A, Nagasawa T, Ui-Tei K. Human TNRC6A is an argonaute-navigator protein for microRNA-mediated gene silencing in the nucleus. *RNA*. 2013;**19**(1):17-35

[39] Volpe TA, Kidner C, Hall IM, Teng G, Grewal SI, Martienssen RA. Regulation of heterochromatic silencing and histone H3 lysine-9 methylation by RNAi. *Science*. 2002;**297**(5588):1833-1837

[40] Benhamed M, Herbig U, Ye T, Dejean A, Bischof O. Senescence is an endogenous trigger for microRNA-directed transcriptional gene silencing in human cells. *Nature Cell Biology*. 2012;**14**(3):266-275

[41] Xiao M, Li J, Li W, Wang Y, Wu F, Xi Y, et al. MicroRNAs activate gene transcription epigenetically as an enhancer trigger. *RNA Biology*. 2017;**14**(10):1326-1334

[42] Allo M, Agirre E, Bessonov S, Bertucci P, Gomez Acuna L, Buggiano V, et al. Argonaute-1 binds transcriptional enhancers and controls constitutive and alternative splicing in human cells. *Proceedings of the National Academy of Sciences of the United States of America*. 2014;**111**(44):15622-15629

[43] Bottini S, Hamouda-Tekaya N, Mategot R, Zaragosi LE, Audebert S, Pisano S, et al. Post-transcriptional gene silencing mediated by microRNAs is controlled by nucleoplasmic Sfpq. *Nature Communications*. 2017;**8**(1):1189

[44] Iftikhar H, Carney GE. Evidence and potential in vivo functions for biofluid miRNAs: From expression profiling to functional testing: Potential roles of extracellular miRNAs as indicators of physiological change and as agents of intercellular information exchange. *BioEssays*. 2016;**38**(4):367-378

[45] Khurana JS, Theurkauf W. piRNAs, transposon silencing, and *Drosophila* germline development. *The Journal of Cell Biology*. 2010;**191**(5):905-913

[46] Sienski G, Donertas D, Brennecke J. Transcriptional silencing of transposons by Piwi and maelstrom and its impact on chromatin state and gene expression. *Cell*. 2012;**151**(5):964-980

[47] Grimaud C, Bantignies F, Pal-Bhadra M, Ghana P, Bhadra U, Cavalli G. RNAi components are required for nuclear clustering of polycomb group response elements. *Cell*. 2006;**124**(5):957-971

[48] Pal-Bhadra M, Leibovitch BA, Gandhi SG, Chikka MR, Bhadra U, Birchler JA, et al. Heterochromatic silencing and HP1 localization in *Drosophila* are dependent on the RNAi machinery. *Science*. 2004;**303**(5658):669-672

[49] Luteijn MJ, Ketting RF. PIWI-interacting RNAs: From generation to transgenerational epigenetics. *Nature Reviews. Genetics*. 2013;**14**(8):523-534

[50] Jones BC, Wood JG, Chang C, Tam AD, Franklin MJ, Siegel ER, et al. A somatic piRNA pathway in the *Drosophila* fat body ensures metabolic homeostasis and normal lifespan. *Nature Communications*. 2016;**7**:13856

- [51] Moyano M, Stefani G. piRNA involvement in genome stability and human cancer. *Journal of Hematology & Oncology*. 2015;**8**:38
- [52] Ross RJ, Weiner MM, Lin H. PIWI proteins and PIWI-interacting RNAs in the soma. *Nature*. 2014;**505**(7483):353-359
- [53] Mestdagh P, Hartmann N, Baeriswyl L, Andreasen D, Bernard N, Chen C, et al. Evaluation of quantitative miRNA expression platforms in the microRNA quality control (miRQC) study. *Nature Methods*. 2014;**11**(8):809-815
- [54] Wang Z, Gerstein M, Snyder M. RNA-Seq: A revolutionary tool for transcriptomics. *Nature Reviews. Genetics*. 2009;**10**(1):57-63
- [55] Londin E, Loher P, Telonis AG, Quann K, Clark P, Jing Y, et al. Analysis of 13 cell types reveals evidence for the expression of numerous novel primate- and tissue-specific microRNAs. *Proceedings of the National Academy of Sciences of the United States of America*. 2015;**112**(10):E1106-E1115
- [56] Martinez VD, Vucic EA, Thu KL, Hubaux R, Enfield KS, Pikor LA, et al. Unique somatic and malignant expression patterns implicate PIWI-interacting RNAs in cancer-type specific biology. *Scientific Reports*. 2015;**5**:10423
- [57] Bracken CP, Scott HS, Goodall GJ. A network-biology perspective of microRNA function and dysfunction in cancer. *Nature Reviews. Genetics*. 2016;**17**(12):719-732
- [58] Enfield KS, Pikor LA, Martinez VD, Lam WL. Mechanistic roles of noncoding RNAs in lung cancer biology and their clinical implications. *Genetics Research International*. 2012;**2012**:737416
- [59] Vucic EA, Thu KL, Pikor LA, Enfield KS, Yee J, English JC, et al. Smoking status impacts microRNA mediated prognosis and lung adenocarcinoma biology. *BMC Cancer*. 2014;**14**:778
- [60] Calin GA, Croce CM. MicroRNA signatures in human cancers. *Nature Reviews. Cancer*. 2006;**6**(11):857-866
- [61] Hammond SM. An overview of microRNAs. *Advanced Drug Delivery Reviews*. 2015;**87**:3-14
- [62] Marshall EA, Sage AP, Ng KW, Martinez VD, Firmino NS, Bennewith KL, et al. Small non-coding RNA transcriptome of the NCI-60 cell line panel. *Scientific Data*. 2017;**4**:170157
- [63] Minatel BC, Martinez VD, Ng KW, Sage AP, Tokar T, Marshall EA, et al. Large-scale discovery of previously undetected microRNAs specific to human liver. *Human Genomics*. 2018;**12**(1):16
- [64] Sage AP, Minatel BC, Marshall EA, Martinez VD, Stewart GL, Enfield KSS, et al. Expanding the miRNA transcriptome of human kidney and renal cell carcinoma. *International Journal of Genomics*. 2018;**2018**:6972397
- [65] Friedlander MR, Mackowiak SD, Li N, Chen W, Rajewsky N. miRDeep2 accurately identifies known and hundreds of novel microRNA genes in seven animal clades. *Nucleic Acids Research*. 2012;**40**(1):37-52
- [66] Griffiths-Jones S, Saini HK, van Dongen S, Enright AJ. miRBase: Tools for microRNA genomics. *Nucleic Acids Research*. 2008;**36**(Database issue):D154-D158
- [67] Mutz KO, Heilkenbrinker A, Lonne M, Walter JG, Stahl F. Transcriptome analysis using next-generation sequencing. *Current Opinion in Biotechnology*. 2013;**24**(1):22-30

- [68] Peterson SM, Thompson JA, Ufkin ML, Sathyanarayana P, Liaw L, Congdon CB. Common features of microRNA target prediction tools. *Frontiers in Genetics*. 2014;**5**:23
- [69] Ozata DM, Gainetdinov I, Zoch A, O'Carroll D, Zamore PD. PIWI-interacting RNAs: Small RNAs with big functions. *Nature Reviews. Genetics*. 2019;**20**:89-108
- [70] Cox DN, Chao A, Lin H. Piwi encodes a nucleoplasmic factor whose activity modulates the number and division rate of germline stem cells. *Development*. 2000;**127**(3):503-514
- [71] Krishnan P, Damaraju S. The challenges and opportunities in the clinical application of noncoding RNAs: The road map for miRNAs and piRNAs in cancer diagnostics and prognostics. *International Journal of Genomics*. 2018;**2018**:5848046
- [72] Zuo L, Wang Z, Tan Y, Chen X, Luo X. piRNAs and their functions in the brain. *International Journal of Human Genetics*. 2016;**16**(1-2):53-60
- [73] Lee EJ, Banerjee S, Zhou H, Jammalamadaka A, Arcila M, Manjunath BS, et al. Identification of piRNAs in the central nervous system. *RNA*. 2011;**17**(6):1090-1099
- [74] Girard A, Sachidanandam R, Hannon GJ, Carmell MA. A germline-specific class of small RNAs binds mammalian Piwi proteins. *Nature*. 2006;**442**(7099):199-202
- [75] Keam SP, Young PE, McCorkindale AL, Dang TH, Clancy JL, Humphreys DT, et al. The human Piwi protein Hiwi2 associates with tRNA-derived piRNAs in somatic cells. *Nucleic Acids Research*. 2014;**42**(14):8984-8995
- [76] Wang J, Zhang P, Lu Y, Li Y, Zheng Y, Kan Y, et al. piRBase: A comprehensive database of piRNA sequences. *Nucleic Acids Research*. 2019;**47**(D1):D175-D180
- [77] Sai Lakshmi S, Agrawal S. piRNABank: A web resource on classified and clustered Piwi-interacting RNAs. *Nucleic Acids Research*. 2008;**36**(Database issue):D173-D177
- [78] Balatti V, Nigita G, Veneziano D, Drusco A, Stein GS, Messier TL, et al. tsRNA signatures in cancer. *Proceedings of the National Academy of Sciences of the United States of America*. 2017;**114**(30):8071-8076
- [79] Watanabe T, Lin H. Posttranscriptional regulation of gene expression by Piwi proteins and piRNAs. *Molecular Cell*. 2014;**56**(1):18-27
- [80] Batista PJ, Ruby JG, Claycomb JM, Chiang R, Fahlgren N, Kasschau KD, et al. PRG-1 and 21U-RNAs interact to form the piRNA complex required for fertility in *C. Elegans*. *Molecular Cell*. 2008;**31**(1):67-78
- [81] Toombs JA, Sytnikova YA, Chirn GW, Ang I, Lau NC, Blower MD. *Xenopus* Piwi proteins interact with a broad proportion of the oocyte transcriptome. *RNA*. 2017;**23**(4):504-520
- [82] Zhang D, Tu S, Stubna M, Wu WS, Huang WC, Weng Z, et al. The piRNA targeting rules and the resistance to piRNA silencing in endogenous genes. *Science*. 2018;**359**(6375):587-592
- [83] Enright AJ, John B, Gaul U, Tuschl T, Sander C, Marks DS. MicroRNA targets in *Drosophila*. *Genome Biology*. 2003;**5**(1):R1
- [84] Hashim A, Rizzo F, Marchese G, Ravo M, Tarallo R, Nassa G, et al. RNA sequencing identifies specific PIWI-interacting small non-coding RNA expression patterns in breast cancer. *Oncotarget*. 2014;**5**(20):9901-9910

- [85] Yang Y, Zhang X, Song D, Wei J. Piwil2 modulates the invasion and metastasis of prostate cancer by regulating the expression of matrix metalloproteinase-9 and epithelial-mesenchymal transitions. *Oncology Letters*. 2015;**10**(3):1735-1740
- [86] Schudrowitz N, Takagi S, Wessel GM, Yajima M. Germline factor DDX4 functions in blood-derived cancer cell phenotypes. *Cancer Science*. 2017;**108**(8):1612-1619
- [87] Mei Y, Wang Y, Kumari P, Shetty AC, Clark D, Gable T, et al. A piRNA-like small RNA interacts with and modulates p-ERM proteins in human somatic cells. *Nature Communications*. 2015;**6**:7316
- [88] Firmino N, Martinez VD, Rowbotham DA, Enfield KSS, Bennewith KL, Lam WL. HPV status is associated with altered PIWI-interacting RNA expression pattern in head and neck cancer. *Oral Oncology*. 2016;**55**:43-48
- [89] Krishnan AR, Korrapati A, Zou AE, Qu Y, Wang XQ, Califano JA, et al. Smoking status regulates a novel panel of PIWI-interacting RNAs in head and neck squamous cell carcinoma. *Oral Oncology*. 2017;**65**:68-75
- [90] Nogueira Jorge NA, Wajnberg G, Ferreira CG, de Sa Carvalho B, Passetti F. snoRNA and piRNA expression levels modified by tobacco use in women with lung adenocarcinoma. *PLoS One*. 2017;**12**(8):e0183410
- [91] Weng W, Liu N, Toiyama Y, Kusunoki M, Nagasaka T, Fujiwara T, et al. Novel evidence for a PIWI-interacting RNA (piRNA) as an oncogenic mediator of disease progression, and a potential prognostic biomarker in colorectal cancer. *Molecular Cancer*. 2018;**17**(1):16
- [92] Li Y, Wu X, Gao H, Jin JM, Li AX, Kim YS, et al. Piwi-interacting RNAs (piRNAs) are Dysregulated in renal cell carcinoma and associated with tumor metastasis and cancer-specific survival. *Molecular Medicine*. 2015;**21**:381-388
- [93] Huang G, Hu H, Xue X, Shen S, Gao E, Guo G, et al. Altered expression of piRNAs and their relation with clinicopathologic features of breast cancer. *Clinical & Translational Oncology*. 2013;**15**(7):563-568
- [94] Cheng J, Guo JM, Xiao BX, Miao Y, Jiang Z, Zhou H, et al. piRNA, the new non-coding RNA, is aberrantly expressed in human cancer cells. *Clinica Chimica Acta*. 2011;**412**(17-18):1621-1625
- [95] Yan H, Wu QL, Sun CY, Ai LS, Deng J, Zhang L, et al. piRNA-823 contributes to tumorigenesis by regulating de novo DNA methylation and angiogenesis in multiple myeloma. *Leukemia*. 2015;**29**(1):196-206
- [96] Das B, Roy J, Jain N, Mallick B. Tumor suppressive activity of PIWI-interacting RNA in human fibrosarcoma mediated through repression of RRM2. *Molecular Carcinogenesis*. 2018;1-14
- [97] Wang QE, Han C, Milum K, Wani AA. Stem cell protein Piwil2 modulates chromatin modifications upon cisplatin treatment. *Mutation Research*. 2011;**708**(1-2):59-68
- [98] Wang Y, Gable T, Ma MZ, Clark D, Zhao J, Zhang Y, et al. A piRNA-like small RNA induces chemoresistance to cisplatin-based therapy by inhibiting apoptosis in lung squamous cell carcinoma. *Molecular Therapy. Nucleic Acids*. 2017;**6**:269-278
- [99] Balaratnam S, West N, Basu S. A piRNA utilizes HILI and HIWI2 mediated pathway to down-regulate ferritin heavy chain 1 mRNA in human somatic cells. *Nucleic Acids Research*. 2018;**46**(20):10635-10648
- [100] Lithwick-Yanai G, Dromi N, Shtabsky A, Morgenstern S, Strenov Y,

Feinmesser M, et al. Multicentre validation of a microRNA-based assay for diagnosing indeterminate thyroid nodules utilising fine needle aspirate smears. *Journal of Clinical Pathology*. 2017;**70**(6):500-507

[101] Martinez VD, Enfield KS, Rowbotham DA, Lam WL. An atlas of gastric PIWI-interacting RNA transcriptomes and their utility for identifying signatures of gastric cancer recurrence. *Gastric Cancer*. 2016;**19**(2):660-665

[102] Xu R, Rai A, Chen M, Suwakulsiri W, Greening DW, Simpson RJ. Extracellular vesicles in cancer—Implications for future improvements in cancer care. *Nature Reviews. Clinical Oncology*. 2018;**15**(10):617-638

[103] Anfossi S, Babayan A, Pantel K, Calin GA. Clinical utility of circulating non-coding RNAs—An update. *Nature Reviews. Clinical Oncology*. 2018;**15**(9):541-563

[104] Umu SU, Langseth H, Bucher-Johannessen C, Fromm B, Keller A, Meese E, et al. A comprehensive profile of circulating RNAs in human serum. *RNA Biology*. 2018;**15**(2):242-250

[105] Ren J, Zhou Q, Li H, Li J, Pang L, Su L, et al. Characterization of exosomal RNAs derived from human gastric cancer cells by deep sequencing. *Tumour Biology*. 2017;**39**(4):1010428317695012

[106] Yuan T, Huang X, Woodcock M, Du M, Dittmar R, Wang Y, et al. Plasma extracellular RNA profiles in healthy and cancer patients. *Scientific Reports*. 2016;**6**:19413

[107] Kleivi Sahlberg K, Bottai G, Naume B, Burwinkel B, Calin GA, Borresen-Dale AL, et al. A serum microRNA signature predicts tumor relapse and survival in triple-negative breast cancer patients. *Clinical Cancer Research*. 2015;**21**(5):1207-1214

[108] Vychytilova-Faltejskova P, Stitkovcova K, Radova L, Sachlova M, Kosarova Z, Slaba K, et al. Circulating PIWI-interacting RNAs piR-5937 and piR-28876 are promising diagnostic biomarkers of colon cancer. *Cancer Epidemiology, Biomarkers & Prevention*. 2018;**27**(9):1019-1028