We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Retinoic Acid in Ocular Growth Regulation

Jody A. Summers

Abstract

All-*trans*-retinoic acid (atRA) is a metabolite of vitamin A (retinol) and is required for growth and development of a variety of organ systems in all higher animals from fish to humans. Evidence is accumulating to suggest that atRA may also be an important molecular signal in the postnatal control of eye size. Choroidal synthesis of atRA is modulated during periods of visually-induced changes in ocular growth and has pronounced effects on eye growth and refraction in several animal models of myopia. Choroidal atRA synthesis is exclusively regulated by expression of the enzyme, retinaldehyde dehydrogenase 2 (RALDH2). In chicks and humans, RALDH2 is synthesized by a unique population of uncharacterized extravascular stromal cells concentrated in the proximal choroid. The identification of choroidal atRA and RALDH2 as visually induced ocular growth regulators provides the potential for new therapeutic targets for the treatment of childhood myopia. The objective of this chapter is to discuss what is presently known about atRA biosynthesis and transport in the eye during visually guided eye growth and how this research can contribute to a better understanding of the mechanisms underlying the development of myopia.

Keywords: retinoic acid, choroid, myopia, sclera, RALDH2, emmetropization

1. Introduction

All-*trans*-retinoic acid (atRA) is the transcriptionally active derivative of vitamin A. atRA is an essential signaling molecule for developmental processes of numerous organ systems including those of the brain, limbs, lungs, pancreas, heart, and eye in many organisms from fish to humans [1, 2]. With the advent of increasingly sensitive methods to measure endogenous concentrations of atRA, data is accumulating to suggest that atRA is also important in the growth and maintenance of a number of organ systems during postnatal and adult life [3–7]. Within the postnatal eye, atRA has been detected in the retina, where its synthesis was shown to be mediated by oxidation of the chromophore all-*trans*-retinaldehyde, released from bleached rhodopsin in the photoreceptor outer segments following exposure to light [8]. More recently, the choroid (the highly vascular layer between the retina and the sclera) has been shown to synthesize and accumulate high levels of atRA [9, 10]. A number of studies in several animal models suggest that choroidal atRA may be an important molecular signal for the control of postnatal ocular growth [9, 11–13]. We and others have demonstrated that in response to visual stimuli, ocular atRA synthesis is regulated exclusively via choroidal expression of the atRA synthesizing enzyme, retinaldehyde dehydrogenase 2 (RALDH2) by a unique population of cells [10, 14]. Furthermore, choroidally derived retinoic acid is transported to the

sclera (the outer connective tissue shell of the eye) by apolipoprotein A-1 (ApoA-1) which functions as a specific extracellular atRA-binding and transport protein in the eye [15, 16]. Once delivered to the sclera, we speculate that atRA regulates the transcription of many genes in the sclera to effect changes in scleral extracellular matrix remodeling, ocular size, and refraction.

This chapter therefore focuses primarily on the potential role of atRA on the control of postnatal growth of the eye, and implications for the development of new therapies for the control of myopia in children.

2. Retinoic acid is a vitamin A derivative

atRA is synthesized in two steps from vitamin A (all-*trans*-retinol). The first step produces all-*trans*-retinaldehyde from all-*trans*-retinol through the action of cytosolic or membrane bound alcohol dehydrogenases (ADH). The second, irreversible, step in atRA synthesis involves the oxidization of all-*trans*-retinaldehyde to atRA through the actions of the cytosolic retinaldehyde dehydrogenases (RALDH1, RALDH2, RALDH3; a.k.a. Aldh1 α 1, Aldh1 α 2, Aldh1 α 3) [16, 17]. Tissue concentrations of atRA are regulated by the activities of these synthesizing enzymes, as well as the atRA-metabolizing enzyme CYP26, a member of the cytochrome P450 family [17]. Cyp1B1 may also contribute to atRA synthesis in the chick embryo [18] via synthesis of all-*trans*-retinaldehyde and atRA from all-*trans*-retinol. Furthermore, the rate of these reactions is regulated by the availability of the substrates, the accessibility of the enzymes to their substrates, and the catalytic activity of the enzymes [19]. Once synthesized, atRA can act within its own cell of synthesis (autocrine signaling) or be transported to nearby cells (paracrine signaling) and bind with nuclear retinoic acid receptor complexes to directly control the transcriptional activity of more than 100 target genes [20].

3. Visual regulation of intraocular retinoic acid synthesis

3.1 Emmetropization: vision-dependent ocular growth regulation

Clinical and experimental evidence have indicated that postnatal eye growth is regulated, at least in part, by a vision-dependent “emmetropization” mechanism that acts to minimize refractive error through the coordinated regulation of the growth of the ocular tissues [21, 22]. Interruption of emmetropization in animal models, such as the chick, primate, and guinea pig, through the application of translucent occluders (form deprivation) causes a distortion in visual quality, which results in ocular growth and myopia through changes in the regulation of scleral extracellular matrix (ECM) remodeling [23–27]. Form deprivation-induced myopia is reversible; removal of the occluder and subsequent detection of myopic defocus results in a rapid cessation of axial growth and the eventual reestablishment of emmetropia (recovery) [24]. Even stronger evidence for the presence of an emmetropization mechanism comes from studies in which animals are fitted with either concave (minus) lenses or convex (plus) lenses that shifts the focal plane behind (hyperopic defocus) or in front of (myopic defocus) the retinal photoreceptors, respectively. In animals with functional emmetropization, the axial length of the lens-treated eye will increase or decrease until the retinal location has shifted to match that of the new focal plane [28–31]. The emmetropization mechanism does not require the central nervous system and appears to be regulated by locally produced chemical signals within the eye itself. When visual form deprivation is restricted to nasal or temporal visual

fields, excessive growth of the sclera is limited to that portion corresponding to the visually deprived part of the retina [32, 33]. Furthermore experimental myopia can be induced in animals lacking a functional optic nerve [34–36], suggesting that the central nervous system is not required for the development of myopia. It is now generally accepted that visually guided eye growth is regulated by a series of locally generated chemical events that begin in the retina in response to specific visual stimuli and terminate in the sclera where they result in scleral extracellular matrix (ECM) remodeling, changes in ocular length and refractive status [37–42]. Therefore the elucidation of the chemical events responsible for visually-induced changes in ocular growth is of great interest as it may provide new avenues for the development of therapies to slow or prevent the progression of myopia.

3.2 Choroidal retinoic acid: a potential ocular growth regulator

Several studies in a variety of animal models indicate that all-*trans*-retinoic acid (atRA) may be one of the chemical signals required for the regulation of eye growth during emmetropization [9, 11–13]. Mertz and Wallman [9] were the first to show that choroidal synthesis of atRA was increased in chick eyes during recovery from form deprivation myopia and following application of positive lenses (imposed myopic defocus), two visual conditions that cause a deceleration in ocular growth rates. Moreover, atRA was shown to be decreased in eyes undergoing form deprivation myopia and compensation for hyperopic defocus compared with the fellow control eye, conditions that stimulate ocular elongation. It was therefore suggested that choroidal atRA could act as a locally produced (within the eye) scleral growth modulator during visually guided ocular growth. atRA is an attractive candidate for a visually regulated ocular growth regulator because it is readily diffusible, has pronounced effects on scleral extracellular matrix metabolism, and exerts its effects through highly regulated, locally controlled synthesis and degradation.

Studies by Simon et al. [43] and Rada et al. [10] identified transcriptional changes in choroidal *RALDH2* in response to imposed defocus or recovery from induced myopia. *RALDH2* mRNA concentration was found to decrease in the choroid following treatment with negative lenses and to increase with positive lenses or during recovery from induced myopia. No changes were observed in the expression of the atRA metabolizing enzymes, *RALDH3*, *RDH10*, *CYP1B1*, *CYP26*, and transcript levels of choroidal *RALDH1* were undetectable [10]. Additionally, changes in choroidal *RALDH2* protein concentrations and enzymatic activity in recovering eyes were reflective of the transcriptional changes in choroidal *RALDH2* [14] suggesting that, in response to myopic defocus or recovery from induced myopia, the concentration of choroidal *RALDH2* increases which, in turn, results in increased production of atRA. No *RALDH* activity was detected in the sclera or retina/RPE of control or treated chick eyes, indicating that the choroid is responsible for the majority of atRA synthesized in the chick eye [14].

3.3 Choroidal *RALDH2*⁺ cells: a novel cell type

In chicks and humans, *RALDH2* is synthesized by a population of extravascular choroidal stromal cells, some of which are closely associated with blood vessels (**Figure 1**) [10, 14, 44]. In chicks, *RALDH2*⁺ cells increase in number markedly over 1–7 days of recovery due, in part, to cellular proliferation (**Figure 1F and G**) and become concentrated on the proximal (RPE) side of the choroid [14]. Immunohistochemical analyses of chick choroids indicate that many of *RALDH2*⁺ express pro-collagen type IA (**Figure 1B and C**), similar to activated pericytes (a.k.a. perivascular stromal cells) within the CNS perivascular space [45]. Additionally *RALDH2* is expressed in the chick

Figure 1. Choroidal RALDH2 positive (+) cells are heterogeneous. (A–E) RALDH2+ cells (green) are identified following labeling of 4 day recovering choroids with anti-RALDH2, together with anti-TCR $\gamma\delta$ ($\gamma\delta$), pro-collagen IA (COL1A), or Ia antigen (IA) (red). Double-labeled cells are indicated by arrows. Asterisks in (C) indicate RALDH2+ cells that co-express Col1A. (F) Proliferating RALDH2+ cells were labeled with BrdU and identified with anti-BRDU (red). (G) Percentage of RALDH2+ / BRDU+ cells is $\approx 3\times$ higher in 4 day recovering choroids, suggesting that RALDH2 activity is partially controlled by proliferation of RALDH2+ cells. Bar = 20 μ M in A, 40 μ M in B–F. ***Student's *t* test, *p* < 0.0001. Error bars = SEM. Nuclei in A–F are stained with DAPI. BV, blood vessel.

choroid by a small population of round cells that are positive for the Ia antigen [46, 47], indicating similarities with thymic macrophages/dendritic cells (**Figure 1D and E**), but are negative for the macrophage markers KuL01, MHC-II, and IgY [48]. A sub-population of RALDH2+ cells also express α -smooth muscle actin (α SMA) [10, 14], but are negative for the smooth muscle/myofibroblast proteins, smoothelin, desmin and myocardin. RALDH2+ cells do not co-localize with CD-45 [14], TCR $\delta\gamma$ (**Figure 1A**), CD5, or GRL(2) positive cells [49, 50], indicating they are not of hematopoietic origin. RALDH2+ cells also do not co-localize with neuron-specific beta III tubulin, NOS (pan), or tyrosine hydroxylase, indicating they are not of neuronal origin. Negative results were also obtained using anti-NG2 (a pericyte marker), vimentin, and PECAM-1 (an endothelial marker). Similarly, RALDH2+ cells in the human choroid were negative for the endothelial cell marker, CD31, the pericyte markers, NG2 and CD146, α -smooth muscle actin, the macrophage markers CD68 and LYVE1, IBA1 (microglia) and the pan-neuronal marker PGP9.5 [51]. Unlike results in the chick, some RALDH2+ cells in the human choroid co-localized with vimentin, suggesting a mesenchymal origin [51]. Based on the markers used in these studies, RALDH2+ cells seem to represent an independent cell-population. Studies are in progress using additional markers as well as transcriptome analyses on RALDH2+ cells isolated from chick and human choroids to further classify this new cell-population as this cell type may represent a potential target for therapies to slow or prevent myopia in children.

4. Retinoic acid on scleral proteoglycan synthesis

The retina, choroid and sclera are three possible tissue targets for choroidally generated atRA within the eye. Of these three targets, the sclera is a leading candidate. Based on results using a specific inhibitor of proteoglycan synthesis

Figure 2. Choroidal retinoic acid synthesis and scleral proteoglycan synthesis during recovery from form deprivation myopia. (A) Changes in choroidal all-trans-retinoic acid (atRA) synthesis during recovery from experimental myopia. (B) Changes in scleral proteoglycan synthesis in during recovery from experimental myopia. A from: Mertz and Wallman [9]. B adapted from: Summers and Hollaway [53]. Reproduced with permission © Elsevier.

(*p*-nitrophenyl-beta-D-xylopyranoside), we previously demonstrated that the rate of proteoglycan synthesis in the cartilaginous layer of the chick sclera is largely responsible for visually guided changes in eye size [27, 52]. Moreover, proteoglycan synthesis is rapidly upregulated in the sclera in response to visual form deprivation and is rapidly downregulated in the posterior sclera to levels significantly below those of fellow controls within 12 hours upon restoration of unrestricted vision (=recovery from induced myopia) [53, 54]. Interestingly, the time course of the increase in choroidal atRA synthesis during recovery from induced myopia [9] was remarkably similar to that of the decrease in rate of sclera proteoglycan synthesis observed in the early phase of recovery from induced myopia [53] suggesting a causal relationship between choroidal atRA synthesis and scleral proteoglycan synthesis (**Figure 2**). Moreover, *RALDH2* mRNA pools in the chick choroid were shown to increase and decrease in a manner that inversely correlated with changes in scleral proteoglycan synthesis in recovering eyes [10], suggesting that *RALDH2* gene expression in the choroid regulate choroidal atRA synthesis during visually induced ocular growth. It is well known that atRA is a potent inhibitor of proteoglycan biosynthesis by chondrocytes [55, 56] and that it facilitates cartilage catabolism through the increased synthesis of matrix-degrading enzymes [55, 57, 58]. atRA inhibits scleral proteoglycan synthesis in a dose-dependent manner with an IC_{50} of 8×10^{-9} M, which is similar to the measured endogenous levels of atRA in choroid organ cultures (4×10^{-9} to 7×10^{-9} M) [10]. At this concentration, atRA would be able to regulate scleral growth matrix remodeling through the stimulation or repression of transcription factors, extracellular matrix constituents, and MMPs or TIMPs.

5. Identification of apolipoprotein A-1 as a retinoic acid binding protein

Due to its hydrophobicity, atRA cannot diffuse freely in the hydrophilic extracellular microenvironment. Therefore, the requirement for carrier proteins capable of forming a soluble complex with atRA and transporting atRA to target cells is necessary to achieve high efficiency and specificity while avoiding toxicity associated with random diffusion. Mertz and Wallman [9] and our lab [15] identified a secreted protein of $Mr = 27,000$ that was the major atRA binding protein present in choroid and sclera conditioned medium. This $Mr 27,000$ protein did not correspond in size to any of the previously identified atRA binding proteins [59]. We subsequently determined that the $Mr 27,000$ protein was apolipoprotein a-1 (ApoA-1) [15] (**Figure 3**).

We have also shown that choroidal expression of ApoA-1 is transcriptionally regulated by atRA, and choroidal ApoA-1 mRNA and protein synthesis are upregulated during recovery from induced myopia, suggesting the presence of a regulatory feedback mechanism to regulate atRA transport and activity [15]. We postulate that ApoA-1 functions to transport atRA from its site of synthesis by RALDH2+ cells in the proximal choroid to the sclera for the regulation of scleral ECM remodeling. This idea is supported by our observation that the chick sclera (which is avascular) also releases significant amounts of ApoA-1 into culture medium, despite undetectable *de novo* protein synthesis [15]. These data provide further evidence that choroidally derived ApoA-1 accumulates in the sclera, presumably as a consequence of retinoid transport.

Figure 3. ApoA-1 is a specific atRA-binding protein. (A) Retinoic acid (atRA) lacks intrinsic fluorescence (not shown), but can quench intrinsic protein fluorescence excited at 290 nm due to energy transfer from tryptophan residues on ApoA-1. Increasing atRA concentrations cause decreased fluorescence emission following excitation at 290 nm. (B) Titration of ApoA-1 with various retinoids by measuring quenching of protein fluorescence (emission = 340 nm). Significant quenching of protein fluorescence was observed only for atRA, indicating that ApoA-1 is a specific atRA-binding protein. atRA, all-trans-retinoic acid; RAL, all-trans-retinaldehyde; 9-cis, 9-cis-retinoic acid; 13-cis, 13-cis-retinoic acid. This research was originally published in the *Journal of Biological Chemistry* [70]. © The American Society for Biochemistry and Molecular Biology.

6. Role of retinoic acid in postnatal ocular growth

To elucidate the role of atRA in the regulation of postnatal ocular growth, several studies have been carried out in which either atRA or non-specific atRA synthesis inhibitors (i.e., citral, disulfiram) were administered either systemically or locally in several animals undergoing visually induced changes in eye growth [12, 60, 61]. Results of studies using chicks and mammals to examine the role of atRA in emmetropization, myopia development and postnatal ocular growth are difficult to interpret due to species differences in the processes of scleral remodeling and in the mechanisms by which ocular length and refraction are modulated by visual stimuli [62]. Moreover, these studies are further complicated by the multiple targets of atRA within the eye and pleiotropic cellular responses to retinoid signaling [63]. The mammalian sclera consists of a single fibrous layer that undergoes scleral thinning, and increased distensibility during periods of ocular elongation and myopia development. Scleral thinning during myopia development in mammals is the consequence of decreased sulfated glycosaminoglycan and collagen synthesis [11, 64, 65]. In contrast, the chick sclera consists of both cartilaginous and fibrous scleral layers. Ocular elongation during induced myopia in chicks is the result of growth of the cartilaginous sclera, with increases in sulfated glycosaminoglycan synthesis, increased protein synthesis, and increased total scleral mass [27, 66–68]. In chicks, increased choroidal synthesis of atRA during recovery from form deprivation myopia results in inhibition of scleral proteoglycan synthesis and slowing of the rate of ocular elongation. In primates [11] and guinea pigs [12], choroidal atRA synthesis is increased in treated eyes following induced myopia, a condition that is also associated with decreased proteoglycan synthesis in the posterior sclera but, in contrast to chicks, results in increased ocular elongation and myopia due to weakening of the fibrous sclera and localized ectasia at the posterior ocular pole. Considering the negative effect of atRA on scleral proteoglycan synthesis in animals containing either a single fibrous sclera (i.e., guinea pigs, primates) as well as chicks that contain both cartilaginous and fibrous scleral layers [9, 11], choroidally derived atRA represents a mechanism to regulate ocular length and refraction common to multiple species.

Furthermore, interpretation of experiments in which atRA agonists and atRA synthesis inhibitors are delivered either systemically or intraocularly is complicated by the widespread multicellular effects of atRA. Eye growth is increased following dietary delivery of atRA to chicks and is decreased after oral delivery of citral, a non-specific inhibitor of atRA synthesis [61]. Similarly, intraocular delivery of the non-specific atRA synthesis inhibitor, disulfiram, inhibited the development of form-deprivation myopia in chicks [60], a result generally opposite of what would be predicated if atRA acted to inhibit ocular elongation in chicks. It is likely that untargeted administration of atRA or use of non-specific atRA synthesis inhibitors that also inhibit other aldehyde dehydrogenases lead to multicellular effects that may differ from those mediated by endogenous atRA. We have recently developed a small molecule inhibitor, dichloro-all-*trans*-retinone (DAR) that is an irreversible inhibitor of RALDH1, 2, and 3 that effectively inhibits RALDH1, 2, and 3 in the nanomolar range but has no inhibitory activity against mitochondrial ALDH2 [69]. It is hoped that DAR, or similar compounds can be used to modulate endogenous concentrations of atRA through specific inhibition of the RALDH isoenzymes within the eye for future experimental and clinical studies to elucidate the role of atRA on postnatal ocular growth and myopia development.

7. Conclusions

Although the cause of myopia in humans is complex, clinical and experimental studies indicate that failure of the emmetropization process often leads to the development of myopia. It has been well-established that visually induced changes in ocular length are the result of altered extracellular matrix remodeling of the scleral shell. However no therapeutic targets have been identified and no pharmaceutical or optometric approaches have proven effective for the treatment of high myopia. The increasing prevalence of myopia and earlier age of onset emphasize the need for the development of an effective therapy. The identification of choroidal atRA, RALDH2, and the choroidal cells responsible for atRA synthesis, may provide new targets for the development of effective myopia therapies. Moreover the development of small molecule inhibitors specifically targeting RALDH2 would greatly expand our basic understanding atRA's role in postnatal growth and development as well as provide potential new therapies to slow or prevent the progression of myopia.

Acknowledgements

I would like to thank Dr. Tim Mather (Department of Biochemistry, University of Oklahoma Health Science Center) for his helpful suggestions. The work from the author's laboratory mentioned in this review was supported by grants from the National Institutes of Health (R01 EY009391 and P20 GM103640).

Conflict of interest

The author certifies that she has no conflicts of interest and no affiliations with or involvement in any organization or entity with any financial interest.

Author details

Jody A. Summers
Department of Cell Biology, University of Oklahoma Health Science Center,
Oklahoma City, OK, USA

*Address all correspondence to: jody-summers@ouhsc.edu

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Niederreither K, Dolle P. Retinoic acid in development: Towards an integrated view. *Nature Reviews Genetics*. 2008;**9**(7):541-553
- [2] Rhinn M, Dolle P. Retinoic acid signalling during development. *Development*. 2012;**139**(5):843-858
- [3] Luo T, Wagner E, Crandall JE, Drager UC. A retinoic-acid critical period in the early postnatal mouse brain. *Biological Psychiatry*. 2004;**56**(12):971-980
- [4] Wang TW, Zhang H, Parent JM. Retinoic acid regulates postnatal neurogenesis in the murine subventricular zone-olfactory bulb pathway. *Development*. 2005;**132**(12):2721-2732
- [5] Hagglund M, Berghard A, Strotmann J, Bohm S. Retinoic acid receptor-dependent survival of olfactory sensory neurons in postnatal and adult mice. *The Journal of Neuroscience*. 2006;**26**(12):3281-3291
- [6] Wu JW, Wang RY, Guo QS, Xu C. Expression of the retinoic acid-metabolizing enzymes RALDH2 and CYP26b1 during mouse postnatal testis development. *Asian Journal of Andrology*. 2008;**10**(4):569-576
- [7] Williams JA, Kondo N, Okabe T, Takeshita N, Pilchak DM, Koyama E, et al. Retinoic acid receptors are required for skeletal growth, matrix homeostasis and growth plate function in postnatal mouse. *Developmental Biology*. 2009;**328**(2):315-327
- [8] McCaffery P, Mey J, Drager UC. Light-mediated retinoic acid production. *Proceedings of the National Academy of Sciences of the United States of America*. 1996;**93**(22):12570-12574
- [9] Mertz JR, Wallman J. Choroidal retinoic acid synthesis: A possible mediator between refractive error and compensatory eye growth. *Experimental Eye Research*. 2000;**70**(4):519-527
- [10] Rada JA, Hollaway LY, Li N, Napoli J. Identification of RALDH2 as a visually regulated retinoic acid synthesizing enzyme in the Chick choroid. *Investigative Ophthalmology & Visual Science*. 2012;**53**(3):1649-1662
- [11] Troilo D, Nickla DL, Mertz JR, Summers Rada JA. Change in the synthesis rates of ocular retinoic acid and scleral glycosaminoglycan during experimentally altered eye growth in marmosets. *Investigative Ophthalmology & Visual Science*. 2006;**47**(5):1768-1777
- [12] McFadden SA, Howlett MH, Mertz JR. Retinoic acid signals the direction of ocular elongation in the Guinea pig eye. *Vision Research*. 2004;**44**(7):643-653
- [13] Seko Y, Shimizu M, Tokoro T. Retinoic acid increases in the retina of the chick with form deprivation myopia. *Ophthalmic Research*. 1998;**30**(6):361-367
- [14] Harper AR, Wang X, Moiseyev G, Ma JX, Summers JA. Postnatal Chick choroids exhibit increased retinaldehyde dehydrogenase activity during recovery from form deprivation induced myopia. *Investigative Ophthalmology & Visual Science*. 2016;**57**(11):4886-4897
- [15] Summers JA, Harper AR, Feasley CL, Van-Der-Wel H, Byrum JN, Hermann M, et al. Identification of apolipoprotein A-I as a retinoic acid-binding protein in the eye. *The Journal of Biological Chemistry*. 2016;**291**(36):18991-19005
- [16] Blomhoff R, Blomhoff HK. Overview of retinoid metabolism and function. *Journal of Neurobiology*. 2006;**66**(7):606-630

- [17] Napoli JL. Physiological insights into all-trans-retinoic acid biosynthesis. *Biochimica et Biophysica Acta*. 2012;**1821**(1):152-167
- [18] Chambers D, Wilson L, Maden M, Lumsden A. RALDH-independent generation of retinoic acid during vertebrate embryogenesis by CYP1B1. *Development*. 2007;**134**(7):1369-1383
- [19] Wang C, Kane MA, Napoli JL. Multiple retinol and retinal dehydrogenases catalyze all-trans-retinoic acid biosynthesis in astrocytes. *The Journal of Biological Chemistry*. 2011;**286**(8):6542-6553
- [20] Balmer JE, Blomhoff R. Gene expression regulation by retinoic acid. *Journal of Lipid Research*. 2002;**43**(11):1773-1808
- [21] O'Leary DJ, Millodot M. Eyelid closure causes myopia in humans. *Experientia*. 1979;**35**(11):1478-1479
- [22] Rabin J, Van Sluyters RC, Malach R. Emmetropization: A vision-dependent phenomenon. *Investigative Ophthalmology & Visual Science*. 1981;**20**(4):561-564
- [23] Wallman J, Turkel J, Trachtman J. Extreme myopia produced by modest change in early visual experience. *Science*. 1978;**201**(4362):1249-1251
- [24] Wallman J, Adams JI. Developmental aspects of experimental myopia in chicks: Susceptibility, recovery and relation to emmetropization. *Vision Research*. 1987;**27**(7):1139-1163
- [25] Zhu X, Park TW, Winawer J, Wallman J. In a matter of minutes, the eye can know which way to grow. *Investigative Ophthalmology & Visual Science*. 2005;**46**(7):2238-2241
- [26] Rada JA, Thoft RA, Hassell JR. Increased aggrecan (cartilage proteoglycan) production in the sclera of myopic chicks. *Developmental Biology*. 1991;**147**(2):303-312
- [27] Rada JA, Johnson JM, Achen VR, Rada KG. Inhibition of scleral proteoglycan synthesis blocks deprivation-induced axial elongation in chicks. *Experimental Eye Research*. 2002;**74**(2):205-215
- [28] Schaeffel F, Glasser A, Howland HC. Accommodation, refractive error and eye growth in chickens. *Vision Research*. 1988;**28**(5):639-657
- [29] Hung LF, Crawford ML, Smith EL. Spectacle lenses alter eye growth and the refractive status of young monkeys. *Nature Medicine*. 1995;**1**(8):761-765
- [30] Smith EL 3rd, Hung LF. The role of optical defocus in regulating refractive development in infant monkeys. *Vision Research*. 1999;**39**(8):1415-1435
- [31] Shaikh AW, Siegwart JT Jr, Norton TT. Effect of interrupted lens wear on compensation for a minus lens in tree shrews. *Optometry and Vision Science*. 1999;**76**(5):308-315
- [32] Hodos W, Kuenzel WJ. Retinal-image degradation produces ocular enlargement in chicks. *Investigative Ophthalmology & Visual Science*. 1984;**25**(6):652-659
- [33] Wallman J, Gottlieb MD, Rajaram V, Fugate-Wentzek LA. Local retinal regions control local eye growth and myopia. *Science*. 1987;**237**(4810):73-77
- [34] Raviola E, Wiesel TN. An animal model of myopia. *The New England Journal of Medicine*. 1985;**312**(25):1609-1615
- [35] Norton TT, Essinger JA, McBrien NA. Lid-suture myopia in tree shrews with retinal ganglion cell blockade. *Visual Neuroscience*. 1994;**11**(1):143-153

- [36] Troilo D, Gottlieb MD, Wallman J. Visual deprivation causes myopia in chicks with optic nerve section. *Current Eye Research*. 1987;**6**(8):993-999
- [37] Summers JA. The choroid as a sclera growth regulator. *Experimental Eye Research*. 2013;**114**:120-127
- [38] Wallman J. Retinal influences on sclera underlie visual deprivation myopia. *CIBA Foundation Symposium*. 1990;**155**:126-134 discussion 35-41
- [39] Pendrak K, Nguyen T, Lin T, Capehart C, Zhu X, Stone RA. Retinal dopamine in the recovery from experimental myopia. *Current Eye Research*. 1997;**16**(2):152-157
- [40] Stone RA, Laties AM, Raviola E, Wiesel TN. Increase in retinal vasoactive intestinal polypeptide after eyelid fusion in primates. *Proceedings of the National Academy of Sciences of the United States of America*. 1988;**85**(1):257-260
- [41] Fischer AJ, McGuire JJ, Schaeffel F, Stell WK. Light- and focus-dependent expression of the transcription factor ZENK in the chick retina. *Nature Neuroscience*. 1999;**2**(8):706-712
- [42] Stone RA, Lin T, Laties AM, Iuvone PM. Retinal dopamine and form-deprivation myopia. *Proceedings of the National Academy of Sciences of the United States of America*. 1989;**86**(2):704-706
- [43] Simon P, Feldkaemper M, Bitzer M, Ohngemach S, Schaeffel F. Early transcriptional changes of retinal and choroidal TGFbeta-2, RALDH-2, and ZENK following imposed positive and negative defocus in chickens. *Molecular Vision*. 2004;**10**:588-597
- [44] Harper AR, Wiechmann AF, Moiseyev G, Ma JX, Summers JA. Identification of active retinaldehyde dehydrogenase isoforms in the postnatal human eye. *PLoS ONE*. 2015;**10**(3):e0122008
- [45] Kelly KK, MacPherson AM, Grewal H, Strnad F, Jones JW, Yu J, et al. Col1a1+ perivascular cells in the brain are a source of retinoic acid following stroke. *BMC Neuroscience*. 2016;**17**(1):49
- [46] Guillemot FP, Oliver PD, Peault BM, Le Douarin NM. Cells expressing Ia antigens in the avian thymus. *The Journal of Experimental Medicine*. 1984;**160**(6):1803-1819
- [47] Nagy N, Olah I. Experimental evidence for the ectodermal origin of the epithelial anlage of the chicken bursa of Fabricius. *Development*. 2010;**137**(18):3019-3023
- [48] Summers Rada JA, Hamid S, Harper AR, Forest-Smith L, Wren J. Isolation and transcriptome analyses of choroidal retinaldehyde dehydrogenase-2 (RALDH2) expressing cells. *Investigative Ophthalmology & Visual Science*. 2017;**58**(8). Abs. No. 1100
- [49] Thomas JL, Pourquie O, Coltey M, Vaigot P, Le Douarin NM. Identification in the chicken of GRL1 and GRL2: Two granule proteins expressed on the surface of activated leukocytes. *Experimental Cell Research*. 1993;**204**(1):156-166
- [50] Bodi I, Nagy N, Sinka L, Igyarto BZ, Olah I. Novel monoclonal antibodies recognise Guinea fowl thrombocytes. *Acta Veterinaria Hungarica*. 2009;**57**(2):239-246
- [51] Schroedl F, Kaser-Eichberger A, Trost A, Runge C, Bruckner D, Bogner B, et al. Morphological classification of RALDH2-positive cells in the human choroid. *Investigative Ophthalmology & Visual Science*. 2018;**59**(9). (Abs. # 308)
- [52] Rada JA, Matthews AL, Brenza H. Regional proteoglycan synthesis in

the sclera of experimentally myopic chicks. *Experimental Eye Research*. 1994;**59**(6):747-760

[53] Summers Rada JA, Hollaway LR. Regulation of the biphasic decline in scleral proteoglycan synthesis during the recovery from induced myopia. *Experimental Eye Research*. 2011;**92**(5):394-400

[54] Rada JA, McFarland AL, Cornuet PK, Hassell JR. Proteoglycan synthesis by scleral chondrocytes is modulated by a vision dependent mechanism. *Current Eye Research*. 1992;**11**(8):767-782

[55] Campbell MA, Handley CJ. The effect of retinoic acid on proteoglycan turnover in bovine articular cartilage cultures. *Archives of Biochemistry and Biophysics*. 1987;**258**(1):143-155

[56] Horton WE, Yamada Y, Hassell JR. Retinoic acid rapidly reduces cartilage matrix synthesis by altering gene transcription in chondrocytes. *Developmental Biology*. 1987;**123**(2):508-516

[57] Sugimoto K, Takahashi M, Yamamoto Y, Shimada K, Tanzawa K. Identification of aggrecanase activity in medium of cartilage culture. *Journal of Biochemistry*. 1999;**126**(2):449-455

[58] Nie D, Ishikawa Y, Yoshimori T, Wuthier RE, Wu LN. Retinoic acid treatment elevates matrix metalloproteinase-2 protein and mRNA levels in avian growth plate chondrocyte cultures. *Journal of Cellular Biochemistry*. 1998;**68**(1):90-99

[59] Noy N. Retinoid-binding proteins: Mediators of retinoid action. *The Biochemical Journal*. 2000;**348**(Pt 3):481-495

[60] Bitzer M, Feldkaemper M, Schaeffel F. Visually induced changes in components of the retinoic acid system in fundal layers of the

chick. *Experimental Eye Research*. 2000;**70**(1):97-106

[61] McFadden SA, Howlett MH, Mertz JR, Wallman J. Acute effects of dietary retinoic acid on ocular components in the growing chick. *Experimental Eye Research*. 2006;**83**(4):949-961

[62] Rada JA, Shelton S, Norton TT. The sclera and myopia. *Experimental Eye Research*. 2006;**82**(2):185-200

[63] Cvekl A, Wang WL. Retinoic acid signaling in mammalian eye development. *Experimental Eye Research*. 2009;**89**(3):280-291

[64] Siegwart JT Jr, Norton TT. Regulation of the mechanical properties of tree shrew sclera by the visual environment. *Vision Research*. 1999;**39**(2):387-407

[65] Norton TT, Rada JA. Reduced extracellular matrix in mammalian sclera with induced myopia. *Vision Research*. 1995;**35**(9):1271-1281

[66] Christensen AM, Wallman J. Evidence that increased scleral growth underlies visual deprivation myopia in chicks. *Investigative Ophthalmology & Visual Science*. 1991;**32**(7):2143-2150

[67] McBrien NA, Moghaddam HO, Reeder AP, Moules S. Structural and biochemical changes in the sclera of experimentally myopic eyes. *Biochemical Society Transactions*. 1991;**19**(4):861-865

[68] Gentle A, Truong HT, McBrien NA. Glycosaminoglycan synthesis in the separate layers of the chick sclera during myopic eye growth: Comparison with mammals. *Current Eye Research*. 2001;**23**(3):179-184

[69] Harper AR, Le AT, Mather T, Burgett A, Berry W, Summers JA. Design, synthesis, and ex vivo evaluation of a selective inhibitor for

retinaldehyde dehydrogenase enzymes.
Bioorganic & Medicinal Chemistry.
2018;**26**(22):5766-5779

[70] Summers JA et al. Identification
of apolipoprotein A-1 as a retinoic
acid binding protein in the eye. The
Journal of Biological Chemistry.
2016;**291**:18991-19005