

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Orexin System and Avian Muscle Mitochondria

Kentu Lassiter and Sami Dridi

Abstract

In mammals, orexin A and B (also known as hypocretin 1 and 2) are two orexigenic peptides produced primarily by the lateral hypothalamus that signal through two G-protein-coupled receptors, orexin receptors 1/2, and have been implicated in the regulation of several physiological processes. However, the physiological roles of orexin are not well defined in avian (non-mammalian vertebrate) species. Recently, we made a breakthrough by identifying that orexin and its related receptors 1/2 (ORXR1/2) are expressed in avian muscle tissue and cell line, and appears to be a secretory protein. Functional in vitro studies showed that orexin A and B differentially regulated expression of the orexin system, suggesting that orexins might have autocrine, paracrine, and/or endocrine roles. Administration of recombinant orexin modulated mitochondrial biogenesis, dynamics, function, and bioenergetics. In this chapter, we include a brief overview of the (patho) physiological role of orexin, comparative findings between mammalian and avian orexin, and in-depth analysis of orexin's action on avian muscle mitochondria.

Keywords: orexin system, muscle, avian species, mitochondria, gene expression

1. Introduction

Orexin/hypocretin was originally identified by two different groups and reported in 1998 as an orexigenic feeding-related neuropeptide mainly produced in the rat hypothalamus [1, 2]. Numerous subsequent studies conducted in mammals have shown that orexin and its receptors regulate several physiological processes including food and water intake [3], control of wakefulness [4], circadian clock [5], energy and glucose homeostasis [6–8], lipid metabolism [9, 10], heart rate and blood pressure [11, 12], and neuroendocrine response to stress [13]. Despite these advancements in understanding the orexin system, studies investigating its distribution and function in avian (non-mammalian vertebrate) species are very limited and merit more consideration and further in-depth explorations. Understanding orexin distribution and unraveling its function in avian tissues, particularly in broiler (meat-type chickens) skeletal muscle is of particular importance not only to the poultry industry, but also to the biomedical field.

The poultry (meat and egg) industry supports the livelihoods and food security of billions of people worldwide with an average annual production of 118 million metric tons of meat and 1360 billion eggs in 2017 [14]. This success was mainly achieved by intensive genetic selection for important economic and agricultural traits such as high growth rate, improved feed efficiency (conversion of feed to muscle mass), and increased muscle mass [15]. Today, indeed, modern broilers are

marketed in about half the time and at about twice the body weight compared to 50–60 years ago [16]. This fast growth is largely allocated to the breast muscle [17].

Chickens are naturally hyperglycemic compared with mammals [18], insulin-resistant [19], lack the glucose transporter GLUT4 [20], lack functional brown adipose tissue [21], and are prone to obesity [22], and thereby they represent a highly relevant animal model for biomedical researches.

The proper function of mitochondria is invariably related to muscle growth since these organelles produce a vast majority (~90%) of the ATP needed for tissue growth and maintenance of energy metabolism homeostasis. Previous studies in broiler chickens have shown that mitochondrial perturbations are directly related to a decrease in feed efficiency. Additionally, the absence of orexin's effect on feed intake in chickens [23] suggest that this neuropeptide may be more involved in other peripheral metabolic and physiological processes rather than food/water intake and wakefulness that is seen in mammals. Therefore, if orexin and its receptors are shown to be present in skeletal muscle tissue of avian species, they may be exerting control of energy metabolism and muscle growth (i.e., myogenesis, insulin sensitivity, and glucose transport) in the cell in part by regulating mitochondrial dynamics. The role of orexin in skeletal muscle function could be of interest not only in improving health and feed efficiency of agricultural animals, but also in molecular medicine for pathophysiological understanding and therapeutic perspectives.

2. Identification of orexin and its receptors

Orexin, which regulates wakefulness, energy homeostasis, and appetite/feeding behavior based on nutritional status, is a neuropeptide that was originally discovered in the hypothalamus of rats by two different research groups in 1998. De Lecea's group isolated cDNAs selectively expressed within the hypothalamus and found that two peptides showed high amino acid sequence homology with secretin (the gut peptide hormone), therefore they named the two peptides hypocretin 1 and 2 [1]. Sakurai's group, on the other hand, used reverse pharmacology to identify ligands of orphan G-protein-coupled receptors. Orphan receptors are those whose ligand and physiological actions are unknown [24]. They identified a novel family of neuropeptides that induced feeding behavior, so they named them orexin A and B [2]. The term orexin originates from the Greek word "orexis," meaning appetite. Both orexin A and B are formed by proteolytic cleavage of the precursor prepro-orexin [2]. When initially discovered in rats, the precursor peptide prepro-orexin was shown to be a 130-residue polypeptide from which the mature peptides ORX-A and ORX-B were formed, with ORX-A containing 33 amino acids and a molecular weight 3.562 kDa and ORX-B containing 28 amino acids and a molecular weight of 2.937 kDa [2]. When comparing the two peptides, ORX-B was shown to be 56% identical in amino acid sequence to ORX-A. However, when comparing mammalian species (human, rat, mouse, pig, and cow), the sequence and structure of both peptides is highly conserved [2]. A number of studies have also shown that the structures of ORX-A and ORX-B in chicken and certain types of fish are conserved when compared to their mammalian counterparts [25–28].

ORX-A and ORX-B signal through the G-protein-coupled receptors orexin receptor 1 (ORXR1) and orexin receptor 2 (ORXR2). These two ubiquitously expressed receptors were first identified in human brain tissue through expressed sequence tags combined with database searching using tBLASTn [2, 29]. In humans it has been shown that the amino acid sequence for ORXR1 and ORXR2 is more than

60% identical, making them more similar to each other than to other G-protein-coupled receptors [2]. The same study also showed that both receptors are highly conserved between humans and rats, with the sequence identity being greater than 90% for both. The two orexin peptides have different binding affinities for the two orexin receptors. ORX-A is able to bind to both receptors but has a higher affinity for ORXR1, while ORX-B binds to ORXR2 with the same affinity as ORX-A [28]. Several studies have indicated that the binding of orexins to orexin receptors activates multiple G-proteins. In studies conducted using humans [30, 31] and rats [32], it was shown that the binding of ORXR2 activates Gi, Gs, Go, and Gq proteins in adrenal cortical tissue. It appears that the responses to orexin receptor signaling are highly diverse. The activation of the various G-proteins can lead to a variety of cellular responses such as the regulation of protein/lipid kinases [33, 34]. In the case of orexin stimulation, activation of G-proteins can lead to the excitation of neurons that affect the regulation of ion channels, the activation of signaling cascades that regulate the activity of adenylyl cyclase and phospholipases, and activation of cell death pathways [35, 36].

3. Orexin system in avian species

Significantly fewer studies concerning the orexin system have been conducted in avian species when compared to mammals. Chicken prepro-orexin was first cloned, sequenced, and characterized in 2002 [37]. In that study, chicken orexin cDNA was shown to be expressed in the periventricular and lateral hypothalamic areas and consisting of 658 bp that encode 148 amino acids. Also, chicken ORX-A and ORX-B are evolutionary conserved with their mammalian counterparts, showing ~85 and 65% similarity at the amino acid level [37]. Characterization of the chicken orexin receptor shows that its cDNA has a length of 1869 bp that encode 501 amino acids, which corresponds to mammalian ORXR2 with an 80% homology [37]. Studies looking at tissue distribution of orexin and orexin receptors in chickens show that the peptides are expressed in the brain [37–40], pituitary gland, adrenal gland, testis and ovary [38], and the stomach and intestine [41].

Orexin does not appear to elicit the same responses in birds as it does in mammals. One of the most noted actions that centrally administered orexin has in mammals is that it stimulates feeding/food intake [2, 3, 28, 42]. However, central administration of ORX-A or ORX-B did not stimulate feed intake in neonatal broiler and layer chicks [23, 43] or adult pigeons [44]. Studies examining mRNA expression of prepro-orexin in the hypothalamus of chicken [37] and quail [45] following 24 h fasting showed no increase in expression, providing further evidence for the lack of a stimulatory effect on feeding behavior in birds. Another feeding behavior study did show an increase in prepro-orexin mRNA [46], but this was measured after 48 h fasting, which would be an extreme fasting condition for broiler chickens.

As stated previously, another hallmark of orexin function in mammals is its effects on the regulation of sleep/wakefulness, where a dysfunction in the orexin system is associated with the sleep condition narcolepsy [4]. Studies investigating the effects of orexin on arousal in birds have been conducted with mixed results. It has been concluded that either hypothalamic orexin does not play a role in arousal of the sleep/wake cycle [39], or that only ORX-A in conjunction with the enzyme monoamine oxidase-A (MAO-A) increases arousal in layer chicks only and not broiler chicks [43, 47]. Multiple studies investigating orexin in avian species theorize that the peptide appears to be more involved in the regulation of energy metabolism than feed intake and sleep/wake cycles [39, 40, 48].

4. Orexin system in avian muscle

Studies investigating the orexin-producing neurons in the rat/mouse brain report that central administration of the neuropeptide facilitates changes in




Figure 1. Prepro-orexin and its related receptors are expressed in broiler chicken muscle (a, b) and QM7 cell line (c-f). (a) RT-PCR. Total RNA (1 µg) was reverse transcribed and subjected to RT-PCR. Brain, testis, and ovary were used as positive controls. (b) Western blot. 70 µg total protein extracted from each tissue were electrophoresed and blotted onto polyvinylidene difluoride membrane. Prepro-orexin (ORX) and orexin receptor 1 (ORXR1) expression was detected by immunoblot using rabbit anti-mouse ORX and rabbit anti-rat ORXR1 antibodies. Hypothalamus and brain were used as positive controls. The figure is a representative picture from one animal. (c) RT-PCR. Total RNA (1 µg) was isolated from chicken brain and QM7 cells and subjected to RT-PCR. (d) Western blot. Total protein extracted from the cells were electrophoresed and blotted onto polyvinylidene difluoride membrane. Prepro-orexin (ORX) and ORXR1/2 expression was detected by immunoblot using rabbit anti-mouse ORX and rabbit anti-rat ORXR1/2 antibodies. Brain was used as positive control. (e) Northern blot. Total RNA (10 µg) was separated by agarose gel electrophoresis and transferred to a nylon membrane and hybridized with specific biotin-labeled DNA probe toward chicken ORX, and 18S. Hybridization signals were detected by FluorChem M MultiFluor system. (f) Immunofluorescence staining. Intracellular ORX system distribution visualized by fluorescent microscope in the presence of a secondary antibody conjugated with Alexa Fluor 488 (green) and DAPI (blue).

skeletal muscle tone [49, 50] in addition to increasing glucose metabolism and insulin sensitivity of skeletal muscle [51]. This indicates that orexin signaling in the central nervous system regulates muscle glucose metabolism by activating muscle sympathetic nerves. Even though the orexin system has been identified in the peripheral tissues of a variety of vertebrate species [52], very few studies have been conducted to determine whether orexin and its related receptors are expressed in vertebrate skeletal muscle and if so, what effects its presence may have on vertebrate muscle function and physiology. Review of the current literature shows that to date orexin has only been identified as being expressed in the heart muscle of zebrafish [53] and skeletal muscle of goldfish, chicken, and quail [54–56].

We have conducted studies using RT-PCR and Western blot analysis to identify both orexin and its related receptors as being expressed in broiler chicken muscle (**Figure 1a, b**). In addition, RT-PCR, Western blot analysis, Northern blot analysis, and immunofluorescence have been used to illustrate that orexin and its receptors are also expressed in the cytoplasmic compartment of a spontaneously immortalized quail muscle (QM7) cell line (**Figure 1c–f**). In humans, scientific evidence indicates that orexin is a secretory peptide due to its presence in the blood circulation. Additional support for orexin being secreted is given by the first 33 amino acids of human prepro-orexin containing a hydrophobic core followed by residues with small polar side chains, which are characteristics of a secretory signal sequence [57]. Cell culture experiments using the QM7 cell line have been carried out in order to determine whether orexin is also secreted in avian muscle. When QM7 cells are incubated in serum-free growth media in the presence of recombinant human orexin B (rORX-B), an increase in the level of prepro-orexin in the growth media is evident as seen in **Figure 2a**. Further support for the secretion of prepro-orexin is illustrated in **Figure 2b** where the levels of prepro-orexin in the serum-free growth media accumulate over a 72 h period, and in **Figure 2c** where application of the anti-secretory compound brefeldin A causes a buildup of prepro-orexin in the QM7 cell lysate and the subsequent absence of the peptide in the growth media. The expression and secretion of prepro-orexin from avian muscle cells suggests that avian orexin could be a myokine that probably functions in autocrine, paracrine, and/or endocrine roles.

Subsequent experiments were conducted in order to determine whether the orexin system is capable of self-regulation. The effects of 10 and 100 nM of recombinant human orexin A (rORX-A) and rORX-B on mRNA and protein expression of ORX and its related receptors ORXR1 and ORXR2 in QM7 cells are shown in **Figure 2d–g**. A 24 h treatment with either 10 or 100 nM rORX-A upregulated ORX and ORXR1, but not ORXR2 mRNA expression (**Figure 2f**). Treating cells with rORX-B downregulated ORX and ORXR2 and increased ORXR1 mRNA levels (**Figure 2g**). Protein expression levels of ORX, ORXR1, and ORXR2 showed the same patterns as their corresponding genes (**Figure 2d, e**) with the effects on expression levels appearing to be dose-dependent. The ability of rORX-A and rORX-B to differentially regulate gene and protein expression of the orexin system supports the idea that orexins function in an autocrine, paracrine, and/or endocrine role in avian muscle. Although the underlying mechanisms behind the differential regulation of the orexin system is still unknown, the divergent effects of rORX-A and rORX-B on orexin system expression might be related to their structure (presence of disulfide bonds in orexin A and not in orexin B) and their different binding affinity to ORXR2, since they had similar effects on ORXR1 expression.




Figure 2. Secretion of orexin by QM7 cells (a–c) and effect of orexin treatment on orexin system expression in QM7 cells (d–g). (a) Cell monolayers were incubated in serum-free medium with rORX-B (100 nM) for up to 72 h prior to Western blotting. (b) Percent change in ORX levels when compared to control (2 h post-treatment). (c) Cells were incubated in serum-free medium with or without brefeldin A (0.3 µg/ml) for 12 h. Medium and/or cell lysates were subjected to immunoblot analysis using an orexin antibody. (d–g) Cells were treated with 0 (control), 10, or 100 nM of recombinant orexin A and B for 24 h. Total protein and RNA were isolated and relative expression of ORX and ORXR1/2 was determined. Protein levels were measured by Western blot analysis (d, e). mRNA expression was measured by QPCR using $2^{-\Delta\Delta C_t}$ method (f, g). Data are expressed as means \pm SE (n = 6). Significant difference between orexin-treated and control cells ($P < 0.05$).

4.1 The orexin system differentially regulates mitochondrial-related genes and mitochondrial bioenergetics in avian muscle cells

In mammals, orexin has been shown to induce differentiation of brown adipose tissue (BAT), subsequently leading to thermogenesis [58, 59]. One of the effects orexin has in this process is the regulation of genes involved in mitochondrial biogenesis. The treatment of mouse preadipocytes with ORX-A revealed a number of changes in

mitochondrial dynamics, including up-regulation of the expression of a number of genes involved in mitochondrial biogenesis (i.e., PGC-1 α , PGC-1 β , PPAR γ 1, and UCP1) [58]. These findings were further supported when subsequent immunofluorescence staining revealed an increase in mitochondrial abundance of the treated cells. Studies using other cell types treated with ORX-A have also shown effects on mitochondrial function. Human neuroblastoma cells treated with ORX-A had increased mitochondrial membrane potential [60]. Additionally, in studies using human hepatoma cells [61] and human embryonic kidney cells [62], treatment with ORX-A resulted in increased ATP production that shifted from glycolysis in the cytoplasm to oxidative phosphorylation in the mitochondria. Taken together, these studies indicate that orexin is able to enhance mitochondrial function, biogenesis, and ATP production in mammalian vertebrates.

Similar experiments were carried out by treating QM7 cells with recombinant human orexin, as previously described, in order to observe the differential effects on mitochondrial-related genes, transcriptional regulators, and bioenergetics. **Figure 3** illustrates how orexin causes differential expression of mitochondrial genes. rORX-A had no effect on av-UCP mRNA abundance (**Figure 3b**), but it downregulated the expression of av-ANT (mRNA and protein levels), Ski, and NRF-1 in a dose-dependent manner (**Figure 3a, c, d**). rORX-B, however, downregulated the expression of av-UCP and increased the expression of Ski and NRF-1 without altering the expression of av-ANT (**Figure 3a–d**). The mitochondrial transcriptional regulators related to these genes were also differentially regulated following treatment with recombinant orexins as seen in **Figure 3e–h**. Both doses of rORX-A caused a significant downregulation of PPAR δ and FoxO-1 expression (**Figure 3g, h**). A high dose of rORX-A significantly downregulated the expression of PGC-1 β , and both doses did not alter PGC-1 α mRNA abundance (**Figure 3e, f**). rORX-B, however, induced the expression of these transcription factors in a dose-dependent manner, but the effects were statistically significant only for PGC-1 α , PGC-1 β , and FoxO-1 with the high dose (**Figure 3e, f, h**).

Since mitochondria are the powerhouse of the cell with central importance for producing more than 90% of the ATP needed to carry out essential cellular functions, these organelles are important for proper growth and development of skeletal muscle tissue. The changes in expression of mitochondrial-related genes and their transcriptional regulators suggest that orexin may control mitochondrial respiratory function in avian muscle. To gain better insight into the physiological roles of orexins in avian muscle, mitochondrial bioenergetics in QM7 cells treated with 10 and 100 nM of rORX-A and rORX-B were assessed by monitoring basal oxygen consumption rate (OCR) followed by sequential treatment of cells with oligomycin, FCCP (carbonyl cyanide-4-phenylhydrazone), and antimycin A as shown in **Figure 4f**. As described previously [63], the decrease in OCR following oligomycin (which blocks ATP synthase) reveals OCR attributed to ATP synthesis activity. Maximal OCR is revealed in response to the uncoupling compound FCCP, and the difference between maximal OCR and basal OCR (prior to oligomycin) represents mitochondrial oxygen reserve capacity that cells can draw upon when increased energy production is needed. Oxygen consumption that remains following treatment with the electron transport inhibitor antimycin A is attributed to non-mitochondrial OCR (i.e., OCR due to activities other than non-mitochondrial c oxidase activity, such as mitochondrial reactive oxygen species production, oxidase activities, etc.). The amount of OCR attributed to proton leak is determined by the difference between oligomycin and antimycin A-inhibited OCR. When the non-mitochondrial component of cellular OCR was subtracted and by setting maximal OCR following FCCP at 100%, the effects of ORX-A and ORX-B on ATP synthesis, reserve capacity, and proton leak were determined. ATP synthesis was slightly elevated by both orexins, but the effect was not statistically discernable. Analysis of reserve capacity indicated no effect of both doses of rORX-A




Figure 3. Effect of orexin treatment on mitochondrial-related genes (a–d) and mitochondrial-transcriptional regulators (e–h) in QM7 cells. Cells were treated with recombinant orexin A or B (10 and 100 nM) for 24 h and the relative abundance of avian (av)-adenosine nucleotide translocator (ANT; a), UCP (b), Ski (c), nuclear respiratory factor 1 (NRF-1; d), PGC-1α (e), PGC-1β (f), peroxisome proliferator-activated receptor δ (PPARδ; g), and FoxO-1 (h) were determined by QPCR. Untreated cells were used as control. Protein levels of av-ANT and PGC-1α were measured by Western blot analysis. Data are expressed as means ± SE (n = 6). Significant difference between orexin-treated and control cells (P < 0.05).

and rORX-B; however, proton leak was significantly decreased by 10 nM of rORX-A, and by 100 nM of rORX-B (**Figure 4g**). The combined data from these experiments illustrate that orexins lead to the alteration of mitochondrial-related genes, their transcriptional regulators, and respiratory function in avian muscle cells. These changes suggest that orexin might also control mitochondrial dynamics (i.e., fusion/fission of mitochondria) in avian muscle.




Figure 4. Effect of orexin treatment on mitochondrial distribution (a), mitochondrial DNA and mass (b–e), and mitochondrial bioenergetics (f, g) in QM7 cells. a: Cells were cultured in chamber slides and treated with 100 nM of rORX-A or rORX-B for 24 h. Mitochondria were visualized with MitoTracker Red CMX Ros (75 nM) under a fluorescent microscope. Representative images acquired and deconvoluted are shown. (b–e) QM7 cells were treated with orexins (10 and 100 nM) for 24 h. The levels of mtDNA (b) and the relative expression of mtSSBP1 (c), mitochondrial markers CoxIV (d), and Cox5a (e) were determined by real-time PCR. (f, g) Oxygen consumption rate (OCR) (f) and the percentage of OCR due to proton leak (g) was determined using an XF24 flux Analyzer (Agilent Technologies). The values represent the means \pm SE (n = 6). *Significant difference between orexin-treated and control cells (P < 0.05).

4.2 Orexins differentially regulate mitochondrial biogenesis and dynamics in avian muscle cells

Since mtDNA replication and quantitation are a necessary component of mitochondrial biogenesis, the expression of mtDNA and mtSSBP1 was measured in orexin-treated QM7 cells as shown in **Figure 4b, c**. In contrast to rORX-A, in which both doses significantly downregulated mtDNA and upregulated mtSSBP1 expression, rORX-B (high dose) significantly increased mtDNA expression without




Figure 5. Effect of orexin treatment on mitochondrial dynamics-related genes in QM7 cells. QM7 cells were treated with orexins (10 and 100 nM) for 24 h. The relative expression of four genes involved in mitochondrial fusion, MFN1 (a), MFN2 (b), OPA1 (c), OMA1 (d) and three genes involved in mitochondrial fission, MTFP1 (f), DNMI1 (g), and MTFR1 (h) was determined by real-time PCR. The protein levels of MFN1, MFN2, and OPA1 were determined by Western blot analysis (e). The values represent the means \pm SE ($n = 6$). *Significant difference between orexin-treated and control cells ($P < 0.05$).

affecting mtSSBP1 levels. Consistent with these observations and in contrast to rORX-A, rORX-B increased mitochondrial content as visualized with MitoTracker Red probe staining (Figure 4a). Neither rORX-A nor rORX-B affected the expression of the mitochondrial transcription factor TFAM (data not shown). The expression of Cox IV and Cox 5a genes, commonly used markers for mitochondrial mass

and biogenesis, was determined. The high dose (100 nM) of rORX-A decreased Cox IV gene expression; however, the high-dose of rORX-B significantly increased Cox IV and Cox 5a mRNA levels compared with untreated cells (**Figure 4d, e**).

Mitochondria are dynamic organelles in the cell that constantly fuse and divide, forming constantly changing tubular networks, according to the needs of the organism, thus leading to alterations in their morphology and function [64]. Since orexin is shown to alter expression of mitochondrial-related genes and transcriptional regulators, it may also control avian muscle mitochondrial dynamics. The molecular mechanisms that control mitochondrial dynamics are complex and require participation and coordination of both the nuclear and mitochondrial genomes. In mammals this network has been partially unraveled after the identification of some of the genes responsible for mitochondrial fusion [mitofusins (MFN1 and MFN2), and optic atrophy 1 (OPA1)] and fission [dynamamin-related protein 1 (Drp1 or DNM1), fission 1 (FIS1), and mitochondrial protein 18 kDa]. Up until the current study, the integration of such a mitochondrial network is unknown in avian species. Following orexin treatment, the expression of four genes related to mitochondrial fusion and three genes related to mitochondrial fission were measured as shown in **Figure 5**. Recombinant ORX-B at high dose significantly induced the expression of MFN2, OPA1, and OMA1, but decreased the mRNA levels of MFN1 (**Figure 5a–d**). The same effect was observed at the protein levels (**Figure 5e**). However, rORX-A significantly downregulated the expression of MFN1 and OMA1 with both doses, and OPA1 with the high dose, but did not affect that of MFN2 (**Figure 5a–d**). Interestingly, and in contrast to rORX-B, where no significant effects were observed, rORX-A upregulated the expression of mitofission-related genes MTFP1, DNM1, and MTFR1 (**Figure 5f–h**). These orexin-induced changes in the expression of dynamics-related genes may serve in regulating mitochondrial metabolism in muscle cells in response to the needs of the animal during stages of growth and development.

5. Conclusions

Orexins are originally identified as hypothalamic neuropeptides that have potent orexigenic effects on appetite and feeding behavior in mammals. In avian species, however, orexins seem to be myokines that regulate the expression of their own system as well as muscle mitochondrial function, biogenesis, bioenergetics, and dynamics. As intensive genetic selection for fast growth rate, driven by human nutritional needs and economic demands, have resulted in dramatic increase in chicken body weight arising mainly from increased muscle mass, these findings open new vistas on the role of orexin system in muscle development and energy metabolism. Further in depth investigations are warranted to understand the relationship between orexin system, mitochondrial network, and muscle growth and development which, in turn, will be beneficial to the poultry industry as it has the potential to lead to increased production efficiency and reduced economic costs.

IntechOpen

IntechOpen

Author details

Kentu Lassiter* and Sami Dridi
Department of Poultry Science, Center of Excellence for Poultry Science,
University of Arkansas, Fayetteville, Arkansas

*Address all correspondence to: klassit@uark.edu

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] de Lecea L, Kilduff TS, Peyron C, Gao X-B, Foye PE, Danielson PE, et al. The hypocretins: Hypothalamus-specific peptides with neuroexcitatory activity. *Proceedings of the National Academy of Sciences of the United States of America*. 1998;**95**:322-327. DOI: 10.1073/pnas.95.1.322
- [2] Sakurai T, Amemiya A, Ishii M, Matsuzaki I, Chemelli RM, Tanaka H, et al. Orexins and orexin receptors: A family of hypothalamic neuropeptides and G protein-coupled receptors that regulate feeding behavior. *Cell*. 1998;**92**:573-585. DOI: 10.1016/S0092-8674(00)80949-6
- [3] Sakurai T. Orexins and orexin receptors: Implication in feeding behavior. *Regulatory Peptides*. 1999;**85**:25-30. DOI: 10.1016/S0167-0115(99)00076-2
- [4] Chemelli RM, Willie JT, Sinton CM, Elmquist JK, Scammell T, Lee C, et al. Narcolepsy in orexin knockout mice: Molecular genetics of sleep regulation. *Cell*. 1999;**98**:437-451. DOI: 10.1016/S0092-8674(00)81973-X
- [5] Belle MD, Hughes AT, Bechtold DA, Cunningham P, Pierucci M, Burdakov D, et al. Acute suppressive and long-term phase modulation actions of orexin on the mammalian circadian clock. *The Journal of Neuroscience*. 2014;**34**:3607-3621. DOI: 10.1523/JNEUROSCI.3388-13.2014
- [6] Tsuneki H, Murata S, Anzawa Y, Soeda Y, Tokai E, Wada T, et al. Age-related insulin resistance in hypothalamus and peripheral tissues of orexin knockout mice. *Diabetologia*. 2008;**51**:657-667. DOI: 10.1007/s00125-008-0929-8
- [7] Tsuneki H, Wada T, Sasaoka T. Role of orexin in the regulation of glucose homeostasis. *Acta Physiologica (Oxford, England)*. 2010;**198**:335-348. DOI: 10.1111/j.1748-1716.2009.02008.x
- [8] Tsuneki H, Wada T, Sasaoka T. Role of orexin in the central regulation of glucose and energy homeostasis. *Endocrine Journal*. 2012;**59**:365-374. DOI: 10.1507/endocrj.EJ12-0030
- [9] Shen Y, Zhao Y, Zheng D, Chang X, Ju S, Guo L. Effects of orexin A on GLUT4 expression and lipid content via MAPK signaling in 3T3-L1 adipocytes. *The Journal of Steroid Biochemistry and Molecular Biology*. 2013;**138**:376-383. DOI: 10.1016/j.jsbmb.2013.07.005
- [10] Skrzypski M, Le TT, Kaczmarek P, Pruszyńska-Oszmalek E, Pietrzak P, Szczepankiewicz D, et al. Orexin A stimulates glucose uptake, lipid accumulation and adiponectin secretion from 3T3-L1 adipocytes and isolated primary rat adipocytes. *Diabetologia*. 2011;**54**:1841-1852. DOI: 10.1007/s00125-011-2152-2
- [11] Ciriello J, Li Z, de Oliveira CV. Cardioacceleratory responses to hypocretin-1 injections into rostral ventromedial medulla. *Brain Research*. 2003;**991**:84-95. DOI: 10.1016/j.brainres.2003.08.008
- [12] Zhang W, Fukuda Y, Kuwaki T. Respiratory and cardiovascular actions of orexin-A in mice. *Neuroscience Letters*. 2005;**385**:131-136. DOI: 10.1016/j.neulet.2005.05.032
- [13] Samson WK, Bagley SL, Ferguson AV, White MM. Hypocretin/orexin type 1 receptor in brain: Role in cardiovascular control and the neuroendocrine response to immobilization stress. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*. 2007;**292**:R382-R387. DOI: 10.1152/ajpregu.00496.2006
- [14] Watt Executive Guide to World Poultry Trends [Internet]. 2017. Available from: <http://www.poultrytrends.com/201711/index>.

php?startid=9#/1 [Accessed: 7 December 2018]

[15] Havenstein GB, Ferket PR, Qureshi MA. Growth, livability, and feed conversion of 1957 versus 2001 broilers when fed representative 1957 and 2001 broiler diets. *Poultry Science*. 2003;**82**:1500-1508. DOI: 10.1093/ps/82.10.1500

[16] Barbut S, Sosnicki AA, Lonergan SM, Knapp T, Ciobanu DC, Gatcliffe LJ, et al. Progress in reducing the pale, soft and exudative (PSE) problem in pork and poultry meat. *Meat Science*. 2008;**79**:46-63. DOI: 10.1016/j.meatsci.2007.07.031

[17] Guernec A, Berri C, Chevalier B, Wacrenier-Cere N, Le Bihan-Duval E, Duclos MJ. Muscle development, insulin-like growth factor-I and myostatin mRNA levels in chickens selected for increased breast muscle yield. *Growth Hormone & IGF Research*. 2003;**13**:8-18. DOI: 10.1016/S1096-6374(02)00136-3

[18] Krzysik-Walker SM, Ocon-Grove OM, Maddineni SR, Hendricks GL 3rd, Ramachandran R. Is visfatin an adipokine or myokine? Evidence for greater visfatin expression in skeletal muscle than visceral fat in chickens. *Endocrinology* 2008;**149**: 1543-1550. DOI: 10.1210/en.2007-1301

[19] Akiba Y, Chida Y, Takahashi T, Ohtomo Y, Sato K, Takahashi K. Persistent hypoglycemia induced by continuous insulin infusion in broiler chickens. *British Poultry Science*. 1999;**40**:701-705. DOI: 10.1080/00071669987124

[20] Seki Y, Sato K, Kono T, Abe H, Akiba Y. Broiler chickens (Ross strain) lack insulin-responsive glucose transporter GLUT4 and have GLUT8 cDNA. *General and Comparative Endocrinology*. 2003;**133**:80-87. DOI: 10.1016/S0016-6480(03)00145-X

[21] Barre H, Cohen-Adad F, Duchamp C, Rouanet JL. Multilocular adipocytes from muscovy ducklings differentiated in response to cold acclimation. *The Journal of Physiology*. 1986;**375**:27-38. DOI: 10.1113/jphysiol.1986.sp016103

[22] Hood RL. The cellular basis for growth of the abdominal fat pad in broiler-type chickens. *Poultry Science*. 1982;**61**:117-121. DOI: 10.3382/ps.0610117

[23] Furuse M, Ando R, Bungo T, Ao R, Shimojo M, Masuda Y. Intracerebroventricular injection of orexins does not stimulate food intake in neonatal chicks. *British Poultry Science*. 1999;**40**:698-700. DOI: 10.1080/00071669987115

[24] Stadel JM, Wilson S, Bergsma D. Orphan G protein-coupled receptors: A neglected opportunity for pioneer drug discovery. *Trends in Pharmacological Sciences*. 1997;**18**:430-437. DOI: 10.1016/S0165-6147(97)01117-6

[25] Shibahara M, Sakurai T, Nambu T, Takenouchi T, Iwaasa H, Egashira SI, et al. Structure, tissue distribution, and pharmacological characterization of xenopus orexins. *Peptides*. 1999;**20**:1169-1176. DOI: 10.1016/S0196-9781(99)00120-5

[26] Alvarez CE, Sutcliffe JG. Hypocretin is an early member of the incretin gene family. *Neuroscience Letters*. 2002;**324**:169-172. DOI: 10.1016/S0304-3940(02)00195-7

[27] Sakurai T. Reverse pharmacology of orexin: From an orphan GPCR to integrative physiology. *Regulatory Peptides*. 2005;**126**:3-10. DOI: 10.1016/j.regpep.2004.08.006

[28] Tsujino N, Sakurai T. Orexin/hypocretin: A neuropeptide at the interface of sleep, energy homeostasis, and reward system. *Pharmacological Reviews*. 2009;**61**:162-176. DOI: 10.1124/pr.109.001321

- [29] Soppet DR, Li Y, Rosen CA. Human Genome Sciences Inc. Human Neuropeptide Receptor. 1996. World Patent No. WO9634877
- [30] Karteris E, Randeva HS, Grammatopoulos DK, Jaffe RB, Hillhouse EW. Expression and coupling characteristics of the CRH and orexin type 2 receptors in human fetal adrenals. *The Journal of Clinical Endocrinology and Metabolism*. 2001;**86**:4512-4519. DOI: 10.1210/jcem.86.9.7849
- [31] Randeva HS, Karteris E, Grammatopoulos DK, Hillhouse EW. Expression of orexin-A and functional orexin type 2 receptors in the human adult adrenals: Implications for adrenal function and energy homeostasis. *The Journal of Clinical Endocrinology and Metabolism*. 2001;**86**:4808-4813. DOI: 10.1210/jcem.86.10.7921
- [32] Karteris E, Machado RJ, Chen J, Zervou S, Hillhouse EW, Randeva HS. Food deprivation differentially modulates orexin receptor expression and signaling in the rat hypothalamus and adrenal cortex. *American Journal of Physiology. Endocrinology and Metabolism*. 2005;**288**:E1089-E1100. DOI: 10.1152/ajpendo.00351.2004
- [33] Hepler JR, Gilman AG. G proteins. *Trends in Biochemical Sciences*. 1992;**17**:383-387. DOI: 10.1016/0968-0004(92)90005-T
- [34] Gautam N, Downes GB, Yan K, Kisselev O. The g-protein betagamma complex. *Cellular Signalling*. 1998;**10**:447-455. DOI: 10.1016/S0898-6568(98)00006-0
- [35] Kukkonen JP. Physiology of the orexinergic/hypocretinergic system: A revisit in 2012. *American Journal of Physiology. Cell Physiology*. 2013;**304**(1):C2-C32. DOI: 10.1152/ajpcell.00227.2012
- [36] Kukkonen JP, Leonard CS. Orexin/hypocretin receptor signaling cascades. *British Journal of Pharmacology*. 2014;**171**(2):314-331. DOI: 10.1111/bph.12324
- [37] Ohkubo T, Boswell T, Lumineau S. Molecular cloning of chicken prepro-orexin cDNA and preferential expression in the chicken hypothalamus. *Biochimica et Biophysica Acta*. 2002;**1577**:476-480. DOI: 10.1016/S0167-4781(02)00483-9
- [38] Ohkubo T, Tsukada A, Shamoto K. cDNA cloning of chicken orexin receptor and tissue distribution: Sexually dimorphic expression in chicken gonads. *Journal of Molecular Endocrinology*. 2003;**31**:499-508. DOI: 10.1677/jme.0.0310499
- [39] Miranda B, Esposito V, de Girolamo P, Sharp PJ, Wilson PW, Dunn IC. Orexin in the chicken hypothalamus: Immunocytochemical localization and comparison of mRNA concentrations during the day and night, and after chronic food restriction. *Brain Research*. 2013;**1513**:34-40. DOI: 10.1016/j.brainres.2013.03.036
- [40] Godden KE, Landry JP, Slepneva N, Miguez PV, Pompeiano M. Early expression of hypocretin/orexin in the chick embryo brain. *PLoS One*. 2014;**9**(9):e106977. DOI: 10.1371/journal.pone.0106977
- [41] Arcamone N, D'Angelo L, de Girolamo P, Lucini C, Pelagalli A, Castaldo L. Orexin and orexin receptor like peptides in the gastroenteric tract of gallus domesticus: An immunohistochemical survey on presence and distribution. *Research in Veterinary Science*. 2014;**96**:234-240. DOI: 10.1016/j.rvsc.2014.02.002
- [42] Edwards CM, Abusnana S, Sunter D, Murphy KG, Ghatei MA, Bloom SR. The effect of the orexins on food intake: Comparison with

- neuropeptide Y, melanin-concentrating hormone and galanin. *The Journal of Endocrinology*. 1999;**160**:R7-R12. DOI: 0022-0795/99/0160-00R7
- [43] Katayama S, Hamasu K, Shigemi K, Cline MA, Furuse M. Intracerebroventricular injection of orexin-A, but not orexin-B, induces arousal of layer-type neonatal chicks. *Comparative Biochemistry and Physiology A*. 2010;**157**:132-135. DOI: 10.1016/j.cbpa.2010.05.018
- [44] da Silva ES, dos Santos TV, Hoeller AA, dos Santos TS, Pereira GV, Meneghelli C, et al. Behavioral and metabolic effects of central injections of orexins/hypocretins in pigeons (*Columba livia*). *Regulatory Peptides*. 2008;**147**:9-18. DOI: 10.1016/j.regpep.2007.12.003
- [45] Phillips-Singh D, Li Q, Takeuchi S, Ohkubo T, Sharp PJ, Boswell T. Fasting differentially regulates expression of agouti-related peptide, pro-opiomelanocortin, prepro-orexin, and vasoactive intestinal polypeptide mRNAs in the hypothalamus of Japanese quail. *Cell and Tissue Research*. 2003;**313**: 217-225. DOI: 10.1007/s00441-003-0755-8
- [46] Song Z, Liu L, Yue Y, Jiao H, Lin H, Sheikhahmadi A, et al. Fasting alters protein expression of AMP-activated protein kinase in the hypothalamus of broiler chicks (*Gallus gallus domesticus*). *General and Comparative Endocrinology*. 2012;**178**:546-555. DOI: 10.1016/j.ygcen.2012.06.026
- [47] Katayama S, Shigemi K, Cline MA, Furuse M. Clorgyline inhibits orexin-A-induced arousal in layer-type chicks. *The Journal of Veterinary Medical Science*. 2011;**73**:471-474. DOI: 10.1292/jvms.10-0358
- [48] Song Z, Everaert N, Wang Y, Decuyper E, Buyse J. The endocrine control of energy homeostasis in chickens. *General and Comparative Endocrinology*. 2013;**190**:112-117. DOI: 10.1016/j.ygcen.2013.05.006
- [49] Kiyashchenko LI, Mileykovskiy BY, Lai Y-Y, Siegel JM. Increased and decreased muscle tone with orexin (hypocretin) microinjections in the locus coeruleus and pontine inhibitory area. *Journal of Neurophysiology*. 2001;**85**:2008-2016. DOI: 10.1152/jn.2001.85.5.2008
- [50] Mileykovskiy BY, Kiyashchenko LI, Siegel JM. Muscle tone facilitation and inhibition after orexin-A (hypocretin-1) microinjections into the medial medulla. *Journal of Neurophysiology*. 2002;**87**:2480-2489. DOI: 10.1152/jn.2002.87.5.2480
- [51] Shiuchi T, Haque MS, Okamoto S, Inoue T, Kageyama H, Lee S, et al. Hypothalamic orexin stimulates feeding-associated glucose utilization in skeletal muscle via sympathetic nervous system. *Cell Metabolism*. 2009;**10**: 466-480. DOI: 10.1016/j.cmet.2009.09.013
- [52] Wong KK, Ng SY, Lee LT, Ng HK, Chow BK. Orexins and their receptors from fish to mammals: A comparative approach. *General and Comparative Endocrinology*. 2011;**171**:124-130. DOI: 10.1016/j.ygcen.2011.01.001
- [53] Kaslin J, Nystedt JM, Ostergard M, Peitsaro N, Panula P. The orexin/hypocretin system in zebrafish is connected to the aminergic and cholinergic systems. *The Journal of Neuroscience*. 2004;**24**:2678-2689. DOI: 10.1523/JNEUROSCI.4908-03.2004
- [54] Matsuda K, Azuma M, Kang KS. Orexin system in teleost fish. In: Litwack G, editor. *Vitamins and Hormones*. 1st ed. San Diego: Elsevier; 2012. pp. 341-361. DOI: 10.1016/B978-0-12-394623-2.00018-4
- [55] Lassiter K, Greene E, Piekarski A, Faulkner OB, Hargis BM, Bottje W,

- et al. Orexin system is expressed in avian muscle cells and regulates mitochondrial dynamics. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*. 2015;**308**:R173-R187. DOI: 10.1152/ajpregu.00394.2014
- [56] Nguyen PH, Greene E, Kong BW, Bottje W, Anthony N, Dridi S. Acute heat stress alters the expression of orexin system in quail muscle. *Frontiers in Physiology*. 2017;**8**:1079. DOI: 10.3389/fphys.2017.01079
- [57] Arihara Z, Takahashi K, Murakami O, Totsune K, Sone M, Satoh F, et al. Immunoreactive orexin-A in human plasma. *Peptides*. 2001;**22**:139-142. DOI: 10.1016/S0196-9781(00)00369-7
- [58] Sellayah D, Bharaj P, Sikder D. Orexin is required for brown adipose tissue development, differentiation, and function. *Cell Metabolism*. 2011;**14**: 478-490. DOI: 10.1016/j.cmet.2011.08.010
- [59] Swami M. Metabolism: Orexin acts on brown fat. *Nature Medicine*. 2011;**17**:1356. DOI: 10.1038/nm.2563
- [60] Pasban-Aliabadi H, Esmaeili-Mahani S, Abbasnejad M. Orexin-A protects human neuroblastoma SH-SY5Y cells against 6-hydroxydopamine-induced neurotoxicity: Involvement of PKC and PI3K signaling pathways. *Rejuvenation Research*. 2017;**20**:125-133. DOI: 10.1089/rej.2016.1836
- [61] Wan X, Liu Y, Zhao Y, Sun X, Fan D, Guo L. Orexin A affects HepG2 human hepatocellular carcinoma cells glucose metabolism via HIF-1 α -dependent and -independent mechanism. *PLoS One*. 2017;**12**(9):e0184213. DOI: 10.1371/journal.pone.0184213
- [62] Sikder D, Kodadek T. The neurohormone orexin stimulates hypoxia-inducible factor-1 activity. *Genes & Development*. 2007;**21**: 2995-3005. DOI: 10.1101/gad.1584307
- [63] Hill BG, Awe SO, Vladykovskaya E, Ahmed Y, Liu SQ, Bhatnagar A, et al. Myocardial ischaemia inhibits mitochondrial metabolism of 4-hydroxy-trans-2-nonenal. *The Biochemical Journal*. 2009;**417**:513-524. DOI: 10.1042/BJ20081615
- [64] Chen H, Chan DC. Emerging functions of mammalian mitochondrial fusion and, fission. *Human Molecular Genetics*. 2005;**14**(2):R283-R289. DOI: 10.1093/hmg/ddi270