We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Development of a Novel Antibacterial Medicine that Targets a Characteristic Lipid of the Cell Membranes of *Helicobacter pylori*

Kiyofumi Wanibuchi, Hisashi Masui, Takashi Takahashi
and Hirofumi Shimomura

Abstract

Helicobacter pylori is one of the most prevalent causes of gastritis. This pathogen colonizes for many years human stomach and asymptotically leads the persons to chronic gastritis. The eradication of *H. pylori* from human stomach is, therefore, important in order to prevent the digestive diseases including peptic ulcers and gastric cancer that develop via chronic atrophic gastritis. Wide-spectrum antibiotics such as amoxicillin and metronidazole are used for the treatment for *H. pylori* infectious diseases. However, the *H. pylori* strains resistant to these antibiotics are increasing year by year around the world. On this basis, we need urgently to develop the antibacterial medicines that act on *H. pylori* with a novel mechanism. Recent studies by our group have demonstrated that *H. pylori* shows susceptibility to the bactericidal action of indene compounds derived from decomposition of vitamin D. The bactericidal action of indene compounds is selective not against commonplace bacteria but against *H. pylori*. The indene compounds turned out to target the *H. pylori*'s phosphatidylethanolamine that retains a myristic acid as the saturated fatty acid side chain. These findings will contribute to the development of new antibacterial medicines specialized to the treatment for *H. pylori* infectious diseases.

Keywords: *Helicobacter pylori*, phosphatidylethanolamine, myristic acid, vitamin D, indene compound

1. Introduction

Helicobacter pylori is a Gram-negative microaerophilic helical bacillus equipped with polar flagella as the motility organ. This bacterium colonizes human stomach and causes chronic atrophic gastritis [1]. In addition to gastritis, the patients infected with this pathogen are capable of having various digestive diseases such as gastric ulcer, duodenal ulcer, gastric mucosa-associated lymphoid tissue (MALT) lymphoma, and gastric cancer [2–7]. Therefore, the eradication of *H. pylori* from human stomach is aggressively carried out around the world. Wide-spectrum

antibiotics such as amoxicillin, metronidazole, and clarithromycin are used for the treatment for *H. pylori* infectious diseases. However, the *H. pylori* strains resistant to the wide-spectrum antibiotics have been increasing year after year [8]. Especially, almost all *H. pylori* strains clinically isolated from African people have been acquiring the resistance to amoxicillin and metronidazole. In addition, wide-spectrum antibiotics act not only on *H. pylori* but also on commonplace bacteria inhabiting the mucosa of mouth and intestines. Therefore, the patients infected with *H. pylori*, who orally take wide-spectrum antibiotics for the treatment, often suffer from side effects such as stomatitis, constipation, and loose bowels resulted from the collapse of the balance of either oral bacterial flora or enterobacterial flora. When the side effects are serious, the patients develop pseudomembranous colitis accompanied with bloody feces and are compelled to discontinue the eradication of *H. pylori* [9]. To solve the difficult problems on the chemotherapy, we have to develop a novel antibacterial medicine that acts on only *H. pylori* without affecting the survival of human mucosal bacterial flora.

The assimilation of exogenous cholesterol into the cell membranes is one of the unique biological features of *H. pylori*. A part of cholesterol assimilated into the cell membranes is modified with a α -glucose at the carbon position-3 of its steroid framework, and the cholesteryl glucosides generated are used as the bacterial cell membrane constituents [10]. A previous study by our group has revealed that *H. pylori* possesses at least three types of cholesteryl glucosides, cholesteryl- α -D-glucopyranoside (CGL), cholesteryl-6-O-tetradecanoyl- α -D-glucopyranoside (CAG), and cholesteryl-6-O-phosphatidyl- α -D-glucopyranoside (CPG) (Figure 1) [11]. In addition to the three cholesteryl glucosides, other researchers have identified the lyso-type of CPG [12]. CGL is synthesized by the catalytic action of cholesterol α -glucosyltransferase (CGT) that localizes to the cytoplasm-side of the inner membrane of *H. pylori* [13–15]. CGT transfers a glucose derived from a uridine diphosphate glucose (UDP-Glc) to the cholesterol. Although the enzyme proteins involved in the synthesis of either CAG or CPG remain to be clarified, a recent study by other group has demonstrated that the enzymatic activities for the synthesis of CAG and CPG are detected in *H. pylori*'s outer membrane and inner membrane, respectively [16]. In addition, both enzymatic activities for the synthesis of CAG and CPG turned out to use phosphatidylethanolamine (PE) as the substrate [16]. In sum, the CGL acyltransferase (CGLAT) transfers a fatty acid derived from PE to the CGL and thereby synthesizes CAG. Meanwhile, the CGL phosphatidyltransferase (CGLPT) transfers a phosphatidyl group derived from PE to the CGL and thereby synthesizes CPG.

Figure 1.

Chemical structures of cholesteryl glucosides found in *H. pylori* cell membranes. In addition to the three types of cholesteryl glucosides, *H. pylori* possesses the lyso-type of CPG that dissociated a myristic acid. CGL, cholesteryl- α -D-glucopyranoside; CAG, cholesteryl-6-O-tetradecanoyl- α -D-glucopyranoside; CPG, cholesteryl-6-O-phosphatidyl- α -D-glucopyranoside; C14, myristic acid; C19, phytomonic acid.

A number of studies, including our own, have revealed that *H. pylori* assimilates exogenous cholesterol in order to acquire the resistance to the antibacterial actions of antibiotics and lipophilic compounds [17–20]. Meanwhile, *H. pylori* glucosylates the assimilated cholesterol to evade the host immune systems and/or to detoxify the toxic steroid compounds with 3 β -hydroxyl [12, 21]. The mechanism as for cholesterol uptake of *H. pylori*, however, remained for many years to be clarified. In a study in 2012, it has been revealed that PE of *H. pylori* cell membranes functions as a cholesterol-binding lipid [22]. PE is the most predominant glycerophospholipid component composing Gram-negative bacterial cell membranes. The PE of Gram-negative bacteria such as *Enterobacteriaceae* bacteria and *Pseudomonas aeruginosa* retains a palmitic acid (C_{16:0}) as the saturated fatty acid side chain, whereas the PE of *H. pylori* retains a myristic acid (C_{14:0}) as the saturated fatty acid side chain [22–26]. In sum, the PE molecular species composition of *H. pylori* turned out to completely differ from that of typical Gram-negative bacteria. A previous study by our group has demonstrated that the PE accounts for greater than 60% in the total lipids (excluding lipopolysaccharide) of *H. pylori* in the logarithmic growth phase [27]. Moreover, it has been revealed that the PE molecular species (DMPE) with two myristic acids accounts for approximately 30% in the total PE molecular species of *H. pylori* [22]. Intriguingly, DMPE showed higher binding affinity for cholesterol than for cholesteryl ester (**Figure 2**). In sum, the selective intermolecular interaction was found between the low-molecular-weight hydrophobic compounds.

Based on a number of studies including our own, the overview from the cholesterol assimilation to the cholesterol glucosylation was partially clarified in *H. pylori*: (1) cholesterol binds at least to DMPE of the outer membrane of *H. pylori* and is assimilated into the membranes; (2) a part of cholesterol is glucosylated by the catalytic action of CGT localized to the cytoplasm-side of the inner membrane, and thereby CGL is generated; (3) CGL is next exchanged to CAG and CPG by the enzymatic actions of CGLAT and CGLPT that localize to the outer membrane and the inner membrane, respectively (**Figure 3**). CGT utilizes an UDP-Glc as the glucose donor of cholesterol. Both enzymes of CGLAT and CGLPT utilize PE (myristoyl-PE) as the acyl group donor and phosphatidyl group donor of CGL,

Figure 2. Binding affinity of dimyristoyl-phosphatidylethanolamine of *H. pylori* cell membranes for either cholesterol or cholesteryl ester. Dipalmitoyl-PE with two palmitic acids (C_{16:0}) shows the high binding affinity for both of cholesterol and cholesteryl ester, whereas dimyristoyl-PE (DMPE) with two myristic acids (C_{14:0}) shows the selective high binding affinity only for cholesterol.

Figure 3.

Cholesteryl glucoside synthesis and its enzyme localization in *H. pylori* cell membranes. CGT, cholesterol α -glucosyltransferase; CGLAT, CGL acyltransferase; CGLPT, CGL phosphatidyltransferase; Cho, cholesterol; MPE, myristoyl phosphatidylethanolamine; LPS, lipopolysaccharide; UDP-Glc, uridine diphosphate-glucose.

respectively. Incidentally, cholesterol assimilated into *H. pylori* is distributed to both of the inner and outer membranes, whereas cholesteryl glucosides (CGL, CAG, and CPG) synthesized by *H. pylori* predominantly localize to the outer membrane [17].

As described above, it has been revealed that DMPE is one of the most prevalent lipid components of *H. pylori* cell membranes and shows the unique interaction not with cholesteryl ester but with cholesterol. Apart from this, previous studies by our group have demonstrated that a steroid hormone, progesterone acts on the cholesterol-binding site in the *H. pylori* cell membranes and confers the bactericidal action to *H. pylori* [28, 29]. Although it was unclear as for whether the progesterone shows the selective binding affinity for DMPE, this steroid turned out to destabilize the cell membrane structure of *H. pylori* and to ultimately induce the bacteriolysis. These findings drove us to the investigations of the low-molecular-weight hydrophobic compounds that induce the serious structure change to the DMPE molecule through the specific interaction. This chapter mentions the bactericidal activity of the indene compounds against *H. pylori*.

2. Development of new antibacterial medicines for *H. pylori*

2.1 Finding of an indene compound as the anti-*H. pylori* substance

The anti-*H. pylori* activity of various steroidal compounds was investigated. As a consequence, 25-hydroxyvitamin D₃ and 1,25-dihydroxyvitamin D₃ of secosteroids turned out to confer the strong bacteriolytic action to *H. pylori* [30]. However, these vitamin D₃ derivatives conspicuously attenuated their bactericidal activity by the nonbiological degradation (Table 1). In contrast, *H. pylori* did not almost

Seco-steroid		Vitamin D ₃	25-hydroxyvitamin D ₃	1,25-dihydroxyvitamin D ₃
Bactericidal activity	Intact compound	Weak	Strong	Strong
	Degradation	Augmentation	Attenuation	Attenuation

Table 1.
Bacteriolytic action of vitamin D₃ and its derivatives against *H. pylori*.

succumb to the bacteriolytic action of vitamin D₃. Surprisingly, the nonbiological degradation of vitamin D₃ augmented the bactericidal activity of its secosteroid against *H. pylori*. These results indicated that the vitamin D₃ derivatives directly act as bactericidal substances on *H. pylori*, and that some decomposition product of vitamin D₃ possesses potent bactericidal activity against *H. pylori*. It was, therefore, attempted to extract the anti-*H. pylori* substance from the decomposition products of vitamin D₃. As a consequence, the indene compound species (VDP1), otherwise known as Grundmann’s ketone, was successfully obtained [31, 32]. VDP1 is a low-molecular-weight hydrophobic compound in which the indene consisting of 5- and 6-membered rings of hydrocarbons was modified with alkyl, carbonyl, and methyl (**Figure 4**). Intriguingly, *H. pylori* showed high susceptibility to the bacteriolytic action of VDP1, whereas commonplace bacteria such as *Enterobacteriaceae* bacteria, *P. aeruginosa*, and *Staphylococcus aureus* showed insusceptibility to that of VDP1 [30]. In addition, VDP1 conferred the effective bacteriolytic action to *H. pylori* regardless of the assimilation of cholesterol into the cell membranes. These results indicate the possibility that VDP1 is a beneficial fundamental structure for the development of antibacterial medicines to selectively eradicate *H. pylori* without collapsing the balance of human mucosal bacterial flora.

2.2 Interaction between VDP1 and PE molecular species

The collapse induction activity of VDP1 against lipid vesicles was next examined using the unilamellar vesicles prepared with DMPE, dipalmitoyl-PE (DPPE), and dioleoyl-PE (DOPE). Intriguingly, VDP1 turned out to specifically induce the structure collapse of DMPE unilamellar vesicles without affecting the structural stability of either DPPE unilamellar vesicles or DOPE unilamellar vesicles (**Figure 5**). The structure collapse induction activity of VDP1 against DMPE unilamellar vesicles completely corresponded to the bactericidal activity of the indene compound

Figure 4. Bacteriolytic action of VDP1 against *H. pylori*. *H. pylori* shows high susceptibility to bacteriolytic action of VDP1 derived from decomposition of vitamin D₃. Meanwhile, commonplace bacteria such as *Escherichia coli* and *Staphylococcus aureus* show insusceptibility to bacteriolytic action of VDP1. VDP1, (1*R*,3*aR*,7*aR*)-1-[(1*R*)-1,5-dimethylhexyl]octahydro-7*a*-methyl-4*H*-inden-4-one.

Figure 5. Specific intermolecular interaction between VDP1 and DMPE. Dye-containing unilamellar vesicles were prepared with DMPE, dipalmitoyl-PE (DPPE) or dioleoyl-PE (DOPE) and incubated for 5 minutes in the presence or absence of VDP1. The absorbance (A_{590 nm}) of dye released from the PE unilamellar vesicles was measured (left graph). The data were obtained from the three independent experiments and are denoted with the mean A_{590 nm} ± standard deviation. Statistical significance (P < 0.05) was calculated by the Student's *t*-test. N.S. denotes “no statistical significance.” VDP1 induces the selective conformation change to the DMPE unilamellar vesicles. The intermolecular interactions between VDP1 and PE molecular species were simulated by the computational chemistry (right panel). VDP1 induces the conformational change only to a myristic acid side chain in the PE molecule without affecting the conformational stability of a palmitic acid side chain in the PE molecule.

against *H. pylori* that abundantly contains DMPE in the cell membranes. Based on these results, VDP1 was considered to specifically interact with the DMPE, to induce the serious structure change to the myristic acid side chain in the PE molecule, and to ultimately disrupt the vesicular conformation of DMPE. In addition, these results strongly suggested that VDP1 exerts the bactericidal effect on *H. pylori* by targeting at least DMPE of the cell membranes.

The intermolecular interaction between VDP1, DMPE, and DPPE was, therefore, simulated by the computational chemistry. One of the computer simulations showed that VDP1 induces “the winding-structure change” to a myristic acid side chain in DMPE molecule, while VDP1 induced no structure change to a palmitic acid side chain in DPPE molecule (**Figure 5**). The alkyl of VDP1 seemed to be crucial conformation for the induction of the structure change of the myristic acid side chain of DMPE. In other words, the slight difference of the length of carbon

chain composing the fatty acids in PE molecules appeared to play an important role on the specific interaction between VDP1 and PE molecular species.

2.3 Bactericidal mechanism of the indene compound species

As described earlier, VDP1 was considered to confer the bacteriolytic action to *H. pylori* by the disruption of the cell membranes through the induction of the structure change of DMPE that is one of the most prevalent PE molecular species constructing the bacterial cell membranes. To investigate in detail the crucial conformation of VDP1 for exerting the structure collapse effect on DMPE unilamellar vesicles and for exerting the antibacterial effect on *H. pylori*, various VDP1 derivatives were chemically synthesized [33]. The structure collapse induction activity of VDP1 against DMPE unilamellar vesicles was already ascertained to almost completely correspond to the bactericidal activity of the indene compound against *H. pylori*. When the carbonyl of VDP1 was replaced with a hydroxyl, the indene compound (VD3-1) maintained both activities against DMPE unilamellar vesicles and *H. pylori* (Table 2). As seen in VDP1, VD3-1 turned out to confer no structure collapse induction activity against DPPE unilamellar vesicles. In addition, VD3-1 also had no influence on the viability of commonplace bacteria such as *Enterobacteriaceae* bacteria, *P. aeruginosa*, and *S. aureus*. Intriguingly, VD2-2 that lacks the alkyl chain of VD3-1 was ascertained to induce no structure collapses of either DMPE unilamellar vesicles or DPPE unilamellar vesicles and to completely forfeit the effective bactericidal activity against *H. pylori*. In combination with the result of computer simulation as for the intermolecular interaction between VDP1 and DMPE, these results indicate that the alkyl structure in the indene compound

Indene compound	Collapse induction activity		Bactericidal activity		Summary
	DMPE vesicle	DPPE vesicle	<i>H. pylori</i>	Others*
VDP1	Potent	Lack	Potent	Lack	<div>Crucial conformation on the induction of structure change of the myristic acid side-chain in PE molecule</div> <div>Significant conformation on the stable binding to PE molecule</div> <div>Significant conformation on the reinforcement of electrostatic attraction to the phosphate head of PE molecule</div> <div>R, Carbonyl or Hydroxyl</div>
VD3-1	Potent	Lack	Potent	Lack
VD2-2	Lack	Lack	Lack	Lack	

**Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Serratia marcescens*, *Salmonella* sp., *Pseudomonas aeruginosa*, and *Staphylococcus aureus*. VD3-1, (1R,3aR,7aR)-7a-methyl-1-((R)-6-methylheptan-2-yl)octahydro-1H-inden-4-ol; VD2-2, (1R,3aR,7aR)-1-((S)-1-hydroxypropan-2-yl)-7a-methyloctahydro-1H-inden-4-ol.

Table 2.
Relationship between indene compounds, PE unilamellar vesicles and *H. pylori*.

species plays a crucial role on the induction of the structure change of the myristic acid side chain in DMPE molecule. Moreover, the alkyl structure in the indene compound species turned out to be essential for exerting the effective bacteriolytic action to *H. pylori*. The functional groups such as carbonyl and hydroxyl of the indene compound species seem to be significant conformation for bonding to the phosphate head in the PE molecules with an electrostatic attraction. Meanwhile, the indene-ring structure is guessed to be significant to stabilize the hydrophobic interaction between the indene compound species and PE molecules.

A recent study by our group has demonstrated that VDP1 confers the anti-bacterial action not only to *H. pylori* but also to other *Helicobacter* species [33]. Especially, *Helicobacter felis* showed high susceptibility to the bacteriolytic action of VDP1, as similar to *H. pylori*. *H. felis* is a Gram-negative microaerophilic spiral bacillus possessing bipolar tufts of flagella. This bacterium is isolated from the gastric mucosa of cats and dogs [34–36]. As seen in *H. pylori*, *H. felis* causes chronic gastritis and gastric MALT lymphoma in mouse when it colonizes the mouse stomach [37, 38]. An earlier study by our group has revealed that a myristic acid accounts for approximately 16% in the fatty acid composition of *H. felis* PE, and that the PE molecular species retaining a myristic acid and a palmitic acid accounts for approximately 37% in total PE molecular species of the bacteria [39]. Though *H. felis* completely succumbs to the bacteriolytic action at the same concentration of VDP1 (less than 3 µg/ml) that eradicates *H. pylori*, this *Helicobacter* species did not possess DMPE. On this basis, VDP1 is considered to interact not only with DMPE but also with myristoyl-PE that retains a myristic acid as one of the two fatty acid side chains. In addition, *H. felis* turned out to possess lauryl-PE as the most prevalent PE molecular species. The PE molecular species retaining a lauric acid (C_{12:0}) and a palmitic acid accounted for approximately 40% in total PE molecular species of *H. felis*. In sum, large parts of PE molecular species of *H. felis* bind a palmitic acid and either a myristic acid or a lauric acid as the fatty acid side chains. Given that a lauric acid is shorter in the length of carbon chain than a myristic acid, we can assume that VDP1 collapses the vesicular conformation consisting not only of myristoyl-PE but also of lauryl-PE. In the future, it will need to elucidate the hydrophobic interaction between VDP1 and lauryl-PE in addition to myristoyl-PE.

In contrast, *Helicobacter cinaedi* showed low susceptibility to the bacteriolytic action of VDP1, even though the PE molecular species composition in *H. cinaedi* is similar to the PE molecular species composition in *H. felis* [39]. *H. cinaedi* is a Gram-negative rod-like bacillus equipped with bipolar flagella and isolated from the intestinal tracts and livers of various mammals such as human, dog, cat, and hamster [40–42]. Therefore, this bacterium is classified into the enterohepatic *Helicobacter* species [43, 44]. Meanwhile, *H. pylori* and *H. felis* are classified into the gastric *Helicobacter* species. Most of the persons infected with *H. cinaedi* have no clinical symptoms, but some persons suffer from systematic inflammations, namely phlegmone, arthritis, and meningitis, due to the bacteremia [45]. Although it is unclear as for why *H. cinaedi* is lower in the VDP1-susceptibility than the two *Helicobacter* species, the involvement of lipopolysaccharide (LPS) is considered as one possibility. LPS is a glycolipid constructed of a long polysaccharide chain and fatty acids and is one of the composition components of the outermost layer of the outer membrane of Gram-negative bacteria [46]. The part of polysaccharide chain in LPS comes into direct contact with the outsides of the bacterial cells and limits the membrane permeability of various lipophilic compounds. The LPS contents in *H. cinaedi* may be higher than those in *H. pylori* and *H. felis*. The membrane permeability of VDP1 through the LPS barrier may be, therefore, stricter in *H. cinaedi* than in *H. pylori* and *H. felis*. Further investigation will be necessary to compare the LPS contents between the *Helicobacter* species.

Based on the current studies, the following bactericidal mechanism of the indene compounds synthetically derived from vitamin D in *H. pylori* was proposed: the indene compounds bind to the myristoyl-PE (including DMPE) of *H. pylori* cell membranes, induce “the winding-structure change” to the myristic acid side chain

Figure 6.
A proposed bactericidal mechanism of the indene compound species for *H. pylori*. The indene compound species such as VDP1 and VD3-1 bind to the myristoyl-PE of *H. pylori* cell membranes and induce “the winding-structure change” to the myristic acid side chain of the especially DMPE. *H. pylori* ultimately lyses via the destabilization of the membrane conformation.

Figure 7.
Role of cholesterol and cholesteryl glucosides in *H. pylori*. Cholesterol (Cho) binds to the myristoyl-PE (MPE) including dimyristoyl-PE (DMPE) of the *H. pylori*’s cell membranes but has no influence on the stability of the cell membrane conformation. As similar to cholesterol, no cholesteryl glucosides (CGL, CAG, and CPG) affect the cell membrane stability of *H. pylori*. These steroidal compounds are assimilated as the membrane lipid constituents of *H. pylori* and serve to strengthen the cell membrane lipid barrier. Incidentally, VDP1 confers the bacteriolytic action to *H. pylori* regardless of the cholesterol assimilation into the cell membranes.

in the PE molecules, destabilize the membrane conformation, and ultimately confer the bacteriolytic action to *H. pylori* (**Figure 6**).

In the case of the cholesterol assimilation in *H. pylori*, cholesterol is distributed to both membranes of the outer membrane and the inner membrane, and a part of the cholesterol is, thereafter, metabolized to cholesteryl glucosides (CGL, CAG, and CPG), and these metabolites localize to the outer membrane (**Figure 7**). Cholesterol and cholesteryl glucosides have no influence on the stability of the cell membrane conformation of *H. pylori*, and rather these steroidal compounds serve to strengthen the membrane lipid barrier of the bacteria on the limitation of the permeability of lipophilic compounds. A recent study by other group has demonstrated that cholesteryl glucosides are responsible for the morphological maintenance of *H. pylori*, for the acquirement of resistance to antibiotics such as polymyxin B, colistin, and tetracycline, and for the promotion of biofilm formation [47]. This suggests that cholesteryl glucosides of *H. pylori* play an important role to limit the membrane permeability of various low-molecular-weight compounds. However, VDP1 confers the bacteriolytic action even to *H. pylori* retaining cholesterol and cholesteryl glucosides. Further investigation will be necessary to clarify the detailed intermolecular interactions between myristoyl-PE and steroidal compounds.

3. Conclusions

Almost all hydrophobic drugs are pharmacologically designed to inhibit the functions of either protein molecules or nucleic acids in the target creature species. However, no drugs that target a characteristic lipid molecule in the creatures are discovered for many years. In addition, a number of biochemists on lipid research leave great achievements in the analysis of biosynthetic enzymes of various lipophilic compounds such as fatty acids and complex lipids and in the identification of receptors of various hydrophobic ligands such as steroid hormones and eicosanoids. However, these achievements are not as for lipid itself but as for merely proteins. This chapter described the unique interaction between lipids: the indene compound species specifically disrupt the vesicular conformation consisting of DMPE. These findings will bring the new aspects to the drug discovery research and the lipid biochemistry research.

Acknowledgements

This work was supported by a Grant-in-Aid from the Adaptable and Seamless Technology Transfer Program through Target-Driven R&A (A-STEP), JSPS KAKENHI (grant number 15 K08006), and JKA promotion funds from KEIRIN RACE.

Conflict of interest

The authors declare no conflict of interest.

IntechOpen

Author details

Kiyofumi Wanibuchi¹, Hisashi Masui¹, Takashi Takahashi¹
and Hirofumi Shimomura^{2*}

1 Faculty of Pharmaceutical Sciences, Yokohama University of Pharmacy,
Kanagawa, Japan

2 Independent Researcher, Kumamoto, Japan

*Address all correspondence to: shimomura20090601@gmail.com

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Marshall B, Warren JR. Unidentified curved bacilli on gastric epithelium in active chronic gastritis. *Lancet*. 1983;**1**:1273-1274
- [2] Graham DY. *Helicobacter pylori*: Its epidemiology and its role in duodenal ulcer disease. *Journal of Gastroenterology and Hepatology*. 1991;**6**:105-113
- [3] Forman D. The Eurogast study group. An international association between *Helicobacter pylori* infection and gastric cancer. *Lancet*. 1993;**341**:1359-1363
- [4] Wotherspoon AC, Oriz-Hidalgo C, Falzon MR, Isaacson PG. *Helicobacter pylori*-associated gastritis and primary B-cell gastric lymphoma. *Lancet*. 1999;**338**:1175-1176
- [5] Uemura N, Okamoto S, Yamamoto S, Matsumura M, Yamaguchi S, Yamakido M, et al. *Helicobacter pylori* infection and the development of gastric cancer. *The New England Journal of Medicine*. 2001;**345**:829-832
- [6] Peek JRM, Blaser MJ. *Helicobacter pylori* and gastrointestinal tract adenocarcinoma. *Nature Reviews Cancer*. 2002;**2**:28-37
- [7] Stolte M, Bayerdorffer E, Morgner A, Alpen B, Wundish T, Thiede C, et al. *Helicobacter* and gastric MALT lymphoma. *Gut*. 2002;**50**:III19-III24
- [8] De Francesco V, Giorgio F, Hassan C, Manes G, Vannella L, Panella C, et al. Worldwide *H. pylori* antibiotic resistance: A systematic review. *Journal of Gastrointestinal and Liver Diseases*. 2010;**19**:409-414
- [9] Trifan A, Girleanu I, Cojocariu C, Sfarti C, Singeap AM, Dorobat C, et al. Pseudomembranous colitis associated with a triple therapy for *Helicobacter pylori* eradication. *World Journal of Gastroenterology*. 2013;**19**:7476-7479
- [10] Shimomura H, Hosoda K, Hirai Y. Interaction of *Helicobacter pylori* cell membrane with non-esterified cholesterol and other steroids. *Open Journal of Medical Microbiology*. 2013;**3**:70-79
- [11] Hirai Y, Haque M, Yoshida T, Yokota K, Yasuda T, Oguma K. Unique cholesteryl glucosides in *Helicobacter pylori*: Composition and structural analysis. *Journal of Bacteriology*. 1995;**177**:5327-5333
- [12] Wunder C, Churin Y, Winau F, Warnecke D, Vieth M, Lindner B, et al. Cholesterol glucosylation promotes immune evasion by *Helicobacter pylori*. *Nature Medicine*. 2006;**12**:1030-1038
- [13] Lebrun AH, Wunder C, Hildebrand J, Churin Y, Zähringer U, Lindner B, et al. Cloning of a cholesterol- α -glucosyltransferase from *Helicobacter pylori*. *The Journal of Biological Chemistry*. 2006;**281**:27765-27772
- [14] Lee H, Wang P, Hoshino H, Ito Y, Kobayashi M, Nakayama J, et al. α 1,4-GlcNAc-capped mucin-type O-glycan inhibits cholesterol α -glucosyltransferase from *Helicobacter pylori* and suppress *H. pylori* growth. *Glycobiology*. 2008;**18**:549-558
- [15] Hoshino H, Tsuchida A, Kametani K, Mori M, Nishizawa T, Suzuki T, et al. Membrane-associated activation of cholesterol α -glucosyltransferase, an enzyme responsible for biosynthesis of cholesteryl- α -D-glucopyranoside in *Helicobacter pylori* critical for its survival. *Journal of Histochemistry and Cytochemistry*. 2011;**59**:98-105
- [16] Jan HM, Chen YC, Shih YY, Huang YC, Tu Z, Ingle AB, et al. Metabolic labelling of cholesteryl glucosides in *Helicobacter pylori* reveals how the uptake of human lipids enhances bacterial virulence. *Chemical Science*. 2016;**7**:6208-6216

- [17] Shimomura H, Hosoda K, Hayashi S, Yokota K, Oguma K, Hirai Y. Steroids mediate resistance to the bactericidal effect of phosphatidylcholines against *Helicobacter pylori*. FEMS Microbiology Letters. 2009;**301**:84-94
- [18] McGee DJ, George AE, Trainor EA, Horton KE, Hildebrandt E, Testerman TL. Cholesterol enhances *Helicobacter pylori* resistance to antibiotics and LL-37. Antimicrobial Agents and Chemotherapy. 2011;**55**:2897-2904
- [19] Trainor EA, Horton KE, Savage PE, Testerman TL, McGee DJ. Role of the HefC efflux pump in *Helicobacter pylori* cholesterol-dependent resistance to ceragenins and bile salts. Infection and Immunity. 2011;**79**:88-97
- [20] Correia M, Casal S, Vinagre J, Seruca R, Figueiredo C, Touati E, et al. *Helicobacter pylori*'s cholesterol uptake impacts resistance to docosahexanoic acid. International Journal of Medical Microbiology. 2014;**304**:314-320
- [21] Shimomura H, Hosoda K, McGee DJ, Hayashi S, Yokota K, Hirai Y. Detoxification of 7-dehydrocholesterol fatal to *Helicobacter pylori* is a novel role of cholesterol glucosylation. Journal of Bacteriology. 2013;**195**:359-367
- [22] Shimomura H, Hosoda K, Hayashi S, Yokota K, Hirai Y. Phosphatidylethanolamine of *Helicobacter pylori* functions as a steroid-binding lipid in the assimilation of free cholesterol and 3 β -hydroxyl steroids into the bacterial cell membrane. Journal of Bacteriology. 2012;**194**:2658-2667
- [23] Dunnick JK, O'Leary WM. Correlation of bacterial lipid composition with antibiotic resistance. Journal of Bacteriology. 1970;**101**:892-900
- [24] Kearns DB, Robinson J, Shimkets LJ. *Pseudomonas aeruginosa* exhibits directed twitching motility up phosphatidylethanolamine gradients. Journal of Bacteriology. 2001;**183**:763-767
- [25] Shokri A, Larsson G. Characterisation of the *Escherichia coli* membrane structure and function during fed batch cultivation. Microbial Cell Factories. 2004;**3**:9
- [26] Amine A, Saloua KS, Mouadh M, Alya EIM, Ahmed L. The absence of the "GATC-binding protein SeqA" affects DNA replication in *Salmonella enterica* serovar typhimurium, DNA replication and related cellular processes. In: Kusic-Tisma J, editor. DNA replication and related cellular processes. Rijeka, Croatia: InTech; 2011
- [27] Shimomura H, Hayashi S, Yokota K, Oguma K, Hirai Y. Alteration in the composition of cholesteryl glucosides and other lipids in *Helicobacter pylori* undergoing morphological change from spiral to coccoid form. FEMS Microbiology Letters. 2004;**237**:407-413
- [28] Hosoda K, Shimomura H, Hayashi S, Yokota K, Hirai Y. Steroid hormones as bactericidal agents to *Helicobacter pylori*. FEMS Microbiology Letters. 2011;**318**:68-75
- [29] Amgalanbaatar A, Shimomura H, Hosoda K, Hayashi S, Yokota K, Hirai Y. Antibacterial activity of a novel synthetic progesterone species carrying a linoleic acid molecule against *Helicobacter pylori* and the hormonal effect of its steroid on a murine macrophage-like cell line. The Journal of Steroid Biochemistry and Molecular Biology. 2014;**140**:17-25
- [30] Hosoda K, Shimomura H, Wanibuchi K, Masui H, Amgalanbaatar A, Hayashi H, et al. Identification and characterization of a vitamin D₃ decomposition product bactericidal against *Helicobacter pylori*. Scientific Reports. 2015;**5**:8860. DOI: 10.1038/srep08860

- [31] Windaus A, Grundmann W. Über die konstitution des vitamin D₂. II. Justus Liebigs Annalen der Chemie. 1936;**524**:295-299
- [32] Sicinski RR, DeLuca HF. Ruthenium tetroxide oxidation of Grundmann's ketone derivative from vitamin D₃. Bioorganic and Medical Chemistry Letters. 1995;**5**:159-162
- [33] Wanibuchi K, Hosoda K, Ihara M, Tajiri K, Sakai Y, Masui H, et al. Indene compounds synthetically derived from vitamin D have selective antibacterial action on *Helicobacter pylori*. Lipids. 2018;**53**:393-401
- [34] Lee A, Hazell SL, O'Rourke J, Kouprach S. Isolation of a spiral-shaped bacterium from the cat stomach. Infection and Immunity. 1988;**56**:2843-2850
- [35] Paster BJ, Lee A, Fox JG, Dewhirst FE, Tordoff LA, Fraser GJ, et al. Phylogeny of *Helicobacter felis* sp. Nov., *Helicobacter musterae*, and related bacteria. International Journal of Systematic Bacteriology. 1991;**41**:31-38
- [36] Hansen TK, Hansen PS, Nørgaard A, Nielsen H, Lee A, Andersen LP. *Helicobacter felis* does not stimulate human neutrophil oxidative burst in contrast to '*Gastrospirillum hominis*' and *Helicobacter pylori*. FEMS Immunology and Medical Microbiology. 2001;**30**:187-195
- [37] Enno A, O'Rourke JL, Howlett CR, Jack A, Dixon MF, Lee A. MALToma-like lesions in the murine gastric mucosa after long-term infection with *Helicobacter felis*. The American Journal of Pathology. 1995;**147**:217-222
- [38] Mohammadi M, Redline R, Nedrud J, Czinn S. Role of the host in pathogenesis of *Helicobacter*-associated gastritis: *H. felis* infection of inbred and congenic mouse strains. Infection and Immunity. 1996;**64**:238-245
- [39] Amgalanbaatar A, Hosoda K, Shimomura H. Phosphatidylethanolamine, but not phosphatidylglycerol-cardiolipin, isolated from two species of *Helicobacter* binds cholesterol over cholesteryl ester. Lipids. 2015;**50**:799-804
- [40] Kiehlbauch JA, Brenner DJ, Cameron DN, Steigenwalt AG, Makowski JM, Baker CN, et al. Genotypic and phenotypic characterization of *Helicobacter cinaedi* and *Helicobacter fennelliae* strains isolated from human and animals. Journal of Clinical Microbiology. 1995;**33**:2940-2947
- [41] Gebhart CJ, Fennell CL, Murtaugh MP, Stamm WE. *Campylobacter cinaedi* is normal intestinal flora in hamsters. Journal of Clinical Microbiology. 1989;**27**:1692-1694
- [42] Fox JG, Handt L, Sheppard BJ, Xu S, Dewhirst FE, Motzel S, et al. Isolation of *Helicobacter cinaedi* from the colon, liver, and mesenteric lymph node of a rhesus monkey with chronic colitis and hepatitis. Journal of Clinical Microbiology. 2001;**39**:1580-1585
- [43] Mikkonen TP, Kärenlampi RI, Hänninen ML. Phylogenetic analysis of gastric and enterohepatic helicobacter species based on partial HSP60 gene sequences. International Journal of Systematic and Evolutionary Microbiology. 2004;**54**:753-758
- [44] Fowsantear W, Argo E, Pattinson C, Cash P. Comparative proteomics of *Helicobacter* species: The discrimination of gastric and enterohepatic *Helicobacter* species. Journal of Proteomics. 2013;**97**:245-255
- [45] Kawamura Y, Tomida J, Morita Y, Fujii S, Okamoto T, Akaike T. Clinical and bacteriological characteristics of *Helicobacter cinaedi* infection. Journal of Infection and Chemotherapy. 2014;**20**:517-526

[46] Rietschel ET, Kirikae T, Schade FU, Mamat U, Schmidt G, Loppnow H, et al. Bacterial endotoxin: Molecular relationship of structure to activity and function. *The FASEB Journal*. 1994;**8**:217-225

[47] Qaria MA, Kumar N, Hussain A, Qumar S, Doddam SN, Sepe LP, et al. Roles of cholesteryl- α -glucoside transferase and cholesteryl glucosides in maintenance of *Helicobacter pylori* morphology, cell wall integrity, and resistance to antibiotics. *Molecular Biology and Physiology*. 2018;**9**:e01523-e01518