We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

α -Amylase Production by Toxigenic Strains of *Aspergillus* and *Penicillium*

Adekunle Odunayo Adejuwon
and Victoria Anatolyivna Tsygankova

Abstract

Aflatoxins are produced by a variety of fungal species and these have contributed to devastating health problems globally. However, apart from the capability of the production of aflatoxins, the productions of enzymes by like fungi have been explored. Aflatoxin B1-producing-toxigenic strains of *Aspergillus flavus* (A₁), *Aspergillus parasiticus* (A₂), *Penicillium citrinum* (P₁) and *Penicillium rubrum* (P₂) isolated from rice were grown on a defined medium with varying carbon and nitrogen sources. They were also grown on rice as sole carbon and nitrogen source for fungal growth. In an attempt to purify, the extracellular α -amylases produced were subjected to ammonium sulfate precipitation (40–90% saturation) followed by dialysis. The aflatoxin B1-producing toxigenic strains of *Aspergillus flavus* (A₁), *Aspergillus parasiticus* (A₂), *Penicillium citrinum* (P₁) and *Penicillium rubrum* (P₂) were able to produce α -amylases in both the growth medium with varying C and N sources of fungal and also in the rice medium. The most active α -amylase activity was produced by toxigenic *A. flavus* (A₁) with a value of 3.25 ± 0.15 Units and this was when ammonium sulfate was nitrogen source with starch as carbon source of fungal growth in the defined growth medium. These toxigenic fungal strains can be explored for the industrial production of α -amylases.

Keywords: α -amylase, toxigenic, fungi, aflatoxin B1

1. Introduction

1.1 *Aspergilli* and pathogenicity

Aspergilli are of the taxonomic Division Eumycota, Subdivision Eumycotina, Class Ascomycetes, Order Eurotiales, Family Trichocomaceae [1]. *Aspergillus* is a filamentous cosmopolitan and ubiquitous fungus commonly isolated from soil, plant debris and indoor air environment [2]. While the teleomorphic state exists for *Aspergillus species*, some are accepted to be mitosporic without any known sexual spore [3]. The genus *Aspergillus* includes over 185 species and about 20 species have so far been reported as causing opportunistic infections in man [4, 5]. Among these species, *Aspergillus fumigatus* is the most commonly isolated species followed by *Aspergillus flavus* and *Aspergillus niger*. *Aspergillus clavatus*, *Aspergillus glaucus*, *Aspergillus oryzae*, *Aspergillus terreus*, *Aspergillus ustus* and *Aspergillus versicolor* are

among the other species less commonly isolated as opportunistic pathogens [6, 7]. Food infected by *Aspergillus flavus* may be carcinogenic to humans and animals [8]. *Aspergillus flavus* is a saprophyte of grains. It produces mycotoxins in infected food [2]. Infection of peanuts (*Arachis hypogaea*) seeds by *Aspergillus flavus* and *Aspergillus parasiticus* is a serious problem that can result in aflatoxin contamination in the seed [9]. *Aspergillus flavus* produces aflatoxins B, G and cyclopiazonic acid CPA [10].

Beta-1,3-Glucanase activity in peanut seed is induced by infection with *Aspergillus flavus* [9]. Maize seeds are susceptible to *Aspergillus flavus* infection [11]. *Aspergillus flavus* causes Ear rot in corn with aflatoxin production. Resistance to aflatoxin production can be controlled by epistasis [12]. *Aspergillus flavus* causes kernel infection in maize, the Southwestern corn borer (SWCB) has been reported to substantially increase aflatoxin levels in such infection [13]. Kernels of corn genotype GT-MAS: gk are resistant to *Aspergillus flavus* [14]. A 14-KDa protein in corn kernel makes it resistant to *Aspergillus flavus* infection [15]. *Aspergillus flavus* found around corn storage cribs and bins are point sources of inoculum/infection with *Aspergillus flavus* in the corn agroecosystem [16]. Ear corn rot caused by *Aspergillus flavus* and *Aspergillus parasiticus* is severe in areas with high temperatures and drought [2]. *Aspergillus flavus* causes the post-harvest disease of *Arachis hypogaea* [3].

According to Norton [17], carotenoids in endosperm may decrease the amount of aflatoxin produced by *Aspergillus flavus*. *Aspergillus flavus* can be divided into S and L strains on the basis of sclerotial morphology [18]. Atoxigenic *Aspergillus flavus* L strain reduce formation of both sclerotia and aflatoxin when coinoculated with S strain isolate [18]. *Aspergillus flavus* L strain reduce formation of both sclerotia and aflatoxin when coinoculated with S strain isolate [18]. *Aspergillus flavus* produces aflatoxin in cotton seed with the S strain being highly toxigenic [19]. *Aspergillus parasiticus* isolated from soil from a corn field produced aflatoxin B(1) B(2) and G(1) G(2) [20]. *Aspergillus flavus* produces beta-glucuronidase [21]. *Aspergillus flavus* and *Aspergillus parasiticus* can contaminate agricultural crops with the production of toxic fungal metabolite aflatoxins. An endochitinase which is an inhibitory protein with M(r) of 29,000 is capable of inhibiting growth of *Aspergillus flavus* on maize [22]. Onion seeds stalk and flowers are susceptible to infection by *Aspergillus niger* Tiegh [23, 24]. The black-spored *Aspergillus* isolates that have been found to cause the disease fig smut are *Aspergillus niger* var. *niger*, *Aspergillus niger* var. *awamori*, *Aspergillus japonicus* and *Aspergillus carbonarius* [25]. Epiphytic fungi found on table grapes include *Epicoccum nigrum*, *Cladosporium herbarum*, *Alternaria alternata*, *Aspergillus niger* [26].

1.2 Penicilli and pathogenicity

Penicilli belongs to the taxonomic Division Eumycota, Subdivision Eumycotina, Class Ascomycetes, Order Eurotiales, Family Trichocomaceae [1]. Grape fruit green mold is caused by *Penicillium digitatum* [27]. *Penicillium* is common on citrus, gelly and preservatives. It is abundant in the soil and on decaying materials [3]. *Penicillium* spores are present in the air [3]. Studies have shown that *Penicillium* is important in the production of antibiotics such as Penicillin and Griseofluvin [5]. *Penicillium digitatum* causes the green mold of citrus fruits. Optimum temperature of their mycelia on such fruits is about 25°C [28]. The food borne pathogen *Listeria monocytogenes* has been observed to grow on apple infected with *Penicillium expansum* but not after 5 days [29]. The blue mold of decayed pear fruit is caused by *Penicillium expansum* [30]. *Penicillium expansum* has been observed to cause gray mold disease in apple and blue mold in pear [31]. *Penicillium* spp. have been isolated from pear stem [32] while *Penicillium italicum* cause

citrus blue mold and green mold [33, 34]. Postharvest green mold of oranges is caused by *Penicillium digitatum* [35, 36]. *Penicillium* sp. has been isolated from grapes [26]. *Penicillium expansum* causes the blue mold decay of pear [37]. *Penicillium digitatum* and *Penicillium italicum* cause postharvest green and blue molds of citrus fruits. Sporulation of both molds can be prevented or reduced by gaseous ozone without noticeable ozone phytotoxicity to the fruits [33, 34]. Apple fruits with stem pulls have been reported to be more susceptible to blue mold decay caused by *Penicillium expansum* than fruits with stems [38]. According to Spotts and Holz [39], *Penicillium expansum* can infect and cause disease in grape and plum fruits. Aqueous chlorine has been reported to reduce the viable spores of *Penicillium digitatum*, the causative fungi of the green mold and sour rot of citrus [40]. Sodium bicarbonate has been found to reduce postharvest decay of apple [41]. The level of resistance to decay of apple cultivars, caused by *Penicillium expansum*, varies from cultivar to cultivar [42]. Ziram but not calcium chloride control gray mold and bull's-eye rot, the postharvest decay of pear associated with the pathogen *Penicillium expansum* [43]. *Penicillium digitatum* has been associated with the postharvest green mold of oranges. Soda ash was observed to control this post-harvest disease [35, 36]. According to Smilanick et al. [35, 36], the effectiveness of imazalil for the control of citrus green mold caused by *Penicillium digitatum* improved significantly when the citrus fruits were treated with heated aqueous solutions of the fungicide as compared with the current commercial practice of spraying wax containing imazalil on the fruits. According to Smilanick et al. [44], fungicide applications with thiabendazole (TBZ) and sodium bicarbonate reduce green mold caused by *Penicillium digitatum* of citrus fruits and lemon fruits. It was also observed that pre harvest applications of thiophanate methyl to the fruits controlled postharvest green mold. Blue mold caused by *Penicillium* is an important postharvest disease of apple. *Penicillium expansum* and *Penicillium solitum* have been identified and isolated from rotten apple and pear fruits [45]. According to Sirois et al. [23, 24], onion seeds are affected by species of *Penicillium*.

1.2.1 *α*-Amylases

Amylases are hydrolytic enzymes that catalyze the degradation of starch molecules and other carbohydrates to yield dextrans and progressively smaller polymers composed of glucose units [46, 47]. Based on their pattern of catalysis and yield of products, amylases can be categorized as: alpha (α) amylase (endoamylase) (α -1,4-glucan-4-glucanohydrolase, EC 3.2.1.1); beta (β) amylase (exoamylase) (1,4- α -D-glucan maltohydrolase, EC 3.2.1.2); glucoamylase (exohydrolase) (Glucan-1,4- α -glucosidase, EC 3.2.1.3); pullulanase (α -dextrin endo-1,6- α -glucosidase, EC 3.2.1.41); isoamylase (EC 3.2.1.68). Pullulanases and isoamylases are termed deb-branching enzymes [46, 48, 49]. They can be of plant and microbial sources [50–52].

1.2.2 Microbial *α*-amylases

α -Amylases are produced by bacteria and fungi [53–58]. The two types of amylases commonly encountered in microbial degradation of starch are α and β amylases.

Degradation of substrate is important in enzymatic hydrolysis [59]. Starch is the substrate used in microbial amylase assay [60]. The starch molecules are hydrolyzed into polymers of glucose units [47]. According to Vihinen and Mantsala [61], starch-degrading enzymes are widely distributed among microbes and several activities are required to hydrolyze the starch into glucose units. *Bacillus subtilis* isolated from flour mill wastes produced a thermostable α -amylase in a complex medium containing starch [62]. According to Ajayi and Fagade [63], corn starch

can be used as substrate for β -amylase production by *Bacillus macerans*, *Bacillus licheniformis*, *Bacillus circulans*, *Bacillus coagulans*, *Bacillus megaterium*, *Bacillus polymyxa*, *Bacillus cereus* and *Bacillus subtilis*. According to Reiss et al. [64], approximately 80% of potato starch and 40–50% of grain starch were hydrolyzed by alpha amylase of certain microbes. Lactic acid bacteria have been found to ferment starchy foods to recover RNA though digestion with alpha amylase did not improve extraction [65]. *Rhizopus oligosporus*, a prolific amylase producer can degrade cassava tuber containing 65% starch into glucose [66]. *Thermoactinomyces thalpophilus* isolated from flour mill waste has been found to be capable of hydrolyzing 2% soluble starch [62]. A thermostable α -amylase activity from *Bacillus subtilis* isolated from flour mill waste was found to be more strongly expressed with corn starch than soluble starch [62]. Alpha amylase from *Bacillus licheniformis*, an hyperthermostable enzyme, is able to hydrolyze starch to medium-size oligosaccharides [67]. *Fusarium moniliforme* was found to produce alpha amylase in a culture medium containing starch [68].

Certain environmental (physical) factors affect amylase activity [48]. Lactic acid was found to be produced from *Lactobacillus delbrueckii* subsp. *delbrueckii* and defatted rice bran powder containing starch with coupled saccharification with amylase at 37°C and pH 5.0 [69]. An α -amylase produced by *Bacillus* sp. isolated from soil sample was optimally active at 75–80°C [70]. Alpha-amylase from *Bacillus licheniformis* is able to hydrolyze soluble starch within a temperature range of 60–75°C [71]. A thermophilic, moderately halophilic anaerobic *Halothermothrix orenii* synthesized an amylase similar to *Bacillus megaterium* amylase with optimal activity at 65°C [72]. Thermophilic *Thermus* sp. was reported to produce an extracellular α -amylase able to degrade starch at 70°C [73]. *Bacillus stearothermophilus* was found to produce a thermostable α -amylase active at 43°C [74]. According to Saito [75], *Bacillus licheniformis* produced a thermophilic extracellular α -amylase stable at 25°C but more active at an optimum temperature of 76°C. Manning and Campbell [76] reported that *Bacillus stearothermophilus* synthesized a thermostable α -amylase. *Rhizopus arrhizus* and *Rhizopus oryzae* were found to be capable of hydrolyzing starch at 30°C [77]. *Bacillus halodurans* produced an alkaline active maltohexaose-forming α -amylase active at 60°C. According to Oh et al. [57], *Lactobacillus gasseri* is able to synthesize a maltogenic amylase exhibiting optimum activity for β -CD hydrolysis at 55°C. Based on studies carried out by Najafi and Kembhavi [50], a marine *Vibrio* sp. produced an extracellular α -amylase with maximum activity at 55–60°C. According to Ogasahara et al. [78], *Bacillus stearothermophilus* was able to produce a thermophilic α -amylase with optimum temperature range of 65–73°C.

α -Amylase from *Pyrococcus woesei* has maximal activity at pH 5.6 [79]. A *Bacillus* sp. isolated from piglet caecum produced an extracellular alpha amylase optimally active at pH 7.0 [80]. A microorganism from uncultured soil was observed to produce amylolytic enzyme with optimal pH of 9.0 [81]. A mutant of *Bacillus amyloliquefaciens* has been reported to synthesize alpha amylase with optimal activity at pH 7.0 [82]. An extracellular alpha amylase isolated from cell free broth of *Streptomyces megasporus* grown in glucose, soluble starch and raw starch was stable at a pH range of 5.5–8.5 but with optimum activity at pH 6.0 [83]. Amylases in culture supernatants of an environmentally derived microbial mixed culture selected for its ability to utilize starch-containing plastic films as sole carbon sources produced amylases active at pH 5.5 and 8.0 [84]. Alpha amylase from *Thermoactinomyces vulgaris* had optimum activity at pH 4.8–6.0 [85]. Starch degradation by *Rhizopus oryzae* was favorable at pH 6.0 [77]. Akindahunsi [86] reported that waste water from cassava mash fermented by pure strains of *Saccharomyces cerevisiae*, *Lactobacillus delbrueckii* and *Lactobacillus coryniformis* produced amylase after 3 days with maximal activity at pH 6.0.

According to Mijts and Patel [72], the thermophilic, moderately halophilic anaerobic *Halothermothrix orenii* is able to synthesize alpha amylase active with specific activity of 2232 U mg⁻¹ requiring CaCl₂ for optimum activity and thermostability. The maltooligosaccharide-forming amylase from *Bacillus circulans* is enhanced by CO²⁺ and Mg²⁺ [87]. Amylase synthesized by *Lipomyces starkeyi* was found to be actively stable in a commercial mouthwash [88]. A salt-tolerant thermostable amylase produced by *Bacillus megaterium* was reported to be stable at 5 M NaCl [89]. A thermophilic *Thermoascus aurantiacus* has been observed to produce amylase with thermostability enhanced by calcium chloride [90]. Amylase production from *Bacillus sphaericus* was reported to be maximum with 3 mM divalent cations Mg⁺⁺ and Ca⁺⁺ incorporated in a growth medium [91]. Cadmium, Cobalt, Copper, Manganese, Nickel and Lead incorporated into Czapek-Dox liquid medium supported growth and production of amylase by soil yeasts *Geotrichum capitatum* and *Geotrichum candidum* [92]. Activity of α-amylase from a marine *Vibrio* sp. was found to be restored by Fe²⁺, Mn²⁺, Co²⁺, Ca²⁺, Mg²⁺ and Cu²⁺ to nearly 25–55% [50]. A *Bacillus* sp. produced an alkaliphilic amylase which was enhanced by Na⁺ and Co²⁺ [93]. According to Mishra et al. [94], *Bacillus subtilis* produced an α-amylase. Herbizid has been reported to activate amylase production in culture of *Fusarium oxysporum*, *Mucor niemalis* and *Penicillium chrysogenum* [95].

Amylase from *Fusarium verticillioides* has been found to be inhibited by a hydrophobic 19.7-KDa inhibitor from corn kernel [96, 97]. *Fusarium moniliforme*, a mycotoxigenic fungus has been reported to produce an amylase inhibited by a specific amylase inhibitor found in corn [96, 97]. Alpha-amylase inhibitor has been isolated from culture medium of *Streptomyces parvullus* [98]. *Streptomyces aureofaciens* produces a novel polypeptide inhibitor [99]. A strain of *Streptomyces nigrifaciens* has been reported to produce an amylase inhibitor having inhibitory effects on alpha amylase and glucoamylase [100].

Bacillus subtilis isolated from soil produced a starch degrading amylase with molecular weight of 50 KDa and an isoelectric point of 4.9 [101]. *Streptomyces lividans* has been reported to have a molecular weight of 107,054 KDa [102]. Alpha amylases from some *Bacillus* spp. were detected to possess molecular weight of approximately 65,5854 and 49 KDa [103]. Alpha-amylase of *Clostridium thermosulfurogenes* has been reported to have a molecular mass of 75,112 Da [104]. According to Kang et al. [105], *Bacillus stearothermophilus* produces an alpha-amylase which was glycosylated and with molecular weights of approximately 61–75 KDa.

According to Lorentz [106], Protected 4-Nitrophenyl-1,4–1-D-maltoheptaoside can be used in routine amylase assay. A simple and rapid method using Remazol Brilliant Blue-starch as substrate which is non-destructive allows direct visualization and isolation of amylolytic microorganisms from the environment [107].

Alpha amylase can be used in improving anaerobic solid waste treatment [108]. Carbohydrate-hydrolyzing enzymes have long been used by industrial product markers as major catalysts to transform raw materials into end products in such areas as food processing, beverage production, animal nutrition, leather and textiles [109].

With the advent of new frontiers in biotechnology, the spectrum of amylase application has widened in many fields such as clinical, medicinal and fine-chemical industries, as well as a widespread application of starch saccharification in the textile, food, brewing and distilling industries [110].

1.2.3 Aflatoxin B₁ and α-amylase production

According to Mellon et al. [111], an aflatoxin B₁ producer strain of *Aspergillus flavus* seem to possess the ability to produce numerous extracellular hydrolases

(α -amylase inclusive). Aflatoxin B1 have been detected in groundnut and maize contaminated with *Aspergillus flavus* [112]. The removal of lipids from ground substrates significantly reduced the substrate's potential for aflatoxin B1 (AFB1) production by *Aspergillus flavus*. However, maltose, glucose, arginine, glutamic acid, aspartic acid and zinc significantly induced the AFB1 production up to 1.7–26.6 fold [113]. Aflatoxins B₁, B₂, G₁ and G₂ and α -amylase were detected in *Aspergillus oryzae*, *Aspergillus flavus*, *Talaromyces spectabilis*, *Pacilomyces variotii* and *Lichtheimia* sp. isolated from nuruks in several regions of Korea [114]. According to Fakhoury and Woloshuk [115], a mutant strain of *Aspergillus flavus* failed to produce extracellular α -amylase when the Amy1 gene necessary for the production of α -amylase was disrupted in an aflatoxigenic strain (an aflatoxin B1 producing strain) of the fungus. Mycotoxigenic strains (aflatoxin B1 producing strains) of *Fusarium moniliforme* and *Aspergillus flavus* were capable of α -amylase production in a medium composed of 2% ground corn in milky stage corn [116]. In their attempt to increase aflatoxin B1 resistance in maize, Rajasekaran et al. [117] discovered that the α -amylase inhibitor-like protein (AILP) seem to play a role in the inhibition of *Aspergillus flavus* α -amylase and fungal growth. Fountain et al. [118] reviewed the nature of the interaction occurring between aflatoxin production by *Aspergillus flavus*, the environment in which the fungus thrives and its susceptibility to crop host before harvest. They proposed future directions for elucidating future relationship between resistance and susceptibility to the fungus' colonization, abiotic stress and its relationship to oxidative stress in which its aflatoxin B1 production may function as a form of antioxidant protection to the producing fungus. In a known positive transcriptomic database, E-probe Diagnostic for Nucleic acid Analysis (EDNA), a bioinformatic tool, originally developed to detect plant pathogens in mutagenomic databases, is capable of discriminating between production and non-production of aflatoxin B1 by *Aspergillus flavus* [119]. Substrate-induced lipase gene expression might be indirectly related to aflatoxin formation by providing the basic building block "acetate" for aflatoxin B1 synthesis in aflatoxin-producing *Aspergillus flavus* and *Aspergillus parasiticus* [120]. According to Smith et al. [121], silencing of the aflatoxin gene cluster in a certain strain of aflatoxin B1 producing *Aspergillus flavus* is suppressed by ectopic aflR gene (the transcriptional regulator of the aflatoxin biosynthetic gene cluster) expression.

1.2.4 Rice (*Oryza sativa*)

Rice (*Oryza sativa*) is a monocotyledonous cereal which belongs to the Grass family Gramineae or Poaceae [122]. With over 7000 varieties of rice, its pericarp and embryo contain 70–80% starch, 7% proteins, 1.5% oils, some vitamins (mostly A, B and C) and some essential minerals [3]. According to Sizer and Whitney [123], rice contains fiber and the vitamin folate and provides 80% of the calories consumed by humans worldwide [122]. It contains 12 chromosomes in a haploid set [124]. The domestication of rice formed part of the basis for civilization in the near East, far East and the New World [125]. Feeding more people worldwide than any other crop, rice is the only crop grown exclusively for human consumption [125]. Sedentary irrigated rice production in tropical lowlands can support hundreds of people per square kilometer, explaining the wide spread importance of rice crops in the tropics [126]. The discovery of Gibberellins arose from infected rice [122]. *Oryza sativa* is the main cultivated rice species but over 20 species in the genus are known [125].

This research was designed to examine the production and activity of α -amylases by some toxigenic aflatoxin B1-producing strains of *Aspergillus* and *Penicillium* isolated from deterioration rice. Attempts were made to purify the α -amylases.

1.2.5 Contribution to knowledge

The present research will establish the presence of α -amylases in rice during mycotic spoilage by toxigenic strains of *Aspergillus flavus* (A_1), *Aspergillus parasiticus* (A_2), *Penicillium citrinum* (P_1) and *Penicillium rubrum* (P_2). These fungi, being capable of producing these enzymes can be used in the production of amylases. Rice as substrate can be explored in such production.

Amylases are used in clinical chemistry most especially in diagnosis. Their combination with proteases and lipases are also employed industrially in the bio-remediation of recalcitrants/organic pollutants and the hydrolytic digestion of the peptidoglycan layers of both gram positive and gram negative bacteria in wastewaters before chlorination [109].

2. Materials and methods

2.1 Sources and identification of isolates

The isolates of aflatoxin B1-producing-toxigenic strains of *Aspergillus flavus* (A_1), *Aspergillus parasiticus* (A_2), *Penicillium citrinum* (P_1) and *Penicillium rubrum* (P_2) for this research were from deteriorated rice and identified at the Seed Health Unit of the International Institute for Tropical Agriculture, Ibadan, Nigeria using techniques contained in the illustrated Handbook of fungi [127, 128]. The identification was done by observing cultural and morphological characteristics. Each isolate was cultured on Potato Dextrose agar. The nature of growth, rate of growth, colony color and sporulation patterns were carefully observed. Sporulating mature cultures was used in microscopic examination. Fungal samples were taken from advancing margins and centers of the growth regions with the aid of sterile inoculating needle. The samples were smeared on glass slides and stained with lactophenol cotton blue. After placing the cover slips, macroscopic and microscopic morphological characteristics like arrangement and shape of spores, type of sporangia, type of hyphae, presence or absence of septa on hyphae was examined under the high power objective of a compound binocular microscope.

2.2 Culture conditions and preparation of inocula

The isolates were subcultured and maintained on Potato Dextrose agar plates and slants. Each fungus was further subcultured into test tubes of the same medium and incubated at 25°C. A 96-hr-old culture of toxigenic strains of *Aspergillus flavus* (A_1), *Aspergillus parasiticus* (A_2) and *Penicillium rubrum* (P_2) and 120-h-old culture of *Penicillium citrinum* (P_1) was used as inocula. According to the modified method of Olutiola and Ayres [129], cultures was grown in a defined medium of the under-listed composition: $MgSO_4 \cdot 7H_2O$, K_2HPO_4 , KH_2PO_4 , L-cysteine, biotin, thiamine and $FeSO_4 \cdot 7H_2O$ with added carbon and nitrogen sources (Sigma). Conical flasks (250 ml) containing 100 ml growth medium will be inoculated with 1 ml of an aqueous spore suspension containing approximately 6×10^4 spores per ml of each isolate. Experimental and control flasks was incubated without shaking at 25°C [130].

2.3 Rice as a source of carbon

Rice (Caprice) from Spain was bought at the main market, Bodija, Ibadan, Nigeria. The rice was added to distilled water (1% w/v) and autoclaved at 15lb/in² at

121°C. Experimental Conical flasks (250 ml) containing 100 ml of the rice medium was inoculated with 1 ml of an aqueous spore suspension containing approximately 6×10^4 spores per ml of each isolate. Control flasks contained sterilized rice medium not inoculated with aqueous spore suspension of the isolate. Experimental and control flasks was incubated without shaking at 25°C.

On a daily basis, the contents of each flask was filtered through glass fiber filter paper (Whatman GF/A). The protein content of the filtrates was determined using the method of Lowry et al. [131]. The filtrates were analyzed for amylase activity using the modified methods of Pfueller and Elliott [132] and Xiao et al. [133]. The filtrates were used as crude preparation.

2.4 Ammonium sulfate fractionation

The crude enzymes were treated with ammonium sulfate (analytical grade) within the limits of 40–90% saturation. Precipitation was allowed to continue at 4°C for 24 h. The mixtures were then centrifuged 10,000 g for 30 min at 4°C using a high speed cold centrifuge (Optima LE-80 K Ultracentrifuge, Beckman, USA). The supernatant was discarded. The precipitate was re-dissolved in 0.2 M citrate phosphate buffer, pH 6.0. The protein contents were determined using the Lowry et al. [131] method while amylase activity was determined using the modified methods of Pfueller and Elliott [132] and Xiao et al. [133].

2.5 Dialysis

Using acetylated dialysis tubings (Visking dialysis tubings, Sigma) [134] and a multiple dialyser (Pope Scientific Inc. Model 220, USA), the enzyme preparations were dialysed under several changes of 0.2 M citrate phosphate buffer pH 6.0 at 4°C for 24 h. The protein contents of the dialysed enzymes were determined using the Lowry et al. [131] method while amylase activity was determined using the modified methods of Pfueller and Elliott [132] and Xiao et al. [133].

2.6 Enzyme assay

Both experimental (fungal isolate inoculated) and control (un-inoculated) flasks were assayed for amylase activity.

2.6.1 α -Amylase

α -Amylase activity was determined using the modified methods of Pfueller and Elliott [132] and Xiao et al. [133]. The reaction mixtures consisted 2 ml of 0.1% (w/v) starch (Sigma) in 0.2 M citrate phosphate buffer, pH 6.0 as substrate and 0.5 ml of enzyme. These were the experimentals in the assay procedure. The controls in the assay procedure consisted only 2 ml of the prepared substrate. The contents of both experimental and control tubes were incubated at 35°C for 30 min. The reactions were terminated with 3 ml of 1 N HCl. Enzyme (0.5 ml) was added to the contents of each control. About 2 ml of the mixture from each of the sets of experimentals and controls was transferred into new sets of clean test tubes. About 3 ml of 0.1 N HCl was added into the contents of each test tube after which 0.1 ml of iodine solution was added. Optical density readings were taken spectrophotometrically at 620 nm. Enzyme activity was defined in units and specific activity as enzyme units per mg protein.

One unit of α -amylase activity was defined as the amount of enzyme which produced 0.1% reduction in the intensity of the blue color of starch-iodine complex under conditions of the assay.

3. Results

3.1 Amylase activities of isolates on growth media

Toxigenic strains of *Aspergillus flavus* (A₁), *Aspergillus parasiticus* (A₂), *Penicillium citrinum* (P₁) and *Penicillium rubrum* (P₂) grew and exhibited amylase activities, varyingly, in modified growth medium used for this research.

Using different carbon sources (rice, starch, maltose, sucrose, lactose, glucose and galactose) in the growth medium, amylase activity expressed by each isolate on the tenth day of incubation is shown in **Table 1**.

With different sources of nitrogen (NH₄Cl, urea, KNO₃, ammonium sulfate, glycine, sodium nitrate, tryptone and peptone) in the growth medium, amylase activity expressed varyingly by each isolate on the tenth day of incubation is shown in **Table 2**.

Toxigenic *P. citrinum* (P₁) produced active α-amylase (0.75 ± 0.01 Units) and this was when potassium nitrate was nitrogen source with maltose as carbon source of the defined growth medium. Toxigenic *A. parasiticus* (A₂) also expressed an α-amylase activity value of 0.72 ± 0.04 Units when rice was both carbon and nitrogen source of medium for fungal growth (**Table 1**).

Carbon source	Isolate	Amylase activity (Units)
Rice	<i>Aspergillus flavus</i> (A ₁)	0.54 ± 0.01
	<i>Aspergillus parasiticus</i> (A ₂)	0.72 ± 0.04
	<i>Penicillium citrinum</i> (P ₁)	0.43 ± 0.23
	<i>Penicillium rubrum</i> (P ₂)	0.62 ± 0.06
Galactose	<i>Aspergillus flavus</i> (A ₁)	0.06 ± 0.01
	<i>Aspergillus parasiticus</i> (A ₂)	0.53 ± 0.13
	<i>Penicillium citrinum</i> (P ₁)	0.36 ± 0.05
	<i>Penicillium rubrum</i> (P ₂)	0.09 ± 0.04
Glucose	<i>Aspergillus flavus</i> (A ₁)	0.50 ± 0.04
	<i>Aspergillus parasiticus</i> (A ₂)	0.63 ± 0.08
	<i>Penicillium citrinum</i> (P ₁)	0.44 ± 0.08
	<i>Penicillium rubrum</i> (P ₂)	0.32 ± 0.11
Lactose	<i>Aspergillus flavus</i> (A ₁)	0.10 ± 0.00
	<i>Aspergillus parasiticus</i> (A ₂)	0.66 ± 0.10
	<i>Penicillium citrinum</i> (P ₁)	0.40 ± 0.17
	<i>Penicillium rubrum</i> (P ₂)	0.37 ± 0.08
Maltose	<i>Aspergillus flavus</i> (A ₁)	0.52 ± 0.03
	<i>Aspergillus parasiticus</i> (A ₂)	0.68 ± 0.04
	<i>Penicillium citrinum</i> (P ₁)	0.75 ± 0.01
	<i>Penicillium rubrum</i> (P ₂)	0.58 ± 0.12
Starch	<i>Aspergillus flavus</i> (A ₁)	0.45 ± 0.04
	<i>Aspergillus parasiticus</i> (A ₂)	0.57 ± 0.12
	<i>Penicillium citrinum</i> (P ₁)	0.68 ± 0.03
	<i>Penicillium rubrum</i> (P ₂)	0.60 ± 0.14
Sucrose	<i>Aspergillus flavus</i> (A ₁)	0.46 ± 0.05
	<i>Aspergillus parasiticus</i> (A ₂)	0.69 ± 0.03
	<i>Penicillium citrinum</i> (P ₁)	0.59 ± 0.13
	<i>Penicillium rubrum</i> (P ₂)	0.39 ± 0.06

Each value represents the mean of three replicates with standard error.

Table 1.
Effect of carbon sources on activity of amylase produced by isolates.

Nitrogen source	Isolate	Amylase activity (Units)
Ammonium sulfate	<i>Aspergillus flavus</i> (A ₁)	3.25 ± 0.15
	<i>Aspergillus parasiticus</i> (A ₂)	0.05 ± 0.00
	<i>Penicillium citrinum</i> (P ₁)	0.38 ± 0.13
	<i>Penicillium rubrum</i> (P ₂)	0.13 ± 0.03
Glycine	<i>Aspergillus flavus</i> (A ₁)	0.38 ± 0.13
	<i>Aspergillus parasiticus</i> (A ₂)	2.48 ± 0.03
	<i>Penicillium citrinum</i> (P ₁)	0.00 ± 0.00
	<i>Penicillium rubrum</i> (P ₂)	1.53 ± 0.48
Potassium nitrate	<i>Aspergillus flavus</i> (A ₁)	0.50 ± 0.00
	<i>Aspergillus parasiticus</i> (A ₂)	1.30 ± 0.10
	<i>Penicillium citrinum</i> (P ₁)	0.25 ± 0.25
	<i>Penicillium rubrum</i> (P ₂)	0.68 ± 0.03
Ammonium chloride	<i>Aspergillus flavus</i> (A ₁)	3.02 ± 0.18
	<i>Aspergillus parasiticus</i> (A ₂)	0.05 ± 0.05
	<i>Penicillium citrinum</i> (P ₁)	0.13 ± 0.13
	<i>Penicillium rubrum</i> (P ₂)	0.13 ± 0.03
Peptone	<i>Aspergillus flavus</i> (A ₁)	0.25 ± 0.00
	<i>Aspergillus parasiticus</i> (A ₂)	1.50 ± 0.15
	<i>Penicillium citrinum</i> (P ₁)	0.13 ± 0.13
	<i>Penicillium rubrum</i> (P ₂)	2.48 ± 0.03
Sodium nitrate	<i>Aspergillus flavus</i> (A ₁)	0.38 ± 0.13
	<i>Aspergillus parasiticus</i> (A ₂)	1.48 ± 0.13
	<i>Penicillium citrinum</i> (P ₁)	0.25 ± 0.00
	<i>Penicillium rubrum</i> (P ₂)	0.93 ± 0.73
Tryptone	<i>Aspergillus flavus</i> (A ₁)	0.25 ± 0.00
	<i>Aspergillus parasiticus</i> (A ₂)	0.20 ± 0.05
	<i>Penicillium citrinum</i> (P ₁)	0.38 ± 0.13
	<i>Penicillium rubrum</i> (P ₂)	2.40 ± 0.10
Urea	<i>Aspergillus flavus</i> (A ₁)	0.15 ± 0.00
	<i>Aspergillus parasiticus</i> (A ₂)	2.32 ± 0.03
	<i>Penicillium citrinum</i> (P ₁)	0.00 ± 0.00
	<i>Penicillium rubrum</i> (P ₂)	2.33 ± 0.08

Each value represents the mean of three replicates with standard error.

Table 2.
Effect of nitrogen sources on activity of amylase produced by isolates.

Toxigenic *A. flavus* (A₁) produced the most active α-amylase (3.25 ± 0.15 Units) and this was when ammonium sulfate was nitrogen source with starch as carbon source of the defined growth medium. Toxigenic *A. flavus* (A₁) also expressed an α-amylase activity value of 3.02 ± 0.18 Units when starch was carbon source and ammonium chloride was nitrogen source of the defined fungal growth medium (Table 2).

4. Discussion

The results of this investigation show that the toxigenic strains of *A. flavus* (A₁), *A. parasiticus* (A₂), *P. citrinum* (P₁) and *P. rubrum* (P₂) grew in a synthetic medium with varying carbon and nitrogen sources exhibiting α-amylase activities. α-Amylase activities were detected in the extracts of growth medium with rice as carbon source, infected with the toxigenic strains of *A. flavus* (A₁), *A.*

parasiticus (A₂), *P. citrinum* (P₁) and *P. rubrum* (P₂). When the carbon source was varied, potassium nitrate was the nitrogen source. When the nitrogen source was varied, starch was the carbon source for fungal growth. According to Olutiola [135], *Aspergillus chevalieri* from moldy maize produced extracellular amylase when grown in a liquid medium containing starch as carbon source. According to Barnett and Fergus [136], increasing the amount of starch-yeast extract medium increased the extracellular amylase produced by *Humicola lanuginosa*. Studies carried out by Okafor et al. [137] revealed that *Lactobacillus delbrueckii*, *Lactobacillus coryniformis* and *Saccharomyces* sp., isolated from cassava processing environments were high amylase producers. Among a series of starch sources of carbon, wheat and soluble starch were inducers of a thermostable amylase by a yeast strain isolated from starchy soil [138]. According to Bluhm and Woloshuk [139], amylopectin, an important constituent of starch, induces fumonisin B(1) production in *Fusarium verticillioides* during colonization of maize. According to Coleman [140], extracellular α-amylase was secreted by *Bacillus subtilis* in a complex medium containing maltose, starch, glycerol or glucose as carbon source; the general characteristics of secretion indicated a low but definite production of exoenzyme from the moment the cells of the organism started to grow until the end of the logarithmic phase after which, the rate of increase in cell mass decreased, the rate of enzyme secretion increased to a high linear value which was maintained even in the stationary phase.

4.1 Significance of study

Aflatoxin B1-producing-toxigenic strains of *Aspergillus flavus*, *Aspergillus parasiticus*, *Penicillium citrinum* and *Penicillium rubrum* can be explored industrially for α-amylase production using the specific growth medium and the rice medium used in this investigation. Varying the specific C and N source of this growth medium is of utmost significance in such an exploration.

4.2 Limitations

The genetic make-ups of these aflatoxin B1-producing-α-amylase-producing fungal strains are important in their ability to produce the enzyme α-amylase. Specific genes are necessary and important in the production of this enzyme. Mutant strains lacking the specific genes for α-amylase production will not be ideal in the exploration for production of the enzyme. More so, there seems to be a significant relationship between the ability to produce α-amylase and aflatoxin B1 production in mycotoxigenic fungi from literature.

5. Conclusion

The toxigenic strains of *A. flavus* (A₁), *A. parasiticus* (A₂), *P. citrinum* (P₁) and *P. rubrum* (P₂) can be explored in the industrial production of α-amylases.

Acknowledgements

Authors are thankful to the British Mycology Society (BMS), United Kingdom for Grant supports.

A. Appendix

A.1 Acetylation of cellophane tubings [134]

Material

Visking dialysis tubings (Sigma- Aldrich).

Reagents

- (i) Aqueous ethanol (50% V/V)
- (ii) Absolute ethanol
- (iii) Diethyl ether
- (iv) A mixture of benzene, acetic anhydride and pyridine in the ratio 5:4:2 (V/V)
- (v) 10% KCl (10 g of KCl in 100 ml distilled water)

Procedure

The cellophane tubings were filled with distilled water and soaked in distilled water for 24 hours. The tubings were then soaked in turn, for 30 min each time in 50% ethanol, absolute ethanol and diethyl ether successively. The tubings were thereafter soaked in the mixture of benzene, acetic anhydride, and pyridine, prepared as described above, for 18 hours. Each tubing was then properly rinsed in distilled water and stored in 10% KCl solution at 4°C until required.

A.2 Protein content determination [131]

Reagents

- (i) Reagent A—2% Na_2CO_3 in 0.1 N NaOH
- (ii) Reagent B—0.5% $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ in 1% Sodium Potassium tartrate
- (iii) Reagent C—50 ml of reagent A mixed with 1 ml of reagent B
- (iv) Folin-Ciocalteu's phenol reagent (Sigma-Aldrich Chemie GmbH, Fluka Biochemika) diluted with distilled water in the ratio 1:1(V/V). This is labeled reagent D.

Procedure

5 ml of reagent C was added to 1 ml of the test sample. This was thoroughly mixed and left at room temperature for 10 min. Thereafter, 0.5 ml of reagent D was added and allowed to remain at room temperature for 30 min. Absorbance was determined at 620 nm.

Serial dilutions of Bovine serum albumin (Sigma) were treated likewise and used to plot standard graph. The unknown protein value in each test sample is meant to be extrapolated from the standard graph.

A.3 Iodine solution

(0.3% Iodine in 3% KI)

Reagents

- (i) Iodine
- (ii) Potassium iodide (KI)

Procedure

3 g of KI was dissolved in 100 ml of warm distilled water. 0.3 g of Iodine was thereafter added and allowed to dissolve in the solution by mixing and warming.

Author details

Adekunle Odunayo Adejuwon^{1*} and Victoria Anatolyivna Tsygankova²

1 School of Health Information Management, University College Hospital, Ibadan, Nigeria

2 Department for Chemistry of Bioactive Nitrogen-Containing Heterocyclic Compounds, Institute of Bioorganic Chemistry and Petrochemistry of the National Academy of Sciences of Ukraine (NAS), Kyiv, Ukraine

*Address all correspondence to: ao_adejuwon@yahoo.ca

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Alexopoulos CJ. Introductory Mycology. New York/London: John Wiley and Sons Inc.; 1962. pp. 263-273
- [2] Streets RB. Diseases of the Cultivated Plants of the Southwest. Tucson, Arizona: The University of Arizona Press; 1969
- [3] Dutta AC. Botany for Degree Students. New Delhi: Oxford University Press; 2007. 708pp
- [4] Prescott LM, Harley JR, Klein DA. Microbiology. New York: McGraw Hill; 2005. 992pp
- [5] Tortora GJ, Funke BR, Case CL. Microbiology: An Introduction. San Francisco, California: Pearson Education Inc.; 2004. 898pp
- [6] Brock DT, Madigan MT. Biology of Microorganisms. Prentice-Hall International Inc.; 1991. 900pp
- [7] Brock DT, Madigan MT, Martinko JM, Parker J. Biology of Microorganisms. Eaglewood Cliffs, New Jersey: Prentice-Hall, Inc.; 1994. pp. 528-530
- [8] Willey JM, Linda MS, Woolverton CJ. Prescott, Harley and Klein's Microbiology. New York: McGraw Hill Companies Inc.; 2008. 1088pp
- [9] Liang XQ, Holbrook CC, Lynch RE, Guo BZ. Beta-1,3-glucanase activity in peanut seed (*Arachis hypogaea*) is induced by inoculation with *Aspergillus flavus* and copurifies with a conglutin-like protein. Phytopathology. 2005;**95**:506-511
- [10] Novas MV, Cabral D. Association of mycotoxin and sclerotia production with compatibility groups in *Aspergillus flavus* from peanut in Argentina. Plant Disease. 2002;**86**:215-219
- [11] Windham GL, Williams WP. *Aspergillus flavus* infection and aflatoxin accumulation in resistant and susceptible maize hybrids. Plant Disease. 1998;**82**:281-284
- [12] Walker RD, White DG. Inheritance of resistance to *Aspergillus* ear rot and aflatoxin production of corn from C12. Plant Disease. 2001;**85**:322-327
- [13] Windham GL, Williams WP, Davis FM. Effects of the southwestern corn borer on *Aspergillus flavus* kernel infection and aflatoxin accumulation in maize hybrids. Plant Disease. 1999;**83**:535-540
- [14] Russin JS, Guo BZ, Tubajika KM, Brown RL, Cleveland TE, Widstrom NW. Comparison of kernel wax from corn genotypes resistant or susceptible to *Aspergillus flavus*. Phytopathology. 1997;**87**:529-533
- [15] Chen ZY, Brown RL, Lax AR, Guo BZ, Cleveland TE, Russin JS. Resistance to *Aspergillus flavus* in corn kernels is associated with a 14-KDa protein. Phytopathology. 1998;**88**:276-281
- [16] Olanya OM, Hoyos GM, Tiffany LH. Waste corn as a point source of inoculum for *Aspergillus flavus* in the corn agroecosystem. Plant Disease. 1997;**81**:576-581
- [17] Norton RA. Effect of carotenoids on aflatoxin B₁ synthesis by *Aspergillus flavus*. Phytopathology. 1997;**87**:814-821
- [18] Garber RK, Cotty PJ. Formation of sclerotia and aflatoxins in developing cotton balls infected by the S strain of *Aspergillus flavus* and potential for biocontrol with an atoxigenic strain. Phytopathology. 1997;**87**:940-945
- [19] Orum TV, Bigelow DM, Cotty PJ, Nelson MR. Using predictions based on geostatics to monitor trends in

Aspergillus flavus strain composition.
Phytopathology. 1999;**89**:761-769

[20] McAlpin CE, Wicklow DT, Platis CE. Genotypic diversity of *Aspergillus parasiticus* in an Illinois corn field. Plant Disease. 1998;**82**:1132-1136

[21] Brown RL, Chen Y, Cleveland TE, Russin JS. Advances in the development of host resistance in corn to aflatoxin contamination by *Aspergillus flavus*. Phytopathology. 1999;**89**:113-117

[22] Moore KG, Price MS, Boston RS, Weissinger AK, Payne GA. A chitinase from Tex 6 maize kernels inhibits growth of *Aspergillus flavus*. Phytopathology. 2004;**94**:82-87

[23] Sirois KL, Loparco DP, Lobeer JW. Systemic infection of onion seedlings by *Aspergillus niger* and *Fusarium* sp. Phytopathology. 1999;**89**:S73. Publication No: P-1999-0520-AMA

[24] Sirois KL, Lorbeer JW, Holcomb MA. Onion seed infection levels subsequent to sequential exposure of onion seed stalk and flower parts to *Aspergillus niger*. Phytopathology. 1999;**89**:S73. Publication No.: P-1999-0521-AMA

[25] Doster MA, Michailides TJ, Morgan DP. *Aspergillus species* and mycotoxins in Figs from California orchards. Plant Disease. 1996;**80**:484-489

[26] Thompson JR, Latorre BA. Characterization of *Botrytis cinerea* from table grapes in Chile using RAPD-PCR. Plant Disease. 1999;**83**:1090-1094

[27] Shellie KC, Skaria M. Reduction of green mold on grape fruit after hot force-air quarantine treatment. Plant Disease. 1998;**82**:380-382

[28] Zhang J, Swingle PP. Effects of curing on green mold and stem-end rot of citrus fruit and its potential

application under Florida packing system. Plant Disease. 2005;**89**:834-840

[29] Conway WS, Leverentz B, Saftner RA. Survival and growth of *Listeria monocytogenes* on fresh-cut apple slices and *Penicillium expansum*. Plant Disease. 2000;**84**:177-181

[30] Spotts RA, Cervantes LA. Disease incidence-inoculum close relationships for *Botrytis cinerea* and *Penicillium expansum* and decay of pear fruit using dry, air borne conidia. Plant Disease. 2001;**85**:755-759

[31] Tian S, Fan Q, Xu Y, Liu H. Biocontrol efficacy of antagonist yeasts to gray mold and blue mold on apples and pears in controlled atmospheres. Plant Disease. 2002;**86**:848-853

[32] Meyer UM, Spotts RA. Detection and quantification of *Botrytis cinerea* by ELISA in pear stems during cold storage. Plant Disease. 2000;**84**:1099-1103

[33] Palou L, Smilanick JL, Crisostco CH, Mansour M. Effects of gaseous ozone exposure on the development of green and blue molds on cold stored citrus fruit. Plant Disease. 2001;**85**:632-638

[34] Palou L, Smilanick JL, Usall J, Vinas I. Control of post harvest blue and green molds of oranges by hot water, sodium carbonate and sodium bicarbonate. Plant Disease. 2001;**85**:371-376

[35] Smilanick JL, Mackey BE, Reese R, Usall J, Margosan DA. Influence of concentration of soda ash, temperature and immersion period on the control of post harvest green mold of oranges. Plant Disease. 1997;**81**:379-382

[36] Smilanick JL, Michael IF, Mansour MF, Mackey BE, Margosan DA, Flores D, et al. Improved control of green mold of citrus with imazalil in warm water compared with its use in wax. Plant Disease. 1997;**81**:1299-1304

- [37] Sugar D, Spotts RA. Control of postharvest decay in pear by four laboratory-grown yeasts and two registered biocontrol products. *Plant Disease*. 1999;**83**:155-158
- [38] Janisiewicz WJ, Peterson DL. Susceptibility of the stem pull area of mechanically harvested apples to blue mold decay and its control with a biocontrol agent. *Plant Disease*. 2004;**88**:662-664
- [39] Spotts RA, Holz G. Adhesion and removal of conidia of *Botrytis cinerea* and *Penicillium expansum* from grape and plum fruit surfaces. *Plant Disease*. 1996;**80**:688-691
- [40] Smilanick JL, Aiyabei J, Gabler FM, Doctor J, Sorenson D, Mackey B. Quantification of the toxicity of the aqueous chlorine to spores of *Penicillium digitatum* and *Geotrichum citri-aurantii*. *Plant Disease*. 2002;**86**:509-514
- [41] Janisiewicz WJ, Peterson DL, Yoder KS, Miller SS. Experimental bin drenching system for testing biocontrol agents to control post harvest decay of apple. *Plant Disease*. 2005;**89**:487-490
- [42] Spotts RA, Cervantes LA, Mielke EA. Variability in post harvest decay among apple cultivars. *Plant Disease*. 1999;**83**:1051-1054
- [43] Sugar D, Benbow JM, Powers KA, Basile SR. Orchard sprays on post harvest decay of pear. *Plant Disease*. 2003;**87**:1260-1262
- [44] Smilanick JL, Mansour MF, Sorenson D. Pre- and post harvest treatments to control green mold of citrus fruit during ethylene degreening. *Plant Disease*. 2006;**90**:89-96
- [45] Pianzzola MJ, Moscatelli M, Vero S. Characterization of *Penicillium* isolates associated with blue mold on apple in Uruguay. *Plant Disease*. 2004;**88**:23-28
- [46] Bohinski RC. *Modern Concepts in Biochemistry*. 4th ed. Boston/London/Sydney/Toronto: Allyn and Bacon Inc.; 1983. 531pp
- [47] Reddy NS, Nimmagadda A, Sambasiva Rao KRS. An overview of the microbial α -amylase family. *African Journal of Biotechnology*. 2003;**2**(12):645-648
- [48] Dixon M, Webb EC. *Enzymes*. London: Longmans; 1971. 950pp
- [49] Robyt JF. Enzymes in hydrolysis and synthesis of starch. In: *Starch Chemistry and Technology*. New York: Academic Press; 1984. pp. 88-90
- [50] Najafi MF, Kembhavi A. One step purification and characterization of an extracellular α -amylase from marine *Vibrio* sp. *Enzyme and Microbial Technology*. 2005;**36**(4):535-539
- [51] Rahardjo YSP, Jolink F, Haemers S, Tramper J, Rinzema A. Significance of bed porosity, bran and specific surface area in solid-state cultivation of *Aspergillus oryzae*. *Biomolecular Engineering*. 2005;**22**(4):133-139
- [52] Wu YB, Ravindran V, Pierce J, Hendriks WH. Influence of three phytase preparations in broiler diets based on wheat or corn: *In vitro* measurement of nutrient release. *International Journal of Poultry Science*. 2004;**3**(7):450-455
- [53] Ajayi AA, Adejuwon AO, Olutiola PO. Extracellular α -amylase production by the yam (*Discorea* spp.) rot organism, *Penicillium sclerotigenum* Yamamoto Science. *Focus*. 2005;**10**(3):197-203
- [54] Douzdijian V, Gugliuzza KK. The impact of midline versus transverse incisions on wound complications and outcome in simultaneous pancreas-kidney transplant: A retrospective analysis. *Transplant International*. 1996;**9**(1):62-67

- [55] Hashim SO, Delgado OD, Martinez MA, Kaul RH, Mulaa FJ, Mattiasson B. Alkaline active maltohexaose-forming α -amylase from *Bacillus halodurans* LBK 34. *Enzyme and Microbial Technology*. 2005;**36**(1):139-146
- [56] Hemker M, Stratmann A, Goeke K, Schroder W, Lentz J, Piepersberg W, et al. Identification, cloning, expression and characterization of the extracellular acarbose-modifying glycosyltransferase, AcbD from *Actinoplanes* sp. strain SE 50. *Journal of Bacteriology*. 2001;**183**(15):4484-4492
- [57] Oh KW, Kim MJ, Kim HY, Kim BY, Baik MY, Auh JH, et al. Enzymatic characterization of a maltogenic amylase from *Lactobacillus gasseri* ATCC 33323 expressed in *Escherichia coli*. *FEMS Microbiology Letters*. 2005;**252**(1):175-181
- [58] Papi RM, Chaitidou SA, Triikka FA, Kyriakidis DA. Encapsulated *Escherichia coli* in alginate beads capable of secreting a heterologous pectin lyase. *Microbial Cell Factories*. 2005;**4**:35
- [59] Lehninger AL. *Principles of Biochemistry*. New York: Worth Publishers, Inc.; 1982. 1011pp
- [60] Howling D. Mechanisms of starch enzymolysis. *International Biodeterioration*. 1989;**25**:15-19
- [61] Vihinen M, Mantsala P. Microbial amylolytic enzymes. *Critical Reviews in Biochemistry and Molecular Biology*. 1989;**24**(4):329-418
- [62] Uguru GC, Akinyanju JA, Sani A. The use of sorghum for thermostable amylase production from *Thermoactinomyces thalophilus*. *Letters in Applied Microbiology*. 1997;**25**:13-16
- [63] Ajayi AO, Fagade OE. Utilization of corn starch as substrate for β -amylase by *Bacillus* spp. *African Journal of Biomedical Research*. 2003;**6**(1):37-42
- [64] Reiss M, Heibges A, Metzger I, Hartmeier W. Determination of BOD-values of starch containing waste water by a BOD-biosensor. *Biosensors and Bioelectronics*. 1998;**13**(10):1083-1090
- [65] Ampe F, Ben Omar N, Guyot JP. Recovery of total microbial RNA from lactic acid fermented foods with a high starch content. *Letters in Applied Microbiology*. 1998;**27**(5):270-274
- [66] Doelle HW. Socio-economic microbial process strategies for a sustainable development using environmentally clean technologies: Sagopalm a renewable resource. *Livestock Research for Rural Development*. 1998;**10**(1):23-32
- [67] Rivera MH, Lopez-Munguia A, Soberon X, Saab-Rincon G. Alpha-amylase from *Bacillus licheniformis* mutants near to the catalytic site: Effects on hydrolytic and transglycosylation activity. *Protein Engineering*. 2003;**16**(7):505-514
- [68] Zangrando FEL, Yoko HE. Culture media for amylase production by toxigenic fungi. *Brazilian Archives of Biology and Technology*. 2000;**43**(5):461-467
- [69] Tanaka T, Hoshina M, Tanabe S, Sakai K, Ohtsubo S, Taniguchi M. Production of d-lactic acid from defatted rice bran by simultaneous saccharification and fermentation. *Bioresource Technology*. 2005;**97**(2):211-217
- [70] Sejedi RH, Naderi-Manesh H, Khajeh K, Ahmadvand R, Ranjbar B, Asoodeh A, et al. A Ca-independent α -amylase that is active and stable at low pH from the *Bacillus* sp. KR-8104. *Enzyme and Microbial Technology*. 2005;**36**(5-6):666-671

- [71] Rodriguez VB, Alamenda EJ, Gellegos JFM, Lopez AIG. Thermal deactivation of a commercial α -amylase from *Bacillus licheniformis* used in detergents. *Biochemical Engineering Journal*. 2006;**27**(3):299-304
- [72] Mijts B, Patel BKC. Cloning, sequencing and expression of an alpha-amylase gene, amy A, from the thermophilic halophile *Halothermothrix orenii* and purification and biochemical characterization of the recombinant enzyme. *Microbiology*. 2002;**148**:2343-2349
- [73] Shaw JF, Lin FP, Chen SC, Chen HC. Purification and properties of an extracellular α -amylase from *Thermus* sp. *Botanical Bulletin of Academia Sinica*. 1995;**36**:195-200
- [74] Yutani K. Molecular weight of thermostable α -amylase from *Bacillus stearothermophilus*. *Journal of Biochemistry*. 1973;**74**:581-586
- [75] Saito N. A thermophilic extracellular α -amylase from *Bacillus licheniformis*. *Archives of Biochemistry and Biophysics*. 1973;**155**:290-298
- [76] Manning GB, Campbell LL. Thermostable α -amylase of *Bacillus stearothermophilus*. I. Crystallization and some general properties. *The Journal of Biological Chemistry*. 1961;**236**(11):2952-2957
- [77] Huang LP, Jin B, Lant P, Zhou J. Simultaneous saccharification and fermentation of potato starch waste water to lactic acid by *Rhizopus oryzae* and *Rhizopus arrhizus*. *Biochemical Engineering Journal*. 2005;**23**:265-276
- [78] Ogashara K, Imanishi A, Isemura T. Studies on thermophilic α -amylase from *Bacillus stearothermophilus*. I. Some general and physico-chemical properties of thermophilic α -amylase. *The Journal of Biochemistry*. 1970;**74**:581-596
- [79] Synowiecki J, Grzybowska B, Zdzienbło A. Sources, properties and suitability of new thermostable enzyme in food processing. *Critical Reviews in Food Science and Nutrition*. 2006;**46**(3):197-205
- [80] Peng P, Wu J, Cheng AC, Gao QY, Zhang SZ. Cloning and expression of the alpha-amylase gene from a *Bacillus* sp. WS06 and characterization of the enzyme. *Wei Sheng Wu Xue Bao*. 2005;**45**(6):876-880
- [81] Yun J, Kang S, Park S, Yoon H, Kim MJ, Heu S, et al. Characterization of a novel amylolytic enzyme encoded by a gene from a soil-derived metagenomic library. *Applied and Environmental Microbiology*. 2004;**70**(12):7229-7235
- [82] Bessler C, Schmitt J, Maurer KH, Schmid RD. Directed evolution of a bacterial alpha-amylase: Towards enhanced pH-performance and higher specific activity. *Protein Science*. 2003;**12**(10):2141-2149
- [83] Dey S, Agarwal SO. Characterization of a thermostable alpha-amylase from a thermophilic *Streptomyces megasporus* strain SD12. *Indian Journal of Biochemistry and Biophysics*. 1999;**36**(3):150-157
- [84] Burgess-Cassler A, Imam SH, Gould JM. High-molecular weight amylase activities from bacteria degrading starch-plastic films. *Applied and Environmental Microbiology*. 1991;**57**(2):612-614
- [85] Heese O, Hansen G, Hohne WE, Korner D. A thermostable alpha amylase from *Thermoactinomyces vulgaris*. Purification and characterization. *Biomedica Biochimica Acta*. 1991;**50**(3):225-232
- [86] Akindahunsi AA. Physico-chemical studies on amylases from fermented cassava waste water. In: *Scientific*

Preprints Automized List. Trieste, Italy: The Abdus Salam International Centre for Theoretical Physics; 2001

[87] Dey G, Palit S, Banerjee R, Maiti BR. Purification and characterization of malto oligosaccharide-forming amylase from *Bacillus circulans* GRS 313. *Journal of Indian Microbiology and Biotechnology*. 2002;**28**(4):193-200

[88] Doman K, Ryu SJ, Heo SJ, Kim DW, Kim HS. Characterization of a novel carbohydrase from *Lipomyces starkeyi* KSM 22 for dental application. *Journal of Microbiology and Biotechnology*. 1999;**9**(3):260-264

[89] Jana M, Chattopadhyay DJ, Patti BR. Thermostable, high salt-tolerant amylase from *Bacillus megaterium* VUMB-109. *Acta Microbiologica et Immunologica Hungarica*. 1997;**44**(3):281-289

[90] Ohno N, Fukuda H, Wang H, Kasamura M, Shinoyama H, Fuji T. Amylases produced by a thermophilic fungus. *Thermoascus aurantiacus* and some of their properties. *Seibutsu-Kogaku Kaishi*. 1998;**76**(3):111-117

[91] Shekhar HU, Ali MM, Hossain MA. Extracellular amylase production by locally isolated mosquito-pathogenic *Bacillus sphaericus* N11. *Bangladesh Journal of Microbiology*. 1997;**14**(1-2):17-23

[92] Falih AM. Effect of heavy-metals on amylolytic activity of the soil yeasts *Geotrichum capitatum* and *Geotrichum candidum*. *Bioresource Technology*. 1998;**66**(3):213-217

[93] Bernhardsdotter ECMJ, Ng JD, Garriott DK, Pusey ML. Enzymatic properties of an alkaline, chelator-resistant alpha-amylase from an alkaliphilic *Bacillus* sp. isolate L1711. *Process Biochemistry*. 2005;**40**:2401-2408

[94] Mishra S, Noronha SB, Suraishkumar GK. Increase in enzyme

productivity by induced oxidative stress in *Bacillus subtilis* cultures and analysis of its mechanism using microarray data. *Process Biochemistry*. 2005;**40**(5):1863-1870

[95] El-Said AH, Abdel-Hafez SI, Saleem A. Effect of herbizid and touchdown of some extracellular enzymes. *Acta Microbiologica et Immunologica Hungarica*. 2005;**52**(1):105-130

[96] Figueira ELZ, Hirooka EY, Mendiola-Olanya E, Blanco-Labra A. Characterization of a hydrophobic amylase inhibitor from corn (*Zea mays*) seeds with activity against amylase from *Fusarium verticillioides*. *Phytopathology*. 2002;**93**:917-922

[97] Figueira ELZ, Ono EYZ, Mendiola-Olanya E. New amylase inhibitor present in corn seeds active *in vitro* amylase from *Fusarium verticillioides*. *Plant Disease*. 2002;**87**:233-240

[98] Hofmann O, Vertesy L, Braunitzer G. The primary structure of alpha-amylase inhibitor Z-2685 from *Streptomyces parvullus* FH-1641. Sequence homology between inhibitor and alpha-amylase. *Biological Chemistry Hoppe-Seyler*. 1985;**366**(12):1161-1168

[99] Vertesy L, Tripiet D. Isolation and structure elucidation of an alpha-amylase inhibitor, A1-3688, from *Streptomyces aureofaciens*. *FEBS Letters*. 1985;**185**(1):187-190

[100] Su YC, Chiu RJ, Yu N, Chang WR. The microbial production of amylase inhibitor and its application. I. Isolation and cultivation of *Streptomyces nigrifaciens* NTU-3314. *Proceedings of the National Science Council, Republic of China*. 1984;**B8**(4):292-301

[101] Lin J, Lu C, Yu IT, Wu JF. Effects of starch sources on the production and activity of amylase from *Bacillus subtilis*. *Journal of the*

- Agricultural Association of China. 1999;**185**:22-31
- [102] Yin XH, Gerbaud C, Francou FX, Guerineau M, Virolle MJ. Ami C, another amylolytic gene maps close to the ami B locus in *Streptomyces lividans* TK 24. Gene (Amsterdam). 1998;**215**(1):171-180
- [103] Lo HF, Lin LL, Chiang WY, Chie MC, Hsu WH, Chang CT. Deletion analysis of the C-terminal region of the alpha-amylase of *Bacillus* sp. strain TS-23. Archives of Microbiology. 2002;**178**(2):115-123
- [104] Bahl H, Burchhardt G, Spreinat A, Haeckel K, Wienecke A, Schmidt B. Alpha-amylase of *Clostridium therosulfurigenes* EMI: Nucleotide sequence of the gene, processing of the enzyme and comparison of alpha amylases. Applied and Environmental Microbiology. 1991;**57**(5):1554-1559
- [105] Kang DO, Hwang IK, Kim BY, Ahn SC, Mheen TI, Ahn JS, et al. Secretion of Bacillus alpha-amylase from yeast directed by glucoamylase I signal sequence of *Saccharomyces diastaticus*. Biochemistry and Molecular Biology International. 1996;**39**(1):181-190
- [106] Lorentz K. Routine alpha-amylase assay using protected 4-nitrophenyl-1,4-alpha-D-maltoheptaoside and a novel alpha-glucoside. Clinical Chemistry. 2000;**46**:644-649
- [107] Akpan I, Bankole M, Adesemowo AM. A rapid plate culture method for screening of amylase producing microorganisms. Biotechnology Techniques. 1999;**13**(6):411-413
- [108] Higuchi Y, Ohashi A, Imachi H, Harada H. Hydrolytic activity of alpha-amylase in anaerobic digested sludge. Water Science and Technology. 2005;**52**(1-2):259-266
- [109] Uhlig H. Industrial Enzymes and their Applications. New York: Wiley and Sons; 1998. 472pp
- [110] Pandey A, Nigman P, Soccol CR, Soccol VT, Singh D, Mohan R. Advances in microbial amylase. Biotechnology and Applied Biochemistry. 2000;**31**(2):135-152
- [111] Mellon JE, Cotty PJ, Dowd MK. *Aspergillus flavus* hydrolases: Their roles in pathogenesis and substrate utilization. Applied Environmental Microbiology and Biotechnology. 2007;**77**:497-504
- [112] Jallow EAA, Twumasi P, Mills-Robertson FC, Dumevi R. Assessment of aflatoxin-producing fungi strains and contamination levels of aflatoxin B₁ in groundnut, maize, beans and rice. Journal of Agricultural Science and Food Technology. 2018;**4**(4):71-79
- [113] Liu J, Sun L, Zhang N, Zhang J, Guo J, Li C, et al. Effects of nutrients in substrate of different grains on aflatoxin B₁ production by *Aspergillus flavus*. Biomedical Research International. 2016;**2016**:10. Article ID 7232858. <http://dx.doi.org/10.1155/2016/7232858>
- [114] Kim HR, Kim JH, Bai DH, Ahn BH. Identification and characterization of useful fungi with α -amylase activity from Korean traditional nuruk. Microbiology. 2011;**39**(4):278-282
- [115] Fakhoury A, Woloshuk CR. Amy1, the α -amylase gene of *Aspergillus flavus*: Involvement in aflatoxin biosynthesis in maize kernels. Phytopathology. 1999;**89**(1):908-914
- [116] Figueira ELZ, Hirooka EY. Culture medium for amylase production by toxigenic fungi. Brazilian Archives of Biology and Technology. 2000;**43**(5):461-467
- [117] Rajasekaran K, Sayler RJ, Majumdar R, Sickler CM, Cary JW.

Inhibition of *Aspergillus flavus* growth and aflatoxin production in transgenic maize expressing the α -amylase inhibitor from *Lablab purpureus* L. Journal of Visualized Experiments. 2019;144:e59169. DOI: 10.3791/59169

[118] Fountain JC, Scully BT, Ni X, Kemerait RC, Lee RD, Chen ZY, et al. Environmental influences on maize-*Aspergillus flavus* interactions and aflatoxin production. Frontiers in Microbiology. 2014;5:40. DOI: 10.3389/fmicb.2014.00040

[119] Espindola AS, Schneider W, Cardwell KF, Carrillo Y, Hoyt PR, Marek SM, et al. Inferring the presence of aflatoxin-producing *Aspergillus flavus* strains using RNA sequencing and electronic probes as a transcriptomic screening tool. PLoS One. 16 Oct 2018;13(10):e0198575. DOI: 10.1371/journal.pone.0198575

[120] Yu J, Mohawed SM, Bhatnagar D, Cleveland TE. Substrate-induced lipase gene expression and aflatoxin production in *Aspergillus flavus*. Journal of Applied Microbiology. 2003;95:1334-1342

[121] Smith CA, Woloshuk CP, Robertson D, Payne GA. Silencing of the aflatoxin gene cluster in a diploid strain of *Aspergillus flavus* is suppressed by ectopic aflR expression. Genetics. 2007;176(4):2077-2086

[122] Stern KR, Jansky S, Bidlack JE. Introductory Plant Biology. New York: McGraw Hill Higher Education; 2003. 624pp

[123]Sizer FS, Whitney EN. Nutrition: Concepts and Controversies. London: Wadsworth, Thomson Learning; 2000. 567pp

[124] Mauseth JD. Botany: An Introduction to Plant Biology. London: Saunders College Publishing; 1995. 794pp

[125] Leventin E, McMahon K. Plants and Society. New York: WCB McGraw Hill; 1999. 477pp

[126] Graham LE, Graham JM, Wilcox LW. Plant Biology. Upper Saddle River: Pearson Education, Inc.; 2006. 670pp

[127] Cannon PF, Kirk PM. Fungal Families of the World. Wallingford, Oxfordshire: CAB International Publishing; 2007. 4556pp

[128] Hanlin RT. Illustrated Genera of Ascomycetes. St. Paul, Minnesota: American Phytopathological Society Press; 1990. 263pp

[129] Olutiola PO, Ayres PG. Utilization of carbohydrates by *Rhynchosporium secalis*.I. Growth and sporulation on glucose, galactose and galacturonic acid. Physiologia Plantarum. 1973;29:92-96

[130] Olutiola PO, Nwaogwugwu RI. Growth, sporulation and production of maltase and proteolytic enzymes in *Aspergillus aculeatus*. Transactions of the British Mycological Society. 1982;78(1):105-113

[131] Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the folin phenol reagent. Journal of Biological Chemistry. 1951;193:265-275

[132] Pfueller SL, Elliott WH. The extracellular α -amylase of *Bacillus stearothermophilus*. Journal of Biological Chemistry. 1969;244:48-54

[133] Xiao Z, Storms R, Tsang A. A quantitative starch-iodine method for measuring alpha-amylase and glucoamylase activities. Analytical Biochemistry. 2006;351(1):146-148

[134] Whitaker DR, Hanson KR, Datta PK. Improved procedure for purification and characterization of Myrothecium cellulase. Canadian Journal of Microbiology and Physiology. 1963;41:671-696

[135] Olutiola PO. α -Amylase activity of *Aspergillus chevalieri* from mouldy maize. Indian Phytopathology. 1982;**35**(3):428-433

[136] Barnett EA, Fergus CL. The regulation of extracellular amylase mycelium and time, in some thermophilic and mesophilic *Humicola* species. Mycopathologia et Mycologia Applicata. 1971;**44**(2):131-141

[137] Okafor N, Umeh C, Ibenugbu C. Amelioration of garri, a cassava based fermented food by the inoculation of microorganisms secreting amylase, lysine and linamarase into cassava mash. World Journal of Microbiology and Biotechnology. 1998;**14**(6):835-838

[138] Fossi BT, Tavea F, Ndjouenkeu R. Production and partial characterization of a thermostable amylase from ascomycetes yeast strain isolated from starchy soils. African Journal of Biotechnology. 2005;**4**(1):14-18

[139] Bluhm BH, Woloshuk CP. Amylopectin induces fumonisin B(1) production by *Fusarium verticillioides* during colonization of maize kernels. Molecular Plant-Microbe Interactions. 2005;**18**:1333-1339

[140] Coleman G. Studies on the regulation of extracellular enzyme formation by *Bacillus subtilis*. Journal of General Microbiology. 1967;**49**:421-431