We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Scientific Evidences Supporting the Activation of the Renin-Angiotensin-Aldosterone System during Estral Cycle and Pregnancy in Mares

Katy Satué and Ana Muñoz

Abstract

In women and laboratory animals, local and circulating components of the renin-angiotensin-aldosterone system (RAAS) are related to specific reproductive functions that occur during the estrous cycle, such as folliculogenesis, ovulation, corpus luteum development, and steroidogenesis. Also, in pregnant females of these species, maternal cardiovascular and renal systems undergo intense modifications, with the aim of matching the increased energy requirements of the fetus and fetoplacental unit. Some of these changes can be the origin, and others the consequence of a new endocrine environment. The fetus and the placenta induce endocrine changes, with modifications in the protein, lipid, carbohydrate, and mineral metabolism, together with simultaneous cardiovascular changes derived from the uterine growth and its content. The participation of RAAS during this period is of vital importance to regulate these cardiovascular, hemodynamic, hematological, and metabolic adjustments imposed by pregnancy because they will have a direct influence on the correct development and viability of the fetus. In mares, our research team has been investigating the changes of RAAS in mares during the estral cycle and during pregnancy, and these results are presented in the current chapter, comparing with the data previously reported for women and laboratory animals.

Keywords: estrous cycle, mare, pregnancy, renin, angiotensin, aldosterone

1. Introduction

In nonpregnant females of various species, the components of the renin-angiotensin-aldosterone system (RAAS), i.e., prorenin, renin, angiotensin II (ANG-II), and aldosterone, are expressed in the tissues of reproductive organs, mainly the uterus and ovaries. These hormones have direct physiological relationships with specific reproductive functions, including folliculogenesis, oocyte maturation, ovulation, follicular atresia, corpus luteum development and luteolysis, steroidogenesis, angiogenesis, and expression of certain vasoactive substances [1–8].

A great body of literature has confirmed an increase in plasma activity of renin (PRA) and plasma concentrations of ANG-II and aldosterone in women during the

luteal compared to the follicular period of the estrous cycle [9–19]. The primary source of renin and aldosterone is progesterone (P4) secreted by the corpus luteum. The elevated P4 concentrations during the luteal period lead to increased renal plasma flow, glomerular filtration rate, and natriuresis. This natriuretic effect stimulates in a compensatory way the synthesis and release of renin, ANG-II, and aldosterone [10–12, 15, 16, 20–23]. However, these results do not agree with those described by Szmulowicz et al. [19], who suggested that the synthesis of aldosterone might be independent of renin and ANG-II at the ovarian level. Another physiological event that happens during the estrous cycle of the women is the pre-ovulatory increase of renin [14, 24], ANG-II [25], and aldosterone [10, 12, 25, 26].

Similar results to those described in women have been reported in laboratory animals [27–30]. A potential physiological mechanism related to these changes is the stimulatory effect exerted by estrogens (E2) on the synthesis of angiotensinogen [31]. Additional mechanisms could be the hemodynamic and renal blood flow variations, changes in Na concentrations at the macula densa in the kidney, alterations in local sympathetic activity, and release of corticotropin or adrenocorticotrophic hormone (ACTH). All of these mechanisms act as regulatory factors for aldosterone synthesis [17, 19, 32, 33].

During pregnancy, circulating and tissular RAASs interact closely in order to achieve a successful outcome of the pregnancy. The local RAASs involved in pregnancy are mainly located at the ovaries, uterus (both at the placenta and decidua), and intrarenal level. The nonrenal local RAAS systems have pivotal functions in the implantation and in the placentation as well as in the development of the uteroplacental and umbilical placental circulations. In addition, they contribute directly to the circulating RAAS of the pregnant female, determining in part the normal functionality of the cardiovascular and renal systems [34].

In this review, we summarize the state of the art of the current knowledge of the RAAS in reproductive mares. We provide comparative profiling of the hormones of this system with other species, mainly with women and laboratory animals. The results of our own investigations performed during the last 10 years have demonstrated that an activation of the RAAS happens around ovulation and during pregnancy in the reproductive mare, even though significant differences with the reports made for women and laboratory animals have been found.

2. Changes in the renin-angiotensin-aldosterone system during the estral cycle

2.1 A brief review of the estral cycle in the mare

An exhaustive review of the estral cycle of the mare and the neuroendocrine mechanisms associated has been published recently [35]. Briefly, the mean length of a mare's estral cycle is 21 ± 3 days. The cycle is divided into two physiological periods: the estrous or follicular period and the diestrus or luteal period. The duration of the follicular period is approximately 6 days, but it can take 4–10 days depending on the mare. The luteal period has a duration of 15 days, ranging between 12 and 18 days. Ovulation can occur at any time during the follicular period, although it usually occurs 24–48 h before the end of this period. During the estral cycle, physiological changes in mares include follicular development, selection of the dominant follicle, release of the oocyte from follicle, formation of the corpus luteum, and production of P4. These physiological events are controlled by gonadotropin-releasing hormone (GnRH) that stimulates the hypophysis to secrete follicle-stimulating hormone (FSH) and luteinizing hormone (LH). These hormones control the development of

the follicle and stimulate the secretion of E2 from the follicle to induce ovulation and corpus luteum establishment and development [35, 36].

2.2 Changes in the renin-angiotensin-aldosterone system during the luteal period

Most researchers have documented that aldosterone concentrations rise between 2 and 4 times during the luteal period [11, 37]. Presumably, this increase mirrors the increase experienced by ANG-II concentrations. However, this supposition has not always been supporting, and as a consequence, later it was suggested that the changes in ANG-II might be attributed to other ways independent of the RAAS [19]. Although this hypothesis is currently the most defended in the literature, some authors would not be able to confirm it [32, 38].

A variety of studies have shown that the relationship between PRA and aldosterone from the middle to the end of the luteal period can be adjusted to a linear model [9, 10, 16, 22, 37, 39, 40]. The researches performed in women with luteal insufficiency have helped to clarify complex issues, such as the involvement of the sex hormones in the modifications of the different RAAS components during the different period of the estral cycle. In this sense, several early investigations compared the concentrations of angiotensinogen, aldosterone, gonadotropins, ovarian steroids, and PRA in women with ovulatory cycles and in women with luteal insufficiency. While in physiological estral cycles the peak of PRA was closely related to the increase in aldosterone, in women with anovulatory cycles, these temporal relationships between renin and aldosterone or PRA and aldosterone were not found [9, 10, 41]. Subsequently, it was hypothesized that once the corpus luteum reaches functional maturity, it acts as the primary source of renin and aldosterone. Thereby, there was an explanation for the direct relationship between P4 and other compounds derived from the corpus luteum and the RAAS in physiological estral cycles, but not in women with ovulatory failure, because these women show a significant decrease in P4 concentrations [18, 21, 42].

P4 seems to be directly related to the luteal synthesis of aldosterone [15, 18, 21, 42]. Certain mechanisms dependent on P4, such as increased plasma flow, renal glomerular filtration rate, and Na and Cl excretion, result in natriuresis [16]. The initial natriuresis induced by P4 stimulates in a compensatory way the secretion of renin, ANG-II, and aldosterone [9–11, 20, 37]. The experimental administration of P4 results in natriuresis, which is followed by peaks of aldosterone secretion, aldosterone urinary excretion (AUE), and Na retention [11, 43]. Although PRA, ANG-II, and aldosterone increase after the administration of P4, the rises do not occur simultaneously [43]. These results have been explained in base of the competence between P4 and aldosterone for the mineralocorticoid receptor [12, 44, 45]. In physiological cycles, endogenously secreted P4 exerts a mild anti-aldosterone effect, avoiding an excessive retention of Na and water during the luteal period [46].

The E2 and P4 peaks during the luteal period lead to increases of PRA and ANG-II 2–3 times and also increases of AUE [17, 19]. However, the increase of E2 that happens during the first half of the estral cycle does not result in substantially raised PRA and AUE. Perhaps these events are mediated by the expression of aldosterone receptor protein and angiotensinogen in the absence of changes in ANG-II [46–48]. The lack of correlation between aldosterone and E2, as well as the lack of stimulation of aldosterone production in *in vitro* cultures of glomerulosa cells, could be other findings that might exclude E2 from the synthesis of aldosterone during the luteal period [19].

However, there are additional mechanisms involved in the synthesis of aldosterone independently of the P4 concentrations, such as the uptake of proteins and Na in the diet [49, 50] or a peripheral vasodilation [16]. It is difficult to interpret these

findings without considering the intake of Na, since this electrolyte determines primarily the production of aldosterone via RAAS [49]. However, many previous studies did not control or did not report Na concentrations, which is a limiting factor in the proper interpretation of the results. More recently, Szmuiłowicz et al. [19] showed that circulating and urinary levels of aldosterone increase significantly during the luteal period in response to the infusion of ANG-II in women with high Na concentrations. In these situations, positive correlations between P4 and aldosterone have been found, without parallel changes in PRA and ANG-II. On the contrary, in women with low Na concentrations, these interrelationships do not occur, even though the restriction of Na in the diet is a powerful stimulus for the activation of RAAS [49]. In Na-restricted diets, PRA and aldosterone respond independently between each other, and therefore, the absence of modifications in PRA and ANG-II could rule out that the increase in aldosterone is modulated via RAAS activation during the luteal period.

Another mechanism that has been related to the synthesis of renin and aldosterone during the luteal period is the release of angiotensinogen and prorenin. However, the studies conducted in this field have provided contradictory results. The expression of angiotensinogen is regulated transcriptionally by E2 [31], and consequently, the administration of different ovarian stimulation treatments in women increases the concentrations of E2 and renin [15]. Therefore, the activation of the RAAS during the luteal period could be partially related to the increased synthesis of angiotensinogen [21] or be a direct consequence of the vasodilatory effects of E2 [51]. However, early studies have not revealed variations in angiotensinogen during the luteal period of physiological estral cycles [10, 52, 53]. The combined use of E2 and P4 during the luteal period in ovariectomized rats resulted in increases in PRA without significant simultaneous changes in angiotensinogen [54]. The lack of a significant relationship between PRA and angiotensinogen might rule out E2 as a cause of the modifications in renin concentrations, despite its involvement in the secretion of aldosterone [10, 53, 55].

Although the main source of circulating active renin is the kidney, the ovaries appear to be the source of the cyclic increase of prorenin during the estral cycle [14, 56]. In fact, the release of LH triggers an increase of three times the concentrations of prorenin, remaining elevated during the first half of the luteal period, even though the decrease in P4 at the end of the luteal period reduces simultaneously the prorenin levels [18, 21, 42, 56].

2.3 Changes in the renin-angiotensin-aldosterone system during the follicular period

Preovulatory increases of angiotensinogen, prorenin, PRA, renin, ANG-II, and aldosterone have been reported in women [11, 14, 25, 26] and laboratory animals [29]. According to other authors, the increase in aldosterone concentrations appears to be transient [11], or it is not constant in all cycles [37].

The origin of the preovulatory aldosterone peak is unknown, but it has been speculated that the increase in E2 could exert a stimulatory effect on angiotensinogen in the liver, and this, in turn, increases PRA and aldosterone [52, 53, 55]. Other researches, on the contrary, failed to find any relationship between angiotensinogen, PRA, and aldosterone in physiological cycles [11, 17, 19, 57], ruling out some type of significant influence of endogenous E2 on PRA and AUE during the periovulatory period [9, 10, 41]. Despite these contradictory results, the hypothesis that the increase in angiotensinogen is the cause of PRA and aldosterone peaks is still commonly accepted.

A transient peak of prorenin has been described simultaneously with the elevation of gonadotropins, mainly LH, toward the moment of the ovulation. It has been

speculated that this prorenin peak will be responsible later for the increase of active renin [58]. However, prorenin and LH are not always related to active renin [14, 15], because the most striking effects of this precursor appear toward the middle of the luteal period [18, 21, 42].

Because ACTH is one the main regulators of aldosterone secretion, the pre-ovulatory peak of PRA might be related to the stimulatory effect of E2 on the adrenal gland and to the increase of the 17 α -P4-enzyme, which also shows a preovulatory peak. This enzyme catalyzes the conversion of pregnenolone into 17- α -hydroxypregnenolone, which is a precursor of the sexual steroids [59]. However, although an experimental infusion of ACTH triggers an elevation in PRA [60], the absence of changes in cortisol (CORT) concentrations in physiological cycles could refute these hypotheses [17, 19, 32]. Other factors, including hemodynamic changes in renal blood flow, local sympathetic activity, and variation in Na concentrations at the macula densa level, have also been associated with the preovulatory peak of renin and aldosterone [33, 61].

In broodmares, Satué et al. [62] evaluated the changes in circulating RAAS components around ovulation, considering the 5 days before ovulation (from day 5 to day 1), the day of the ovulation (day 0), and the 5 days immediately after ovulation (from day 1 to day 5). A summary of the main endocrine changes, including those of the RAAS, is presented in **Figure 1**.

A progressive increase in plasma renin concentrations was found before ovulation, peaking at the day of the ovulation [62] (**Figure 2**). The increase in plasma renin before ovulation in other species, as has been previously attributed to the release of LH or to the stimulating effect of human chorionic gonadotropin (hCG) [24]. Another plausible explanation for this rise in plasma renin before ovulation was the increase in E2 concentrations, since a positive correlation between both variables has been found in mares ($r = 0.744$) [62] (**Figure 3**). These results in mares agree with the data provided by Sealy et al. [15] in women with ovarian stimulation, in whom significant simultaneous increases in renin and E2 were detected. Another explanation for the rise in renin concentrations in the mares could be a mild hypovolemia associated with the presence of interstitial fluid in

Figure 1. Representative scheme of the hormonal changes that occur in the mare during the 5 days before ovulation (preovulatory period), the day of the ovulation, and the last first 5 days after ovulation (postovulatory period) (ACTH, adrenocorticotrophic hormone; E₂, estrogens; CORT, cortisol; P₄, progesterone).

Figure 2. Plasma renin concentrations during the 5 days before ovulation, the day of the ovulation, and the first 5 days after ovulation in mares (numbers indicate the days between which significant differences were found at $p < 0.05$).

Figure 3. Significant positive correlations between plasma estrogens (E2) and renin concentrations in mares before ovulation ($r = 0.744$).

abdominal cavity during ovulation. Even though changes in Na and Cl concentrations at the renal juxtaglomerular apparatus significantly influence the release of renin [63], the correlations between renin and these electrolytes appear to be low in mares before ovulation [62]. In the same way, Roussel et al. [64] described cyclic patterns of plasma Na and aldosterone concentrations in cow, but without significant correlations with the components of the RAAS.

After ovulation, a sharp decrease in plasma renin concentrations was detected in mares (**Figure 2**). Similar changes have been reported in women [17]. Ounis-Skali et al.

Figure 4.
Plasma angiotensin-II concentrations during the 5 days before ovulation, the day of the ovulation, and the first 5 days after ovulation in mares (numbers indicate the days between which significant differences were found at $p < 0.05$).

[18] attributed these results to the release of renin from the corpus luteum, reflecting a direct relationship between P4 and renin after ovulation, and similar hypotheses might be extrapolable to the mare. Two studies performed by our research team demonstrated low correlations between P4 and renin, suggesting other factors different from P4 might be considered in order to explain the non-significant trend toward an increase in renin concentrations during the first days after ovulation [65, 66].

Plasma ANG-II concentrations did not show significant variations before ovulation, but it experienced a sharp and progressive decrease after ovulation [62] (**Figure 4**). On the contrary, significant increases in ANG-II have been previously observed in sows [11] and in rats [29] before ovulation. ANG-II concentrations in women after ovulation were almost the double of the values found before ovulation [17–19]. This increase in women was attributed to the modulation of the flow in the ovarian blood vessels during oocyte maturation [15]. The reason why ANG-II concentrations decrease in the mares after ovulation, which is an opposite change to this described for women and laboratory animals, is not currently known.

Plasma aldosterone concentrations increased progressively from the fifth day before ovulation until the day of the ovulation. After this moment, aldosterone concentrations did not change and persisted to be significantly higher than before ovulation [62] (**Figure 5**). Because an increase in renin was also found the day of the ovulation in mares and ANG-II was higher during the 5 days before ovulation, it was speculated that the increased renin was the starting point to activate RAAS, resulting in increased aldosterone concentrations. This hypothesis was proven when the correlations between hormones before and after ovulation were assessed (**Figure 6**). A positive significant correlation ($r = 0.756$) was found between renin and aldosterone concentrations before ovulation. Surprisingly, ANG-II and aldosterone concentrations did not present a significant correlation before ovulation (**Figure 6**). These data might indicate a dissociation between ANG-II and aldosterone before ovulation in the mare. In this case, the synthesis of aldosterone might have occurred in addition via other pathways independent from the activation of the RAAS [62].

Figure 5. Plasma aldosterone concentrations during the 5 days before ovulation, the day of the ovulation, and the first 5 days after ovulation in mares (numbers indicate the days between which significant differences were found at $p < 0.05$).

Figure 6. Correlations between the three main components of the renin-angiotensin-aldosterone system in mares before (A) and after ovulation (B).

The relationships between the diameter of the predominant follicle and the hormones of the RAAS have also been investigated in mares [62] (**Figure 7**). A significant positive correlation was found between renin and aldosterone concentrations and the diameter of the follicle before ovulation in the mares ($r = 0.742$ and 0.804 , respectively) [62] (**Figure 7**). In other species, RAAS regulates the development and growth of the antral follicles. Receptors AT1 and AT2 for ANG-II have been found in the granulosa cells of the antral follicles and in the primordial primary and secondary follicles of sows [67]. Administration of an ANG-II inhibitor in cows avoided the growth of the predominant follicle and also induced a reduction in the concentrations of E2 [68, 69]. In the same way, in women subjected to ovarian stimulation with

Figure 7.
Correlations between the diameter of the predominant follicle (DPF) and the concentrations of renin, angiotensin II, and aldosterone before ovulation in mares.

human menopausal gonadotropin (HMG), concentrations of ANG-II were positively correlated with E2 and the diameter of the predominant follicle [70]. In our research in mares, however, correlations between the diameter of the predominant follicle and ANG-II were very low ($r = 0.366$), despite the significant correlations between this diameter and the concentrations of renin and aldosterone [62] (**Figure 7**). Our results might suggest that circulating ANG-II concentrations do not exert a transcendental effect on the development of the ovarian follicle or, alternatively, some antagonist peptides of the ANG-II might be produced before ovulation in mares [65, 66].

3. Changes in the renin-angiotensin-aldosterone system during the pregnancy

Plasma concentrations of prorenin increase by 5–10 times during the first 10 weeks of pregnancy in women. This increase is simultaneous with the increase in hCG and after, both decreased until the moment of the delivery. Although prorenin is mainly synthesized in the renal juxtaglomerular cells, there are other extrarenal sources, such as the ovary, follicular fluid, and placenta [71]. The prorenin concentrations also increase when the LH concentrations peak at the moment of the ovulation [71, 72]. The combined administration of LH and FSH causes a 390% increase in plasma prorenin, while hCG increases above 1000% [13, 73]. An early study speculated that prorenin could act as a hormone with functions independent of the active renin [13]. Until now, to the authors' best knowledge, the prorenin concentration in plasma/serum has not been quantified in mares.

A substantial increase in the circulating concentrations of angiotensinogen has also been documented during the first weeks of pregnancy in women [74]. Angiotensinogen concentrations remain elevated throughout pregnancy, peaking at term. In sheep, a bimodal pattern of secretion of this hormone has been described, with a first peak at the beginning of the pregnancy, concomitant with the period of the maximum placental growth. The second peak occurs at term, suggesting an association between plasma angiotensinogen and fetal growth [75].

The increased synthesis of angiotensinogen has been attributed to the stimulating effect of E2 at hepatic level [16, 17, 76]. However, the relationships between angiotensinogen and E2 are not always positive [77]. In nonpregnant women undergoing estrogen treatments, angiotensinogen and ANG-II increased in blood, whereas renin concentrations tended to decrease because of the negative feedback exercised by angiotensinogen [76]. However, the angiotensinogen concentrations have not been measured in mares yet.

The total concentration of circulating renin, both its active and inactive forms, increases during pregnancy in women. However, the contribution of the physiologically inactive renin to the total renin is greater than that of the active renin [78]. During the first third period of the pregnancy, renin concentrations increased between 2 and 4 times above baseline [79, 80]. After it reaches a plateau, around the fifth month of pregnancy and after that, the concentrations remained unchanged until the end of the pregnancy [81], being the main reason involved in the rise of PRA during the first and second thirds of the pregnancy, respectively [11, 80, 82–84]. Toward the end of the pregnancy, the renin concentration decreases intensely [78]. Even though the increased renin release contributes greatly to the ANG-II synthesis, the renin concentrations could decrease because of a negative feedback mechanism [15, 85].

The synthesis of renin, in addition to renal juxtaglomerular cells, can also be carried out in other extrarenal tissues, as the uterus. This organ is the priority organ of renin synthesis during pregnancy, leading to the so-called hyperreninemia of the

pregnancy [80, 86]. The renin at the level of the fetal membranes and the uterus is found predominantly in the form of prorenin, and although it is supposed that it can be released into the maternal circulation, the degree of contribution to the maternal circulating levels is unknown. Both active renin and inactive renin have been identified in the uterus, placenta, and myometrium in women and in some laboratory animals [87]. Despite this, amniotic fluid seems to be the main source of renin produced by the chorionic cells during pregnancy [74].

The placenta appears to be the major source for the conversion of angiotensin I into ANG-II, since the prorenin, active renin, angiotensin converting enzyme (ACE), and ANG-II have been identified in the chorion and in the placenta [88]. The physiological roles of these components have not been completely clarified, but it has been speculated that they might participate in the regulation of the blood flow [89–92].

The increase in renin concentrations during pregnancy has also been attributed to the loss of electrolytes as a consequence of the increase in the glomerular filtration rate as well as of the secretion of P4, due to the antagonistic effects of the hormone on aldosterone, because P4 induces natriuresis [74, 76, 93]. On the other hand, the positive influence of E2 on the synthesis of angiotensinogen has also been considered a plausible reason for the renin peak during pregnancy [76]. Surprisingly, it is possible to find increased renin concentrations without a simultaneous increase of ANG-II concentrations. In addition, the sequestration of Na and water at the uterus might be another triggering factor for the release of renin during pregnancy, as documented in pregnant bitches [94].

In pregnant mares, Satué et al. [95] described a significant increase in the renin from the seventh month of pregnancy (**Figure 8**). In pregnant women, as stated before, the increase in renin concentrations occurs earlier than in the pregnant mares [79, 80, 84]. At present, it is unknown if the origin of these differences is the different animal species or it could be due to the influence of other physiological factors. In the mare, ANG-II concentrations did not vary significantly during pregnancy (**Figure 9**), despite the changes in renin concentrations (**Figure 8**). These data might suggest that there is a dissociation between renin and ANG-II during pregnancy in the mares. In pregnant women, it has been shown that the increase in

Figure 8.
Mean (maximum and minimum indicated in bars) concentrations of renin in mares during pregnancy (significant differences between months indicated with numbers at $p < 0.05$).

Figure 9. Mean (maximum and minimum indicated in bars) concentrations of angiotensin II in mares during pregnancy.

ANG-II exerts a negative feedback effect on renin secretion [74, 85, 96]. In addition, it has been proposed that the peak of renin during pregnancy is in part associated with the release of estrogens, which act on the production of angiotensinogen in the liver [15, 76, 97]. Moreover, Satué et al. [98] found a dissociation between renin and E2, in agreement with previous results described for pregnant women [77, 99].

On the contrary to the data found for ANG-II in mares [95], most of the investigations performed in women and laboratory animals showed that during pregnancy, circulating concentrations of ANG-II increased up to two and even three times from nonpregnant values [74, 78, 100, 101]. This increase in women appears from the beginning of the pregnancy, peaking at the seventh month and persisting to be elevated until the moment of the delivery [102].

Pregnant mares showed a marked increase in circulating aldosterone concentrations during pregnancy, with mean values that exceed almost 12 times the physiological reference range for healthy adult horses [103] (**Figure 10**). Similar results have been reported in pregnant women [74, 84, 104], bitches [105–107], and some species of laboratory animals [108]. In pregnant bitches, Robb et al. [94] and in rats, Brochu et al. [106] attributed the increase in aldosterone concentrations to a higher activity of the enzyme aldosterone-synthase and to an increased synthesis of mRNA in the cells of the zona glomerulosa, these cells being the only production site, without fetal and/or placental participation, on the contrary of what appears to happen in women [78].

In mares, our team has found lower aldosterone concentrations during the second month of pregnancy, compared to the fifth, sixth, and seventh months of pregnancy (**Figure 10**). The highest values were observed in the fifth month of pregnancy. In the eighth, ninth, and tenth months of pregnancy, aldosterone concentrations were significantly lower than the mean values found in the fifth month. In pregnant women, it has been speculated that the increased release of aldosterone would occur in an attempt to conserve Na and water to favor the expansion of the fetoplacental unit. In this way, the adequate supply of nutrients to the fetus, the maintenance of the ideal oxygen tension for fetal development, as well as the homeostasis and appropriate blood pressure between the mother and the fetus would be guaranteed [108, 109].

Figure 10. Mean (maximum and minimum indicated in bars) concentrations of aldosterone in mares during pregnancy (significant differences between months indicated with numbers at $p < 0.05$).

Although the evolution of aldosterone concentrations during pregnancy is similar in women, laboratory animals, and mares, there are temporary differences in the moment in which the peaks of concentrations are reached. In women and laboratory animals, the maximum increase in aldosterone appears in the second third of pregnancy, followed by a decrease in the third period of pregnancy. In the mare, on the contrary, as shown in **Figure 10**, the maximum concentrations were found in the fifth month.

The increase in aldosterone during pregnancy in women has been attributed, firstly, to the increase in the sensitivity of the cells of the adrenal gland to the increased synthesis and release of renin [74, 110] and, secondly, to the increase in ANG-II concentrations, a finding that has been associated with an increase in placental lactogen [74]. In women, it is known that placental lactogen plays a regulatory role in metabolic homeostasis, triggering an increase in ANG-II receptors during the second period of the pregnancy [111, 112]. It has also been speculated in women that the increase in the concentrations of aldosterone during pregnancy could be a decisive physiological event in order to prevent the massive natriuresis that could arise from an enhanced glomerular filtration rate. In these cases, aldosterone shows a physiological action opposite to the natriuretic effect of P4 at the level of the distal convoluted tubule, avoiding excessive loss of Na and allowing its gradual accumulation in the fetoplacental and in the extracellular maternal fluids [113]. However, the antagonism between both mineralocorticoids, i.e., aldosterone and P4, during pregnancy does not seem to be constant [114, 115].

4. Conclusions

Our results obtained in healthy reproductive mares demonstrated that there is an activation of the RAAS around the time of ovulation, as indicated by the increased plasma renin and aldosterone concentrations. After ovulation, the rapid decrease in plasma renin and ANG-II concentrations might indicate a modulation of the previously activated RAAS, even though plasma aldosterone concentrations increased during this period, suggesting a dissociation of the components of this system.

In the pregnant mare, there is also an increased activity of various components of the RAAS. Pregnant mares show a progressive increase in circulating renin, with fluctuating changes in aldosterone, peaking at fifth month of pregnancy and without significant changes in ANG-II. The reason for the apparent dissociation between the three main hormones of the RAAS at determined moments of the pregnancy remains unclear. The data obtained in pregnant mares regarding the RAAS seems to indicate that renin is a more intense stimulus for the increased synthesis of aldosterone than ANG-II.

Acknowledgements

The authors would like to express their gratitude to the owners who have allowed, over a period of more than 10 years, the participation of their mares in the studies carried out by our research team.

Conflict of interest

The authors declare that they have no conflicts of interest.

Author details

Katy Satué¹ and Ana Muñoz^{2*}

1 University Cardenal Herrera-CEU, Valencia, Spain

2 University of Córdoba, Spain

*Address all correspondence to: ksatue@uch.ceu.es

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Schauser KH, Nielsen AH, Dantzer V, Poulsen K. Angiotensin-converting enzyme activity in the bovine uteroplacental unit changes in relation to the cycle and pregnancy. *Placenta*. 2001;**22**:852-862. DOI: 10.1053/plac.2001.0740
- [2] Brunswing-Spickenheier B, Mukhopadhyay AK. Local regulatory factors in regulation of ovarian function: Role of prorenin-renin-angiotensin-system. *Indian Journal of Experimental Biology*. 2003;**41**(7):669-681
- [3] Ferreira R, Oliveira J, Fernandes R, Moraes J, Gonçalves P. The role of angiotensin II in the early stages of bovine ovulation. *Reproduction*. 2007;**134**:713-719. DOI: 10.1530/REP-07-0239
- [4] Hurley ME, Critelli M, Johnson A, DenHaese D, Jones C, Olivieri MP. Renin immunofluorescence in the metestrus rat uterus. *The FASEB Journal*. 2009;**23**(1):637.9-637.9. DOI: 10.1096/fasebj.23.1_supplement.637.9
- [5] Pereira VM, Reis FM, Santos RA, Cassali GD, Santos SH, Honorato-Sampaio K, et al. Gonadotropin stimulation increases the expression of angiotensin (1-7) and MAS receptor in the rat ovary. *Reproductive Sciences*. 2009;**16**(12):1165-1174. DOI: 10.1177/1933719109343309
- [6] Pountain SJ, Pipkin FB, Hunter MG. The ontogeny of components of the renin-angiotensin system in the porcine fetal ovary. *Animal Reproduction Science*. 2010;**117**(1-2):119-126. DOI: 10.1016/j.anireprosci.2009.03.006
- [7] Herr D, Duncan WC, Hack G, Konrad R, Kreienberg R, Wulff C. Regulated expression of the renin-angiotensin-system in human granulosa lutein cells: Angiotensin II increases VEGF expression but its synthesis is reduced by hCG. *Archives of Gynecology and Obstetrics*. 2010;**281**(3):409-416
- [8] Sneeringer R, Penzias AS, Barrett B, Usheva A. High levels of mineralocorticoids in preovulatory follicular fluid could contribute to oocyte development. *Fertility and Sterility*. 2011;**95**(1):182-187. DOI: 10.1016/j.fertnstert.2010.06.028
- [9] Sundsfjord JA, Aakvaag A. Plasma angiotensin II and aldosterone excretion during the menstrual cycle. *Acta Endocrinologica*. 1970;**64**:452-458
- [10] Sundsfjord JA, Aakvaag A. Plasma rennin activity, plasma rennin substrate and urinary aldosterone excretion in the menstrual cycle in relation to the concentration of progesterone and oestrogens in the plasma. *Acta Endocrinologica*. 1972;**71**:519-529
- [11] Sundsfjord JA, Aakvaag A. Variations in plasma aldosterone and plasma rennin activity throughout the menstrual cycle, with especial reference to the pre-ovulatory period. *Acta Endocrinologica*. 1973;**73**:499-508
- [12] Oelkers W, Schoneshofer M, Blumel A. Effects of progesterone and four synthetic progestogens on sodium balance and the renin-aldosterone system in man. *The Journal of Clinical Endocrinology and Metabolism*. 1974;**39**(5):882-890. DOI: 10.1210/jcem-39-5-882
- [13] Sealey JE, Atlas SA, Glorioso N, Manapat H, Laragh JH. Cyclical secretion of prorenin during the menstrual cycle: Synchronization with luteinizing hormone and progesterone. *PNAS*. 1985;**82**:8705-8709. DOI: 10.1073/pnas.82.24.8705
- [14] Sealey JE, Cholst I, Glorioso N, Troffa C, Weintraub ID, James G, et al. Sequential changes in plasma luteinizing

hormone and plasma prorenin during the menstrual cycle. *The Journal of Clinical Endocrinology and Metabolism*. 1987;**65**(1):1-5. DOI: 10.1210/jcem-65-1-1

[15] Sealey JE, Itskovitz-Eldor J, Rubattu S, James G, August P, Thaler I, et al. Estradiol and progesterone related increases in the renin aldosterone system: Studies during ovarian stimulation and early pregnancy. *The Journal of Clinical Endocrinology and Metabolism*. 1994;**79**(1):258-264. DOI: 10.1210/jcem.79.1.8027239

[16] Chapman AB, Zamudio S, Woodmansee W, Merouani A, Osorio F, Johnson A, et al. Systemic and renal hemodynamic changes in the luteal phase of the menstrual cycle mimic early pregnancy. *American Journal of Physiology-Renal Physiology*. 1997;**273**(5):777-782

[17] Chidambaram M, Duncan J, Lai V, Cattran D, Floras J, Scholey J, et al. Variation in the rennin angiotensin system throughout the normal menstrual cycle. *Journal of the American Society of Nephrology*. 2002;**13**:446-452

[18] Ounis-Skali N, Mitchell GF, Solomon CG, Solomon SD, Seely EW. Changes in central arterial pressure waveforms during the normal menstrual cycle. *Journal of Investigative Medicine*. 2006;**54**(6):321-326. DOI: 10.2310/6650.2006.05055

[19] Szmuiłowicz E, Adler G, Williams J, Green D, Yao T, Hopkins P, et al. Relationship between aldosterone and progesterone in human menstrual cycle. *The Journal of Clinical Endocrinology and Metabolism*. 2006;**91**(10):3981-3987. DOI: 10.1210/jc.2006-1154

[20] Oelkers W, Berger V, Bolik A, Bahr V, Hazard B, Beier S, et al. Dihydrospirorenone, a new progestogen

with antimineralocorticoid activity: Effects on ovulation, electrolyte excretion, and the renin-aldosterone system in normal women. *The Journal of Clinical Endocrinology and Metabolism*. 1991;**73**(4):837-842. DOI: 10.1210/jcem-73-4-837

[21] Oelkers W. Effects of estrogens and progestogens on the rennin-aldosterone system and blood pressure. *Steroids*. 1996;**61**(4):166-171

[22] Hirshoren N, Tzoran I, Makrienko I, Edoute Y, Plawner MM, Itskovitz-Eldor J, et al. Menstrual cycle effects on the neurohumoral and autonomic nervous systems regulating the cardiovascular system. *The Journal of Clinical Endocrinology and Metabolism*. 2002;**87**(4):1569-1575. DOI: 10.1210/jcem.87.4.8406

[23] Nakamura T, Okamura N, Yagi M, Omatsu H, Yamamori M, Kuwahara A, et al. Effects of ABCB1 3435C>T genotype on serum levels of cortisol and aldosterone in women with normal menstrual cycles. *Genetics and Molecular Research*. 2009;**8**(2):397-403

[24] Loret de Mola JR, Goldfarb JM, Hecht BR, Babbo CJ, Friedlander MA. Gonadotropins induce higher active renin levels in the follicular fluid of normal and hyperstimulated cycles. *Gynecological Endocrinology*. 1999;**13**:155-160

[25] Weir RJ, Doig A, Fraser R, Morton JJ, Parboosing J, Robertson JJ, et al. Studies of the renin-angiotensin-aldosterone system, cortisol, DOC, and ADH in normal and hypertensive pregnancy. *Perspectives in Nephrology and Hypertension*. 1976;**5**:251-261

[26] Frölich JC, Hollifield JW, Dormois JC, Frölich BL, Seyberth H, Michelakis AM, et al. Suppression of plasma renin activity by indomethacin in man. *Circulation Research*. 1976;**39**:447-452. DOI: 10.1161/01.RES.39.3.447

- [27] Howard RB, Pucell AG, Bumpus FM, Husain A. Rat ovarian renin: Characterization and changes during the estrous cycle. *Endocrinology*. 1988;**123**(5):2331-2340. DOI: 10.1210/endo-123-5-2331
- [28] De Vito E, Cabrera RR, Guardia DC. Changes during lactation in the renin-like enzyme concentration in rat luteal tissue. *Acta Endocrinologica*. 1988;**116**(4):526-530
- [29] De Vito E, Guardia DC, Cabrera RR. Cyclical changes in plasma renin during the oestrous cycle in rats: Synchronized effect of oestrogen and progesterone. *The Journal of Endocrinology*. 1989;**121**(2):261-267
- [30] Harewood W, Gillin A, Mohamed S, Willis N, Lazzaro V, Duggin G, et al. Cyclical changes in the rennin-angiotensin-aldosterone system during the menstrual cycle of the baboon (*Papio hamadryas*). *Journal of Medical Primatology*. 1996;**25**(4):267-271
- [31] Gordon MS, Chin WW, Shupnik MA. Regulation of angiotensinogen gene expression by estrogen. *Journal of Hypertension*. 1992;**10**:361-366
- [32] Pechere-Bertschi A, Maillard M, Stalder H, Brunner H, Burnier M. Renal segmental tubular response to salt during the normal menstrual cycle. *Kidney International*. 2002;**61**:425-431. DOI: 10.1046/j.1523-1755.2002.00158.x
- [33] Sealey JE, Laragh JH. Plasma renin activity enzyme-kinetic assay: Protection of angiotensin I from bacterial degradation. *Clinical Chemistry*. 2011;**57**(3):529-530. DOI: 10.1373/clinchem.2010.156596
- [34] Lumbers ER, Pringle KG. Roles of the circulating renin-angiotensin-aldosterone system in human pregnancy. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*. 2014;**306**(2):91-101. DOI: 10.1152/ajpregu.00034.2013
- [35] Satué K, Gardón JC. A review of the estrous cycle and the neuroendocrine mechanisms in the mare. *Journal of Steroids and Hormonal Science*. 2013;**4**:115. DOI: 10.4172/2157-7536.1000115
- [36] McKinnon AO, Squires EL, Vaala WE, Dickson DV. *Equine Reproduction*. 2nd ed. Ames, IA: Wiley Blackwell; 2011. pp. 3288-3299. ISBN: 978-0-813-81971-6
- [37] Katz FH, Romfh P. Plasma aldosterone and renin activity during the menstrual cycle. *The Journal of Clinical Endocrinology and Metabolism*. 1972;**34**:819-822. DOI: 10.1210/jcem-34-5-819
- [38] Parry BL, Javeed S, Laughlin GA, Hauger R, Clopton P. Cortisol circadian rhythms during the menstrual cycle and with sleep deprivation in premenstrual dysphoric disorder and normal control subjects. *Biological Psychiatry*. 2000;**48**(9):920-931
- [39] Michelakis AM, Yoshida H, Dormois JC. Plasma rennin activity and plasma aldosterone during the normal menstrual cycle. *American Journal of Obstetrics and Gynecology*. 1975;**123**(7):724-726
- [40] Stachenfeld NS, DiPietro L, Kokoszka CA, Silva C, Keefe DL, Nadel ER. Physiological variability of fluid-regulation hormones in young women. *Journal of Applied Physiology*. 1999;**86**:1092-1096. DOI: 10.1152/jappl.1999.86.3.1092
- [41] Sundsfjord JA. Plasma renin activity and aldosterone excretion during prolonged progesterone administration. *Acta Endocrinologica*. 1971;**67**(3):483-490
- [42] Garcia MJ, Martinez-Martos JM, Mayas MD, Carrera MP, De la Chica S,

Cortes P, et al. Hormonal status modifies renin-angiotensin system regulating aminopeptidases and vasopressin-degrading activity in the hypothalamus-pituitary-adrenal axis of female mice. *Medicinal Chemistry*. 2008;**4**:336-347

[43] Stachenfeld NS, Taylor HS. Oestrogen effects on urine concentrating response in young women. *The Journal of Physiology*. 2003;**1**(3):869-880. DOI: 10.1113/jphysiol.2003.046920

[44] Myles K, Funder JW. Progesterone binding to mineralocorticoid receptors: In vitro and in vivo studies. *The American Journal of Physiology*. 1996;**270**(4):601-607. DOI: 10.1152/ajpendo.1996.270.4.E601

[45] Quinkler M, Meyer B, Bumke-Vogt C, Grossmann C, Gruber U, Oelkers W, et al. Agonistic and antagonistic properties of progesterone metabolites at the human mineralocorticoid receptor. *European Journal of Endocrinology*. 2002;**146**(6):789-799

[46] Keam S, Wagstaff A. Ethinylestradiol/drospirenone. A review of its use as an oral contraceptive. *Treatments in Endocrinology*. 2002;**2**(1):1-23

[47] Nowaczynski W, Murakami T, Richardson K, Genest J. Increased aldosterone plasma protein binding in women on combined oral contraceptives throughout the menstrual cycle. *Journal of Clinical Endocrinology and Metabolism*. 1978;**47**(1):193-199. DOI: 10.1210/jcem-47-1-193

[48] Williamson PM, Buddle ML, Brown MA, Whitworth JA. Ambulatory blood pressure monitoring (ABPM) in the normal menstrual cycle and in women using oral contraceptives. Comparison with conventional blood pressure

measurement. *American Journal of Hypertension*. 1996;**9**(1):953-958. DOI: 10.1016/0895-7061(96)00150-1

[49] Adler GK, Moore TJ, Hollenberg NK, Williams GH. Changes in adrenal responsiveness and potassium balance with shifts in sodium intake. *Endocrine Research*. 1987;**13**(4):419-445

[50] Woods LL. Mechanisms of renal vasodilation after protein feeding: Role of the renin-angiotensin system. *The American Journal of Physiology*. 1993;**264**(3):601-609. DOI: 10.1152/ajpregu.1993.264.3.R601

[51] Mccaffrey TA, Czaja JA. Diverse effects of estradiol-17 beta: Concurrent suppression of appetite, blood pressure and vascular reactivity in conscious, unrestrained animals. *Physiology & Behavior*. 1989;**45**(3):649-657

[52] Skinner SL, Lumbers ER, Symonds EM. Alteration by oral contraceptives of normal menstrual changes in plasma renin activity, concentration and substrate. *Clinical Science*. 1969;**36**(1):67-76

[53] Katz FH, Romfh P, Smith P. Diurnal variation of plasma aldosterone, cortisol and renin activity in supine man. *Journal of Endocrinology and Metabolism*. 1975;**40**:125-134. DOI: 10.1210/jcem-40-1-125

[54] Ojeda N, Grigore D, Robertson E, Alexander B. Estrogen protects against increased blood pressure in postpubertal female growth restricted offspring. *Hypertension*. 2007;**50**:679-685. DOI: 10.1161/HYPERTENSIONAHA.107.091785

[55] Katz FH, Kappas A. The effects of estradiol and estriol on plasma levels of cortisol and thyroid hormone-binding globulins and on aldosterone and cortisol secretion rates in man. *The Journal of Clinical Investigation*. 1967;**46**(11):1768-1777. DOI: 10.1172/JCI105667

- [56] Sealey JE, Parra D, Rosenstein R, Laragh JH. Effective plasma renin activity: A derived measure for assessing residual plasma renin activity in patients taking angiotensin-converting enzyme inhibitors or angiotensin receptor blockers. *Hypertension*. 2010;**55**(3):16-17. DOI: 10.1161/HYPERTENSIONAHA.109.148023
- [57] Kaulhausen H, Leyendecker G, Benker G, Breuer H. The relationship of the renin-angiotensin-aldosterone system to plasma gonadotropin, prolactin, and ovarian steroid patterns during the menstrual cycle. *Archiv für Gynäkologie*. 1978;**225**:179-200
- [58] Bouhnnik J, Feherentz JA, Galen FX, Seyer R, Evin G, Castro B, et al. Immunologic identification of both plasma and human renal inactive renin as prorenin. *The Journal of Clinical Endocrinology and Metabolism*. 1985;**60**(2):399-401. DOI: 10.1210/jcem-60-2-399
- [59] Strott CA, West CD, Nakagawa K, Kondo T, Tyler FH. Plasma 11-deoxycorticosteroid and ACTH response to metyrapone (plasma metyrapone test). *The Journal of Clinical Endocrinology and Metabolism*. 1969;**29**(1):6-11. DOI: 10.1210/jcem-29-1-6
- [60] Kem DC, Gomez-Sanchez C, Kramer NJ, Holland OB, Higgins JR. Plasma aldosterone and renin activity response to ACTH infusion in dexamethasone-suppressed normal and sodium-depleted man. *The Journal of Clinical Endocrinology and Metabolism*. 1975;**40**(1):116-124. DOI: 10.1210/jcem-40-1-116
- [61] Yang S, Bae L, Zhang L. Estrogen increases eNOS and NOx release in human coronary artery endothelium. *Journal of Cardiovascular Pharmacology*. 2000;**36**(2):242-247
- [62] Satué K, Montesinos P, Muñoz A. Activation of the renin-angiotensin-aldosterone system in mares around ovulation. *Czech Journal of Animal Science*. 2017;**62**(6):259-267. DOI: 10.17221/32/2016-CJAS
- [63] Muñoz A, Riber C, Trigo P, Castejón-Riber C, Castejón FM. Dehydration, electrolyte imbalances and renin-angiotensin-aldosterone-vasopressin axis in successful and unsuccessful endurance horses. *Equine Veterinary Journal*. 2010;**38**:83-90. DOI: 10.1111/j.2042-3306.2010.00211.x
- [64] Roussel JD, Clement TJ, Aranas TJ. Changes of aldosterone in blood serum of dairy cattle during estrous cycle. *Journal of Dairy Science*. 1983;**66**:1734-1747. DOI: 10.3168/jds.S0022-0302(83)81999-7
- [65] Satué K, Montesinos P, Gardón JC. Association between aldosterone and cortisol levels during ovulatory period in Spanish purebred mares. *Reproduction, Fertility, and Development*. 2014;**26**:146. DOI: 10.1071/RDv26n1Ab65
- [66] Satué K, Gardon JC, Montesinos P. Relationship between oestradiol-17beta and renin concentrations during preovulatory period in Spanish purebred mares. *Reproduction in Domestic Animals*. 2012;**47**(3):111. DOI: 10.1111/j.1439-0531.2012.02044.x
- [67] Shuttleworth G, Hunter MG, Robinson G, Brouhton Pipkin F. Immunocytochemical localization of angiotensin II receptor subtypes 1 and 2 in the porcine fetal, prepubertal and postpubertal ovary. *Journal of Anatomy*. 2002;**201**:267-274. DOI: 10.1046/j.1469-7580.2002.00091.x
- [68] Ferreira R, Gasperin B, Rovani M, Santos J, Barreta M, Bohrer R, et al. Angiotensin II signaling promotes follicle growth and dominance in cattle. *Endocrinology*. 2011;**152**(12):4957-4965. DOI: 10.1210/en.2011-1146

- [69] Ferreira R, Gasperin B, Santos J, Rovani M, Santos RA, Gutierrez K, et al. Angiotensin II profile and mRNA encoding RAS proteins during bovine follicular wave. *Journal of the Renin-Angiotensin-Aldosterone System*. 2011;**12**(4):475-482. DOI: 10.1177/1470320311403786
- [70] Jarry H, Meyer B, Holzapfel G, Hinney B, Kuhn W, Wuttke W. Angiotensin II/III and substance P in human follicular fluid obtained during IVF: Relation of the peptide content with follicular size. *Acta Endocrinologica*. 1988;**119**:277-282
- [71] Vrtacnik-Bokal E, Meden-Vrtovec H, Verdenik I. Uterine arterial blood flow and the substances of ovarian renin-angiotensin system in women with polycystic ovaries undergoing in vitro fertilization. *European Journal of Obstetrics & Gynecology and Reproductive Biology*. 2006;**124**(1):77-81
- [72] Vrtacnik-Bokal EV, Vrotec HM, Virant Klun I, Verdenik I. Prolonged HCG action affects angiogenic substances and improves follicular maturation, oocyte quality and fertilization competence in patients with polycystic ovary syndrome. *Human Reproduction*. 2005;**20**:1562-1568. DOI: 10.1093/humrep/deh789
- [73] Itskovitz J, Sealey JE, Glorioso N, Rosenwaks Z. Plasma prorenin response to human chorionic gonadotropin in ovarian-hyperstimulated women: Correlation with the number of ovarian follicles and steroid hormone concentrations. *Proceedings of the National Academy of Sciences of the United States of America*. 1987;**84**(20):7285-7289. DOI: 10.1073/pnas.84.20.7285
- [74] Alhenc-Gelas F, Tache A, Saint-Andre JP, Milliez J, Sureau C, Corvol P, et al. The renin-angiotensin system in pregnancy and parturition. *Advances in Nephrology from the Necker Hospital*. 1986;**15**:25-33
- [75] Dandrea J, Cooper S, Ramsay M, Keller-Woods M, Broughton Pipkin F, Symonds M, et al. The effects of pregnancy and maternal nutrition on the maternal renin angiotensin system in sheep. *Experimental Physiology*. 2002;**87**(3):353-359
- [76] Oelkers W. The renin-aldosterone system and drospirenone. *Gynecological Endocrinology*. 2002;**16**:83-87
- [77] Weinberger MH, Kramer NJ, Grim CE, Petersen LP. The effect of posture and saline loading on plasma renin activity and aldosterone concentration in pregnancy, non-pregnant and oestrogen treated women. *The Journal of Clinical Endocrinology and Metabolism*. 1977;**44**:69-77. DOI: 10.1210/jcem-44-1-69
- [78] Carr B, Gant N. The endocrinology of pregnancy induced hypertension. *Clinics in Perinatology*. 1983;**10**:737-761
- [79] Tapia HR, Johnson CE, Strong CG. Renin angiotensin system in normal and in hypertensive disease of pregnancy. *Lancet*. 1972;**2**:847-850. DOI: 10.1016/S0140-6736(72)92211-8
- [80] Skinner SL. The renin system in fertility and normal human pregnancy. In: Robertson JIS, Nicolls MG, editors. *The Renin Angiotensin System*. London: Gower Medical Publishing; 1993. pp. 1-16
- [81] Langer B, Grima M, Coquard C, Bader AM, Schlaeder G, Imbs JL. Plasma active renin, angiotensin I, and angiotensin II during pregnancy and in preeclampsia. *Journal of Obstetrics and Gynaecology*. 1998;**91**:196-202
- [82] Weinberger MH, Kramer NJ, Petersen LP, Cleary RE, Young PC. Sequential changes in the renin-angiotensin-aldosterone systems and

plasma progesterone concentration in normal and abnormal human pregnancy. *Perspectives in Nephrology and Hypertension*. 1976;5:263-269

[83] Sullivan C, Matrin J. Sodium and pregnancy. *Clinical Obstetrics and Gynecology*. 1994;37:558-573

[84] Bentley-Lewis MD, Graves SW, Seely EW. The renin aldosterone response to stimulation and suppression during normal pregnancy. *Hypertension in Pregnancy*. 2005;24:1-16. DOI: 10.1081/PRG-45765

[85] Broughton Pipkin F, Morrison R, O'Brien PMS. The effect of prostaglandin E1 upon the pressor and hormonal response to exogenous angiotensin II in human pregnancy. *Clinical Science*. 1987;72:351-357

[86] Lindheimer MD, Katz AI. Renal physiology and disease in pregnancy. In: Seldin DW, Giebisch G, editors. *The Kidney: Physiology and Pathophysiology*. New York: Raven; 1992. pp. 3371-3432

[87] Skinner SL, Lumbers ER, Symonds EM. Renin concentration in human fetal and maternal tissues. *American Journal of Obstetrics and Gynecology*. 1968;101:529-533

[88] Kalenga MK, Thomas K, de Gasparo M, De Hertogh R. Determination of renin, angiotensin converting enzyme and angiotensin II levels in human placenta, chorion and amnion from women with pregnancy induced hypertension. *Clinical Endocrinology (Oxf)*. 1996;44:429-433

[89] Pringle KG, Tadros MA, Callister RJ, Lumbers ER. The expression and localization of the human placental prorenin/renin-angiotensin system throughout pregnancy: Roles in trophoblast invasion and angiogenesis? *Placenta*. 2011;32:956-962. DOI: 10.1016/j.placenta.2011.09.020

[90] Lumbers ER, Wang Y, Delforce SJ, Corbisier de Meaultsart C, Logan PC, Mitchell MD, et al. Decidualisation of human endometrial stromal cells is associated with increased expression and secretion of prorenin. *Reproduction of Biological Endocrinology*. 2015;13:129. DOI: 10.1186/s12958-015-0127-8

[91] Pepin E, Dehboneh SS, Raguema N, Esfandarani MT, Lavoie JL. Role of the renin-angiotensin system in healthy and pathological pregnancies, renin-angiotensin system—Past, present and future. In: Anna Naidenova Tolekova. *IntechOpen*; 2017. DOI: 10.5772/66748

[92] Khlestova GV, Romanov AY, Nizyaeva NV, Karapetyan AO, Baev OR, Ivanets TY. Dynamics of renin, angiotensin II, and angiotensin (1-7) during pregnancy and predisposition to hypertension-associated complications. *Bulletin of Experimental Biology and Medicine*. 2018;165:438-439. DOI: 10.1007/s10517-018-4188-5

[93] Jeyabalan A, Conrad KP. Renal function during normal pregnancy and preeclampsia. *Frontiers in Bioscience*. 2007;12:2425-2437

[94] Robb CA, Davis JO, Johnson JA, Blaine EH, Schneider EG, Baumber JS. Mechanisms regulating the renal excretion of sodium during pregnancy. *The Journal of Clinical Investigation*. 1970;49:871-880. DOI: 10.1172/JCI106306

[95] Satué K, Domingo R. Longitudinal study of the renin angiotensin aldosterone system in purebred Spanish broodmares during pregnancy. *Theriogenology*. 2011;75(7):1185-1194. DOI: 10.1016/j.theriogenology.2010.11.029

[96] Laragh JH, Sealey JE. The plasma renin test reveals the contribution of body sodium-volume content (V) and renin-angiotensin (R) vasoconstriction

- to long-term blood pressure. *American Journal of Hypertension*. 2011;**24**(11):1164-1180. DOI: 10.1038/ajh.2011.171
- [97] Brosnihan AL. Systemic and uteroplacental renin-angiotensin system in normal and pre-eclamptic pregnancies. *Therapeutic Advances in Cardiovascular Disease*. 2008;**2**:349-362. DOI: 10.1177/1753944708094529
- [98] Satué K, Domingo R, Redondo JI. Relationship between progesterone, oestrone sulphate and cortisol and the components of renin angiotensin aldosterone system in Spanish purebred broodmares during pregnancy. *Theriogenology*. 2011;**76**(8):1404-1415. DOI: 10.1016/j.theriogenology.2011.06.009
- [99] Weir RJ, Brown JJ, Fraser R, Lever AF, Logan RW, McIlwaine GM, et al. Relationship between plasma renin, renin substrate, angiotensin II, aldosterone and electrolytes in normal pregnancy. *The Journal of Clinical Endocrinology and Metabolism*. 1975;**40**:108-115. DOI: 10.1210/jcem-40-1-108
- [100] Symonds EM. The renin angiotensin system in pregnancy. *Obstetrics & Gynecology Annual*. 1981;**10**:45-67
- [101] Van Dijk DJ, Boner G, Giler S, Erman A. Increased serum angiotensin converting enzyme activity and plasma angiotensin II levels during pregnancy and postpartum in the diabetic rat. *JRAAS*. 2001;**2**:193-198. DOI: 10.3317/jraas.2001.027
- [102] Godard C, Gaillard R, Vallotton MB. The renin angiotensin aldosterone system in mother and fetus at term. *Nephron*. 1976;**17**:353-360. DOI: 10.1159/000180741
- [103] McKeever KH, Hinchcliff KW, Schmall LM, Reed SM, Lamb DR, Muir W III. Plasma renin activity, aldosterone, and vasopressin during incremental treadmill exercise in horses. *American Journal of Veterinary Research*. 1992;**53**:1290-1293
- [104] McCance DR, McKnight JA, Traub AI, Sheridan B, Roberts G, Atkinson AB. Plasma atrial natriuretic factor levels during normal pregnancy and pregnancy complicated by diabetes mellitus and hypertension. *Journal of Human Hypertension*. 1990;**4**:31-35
- [105] Brochu M, Gauvin JP, St. Louis J. Increase of aldosterone secretion in adrenal cortex suspensions derived from pregnant rats. *Proceedings of the Society for Experimental Biology and Medicine*. 1996;**212**:147-152
- [106] Brochu M, Lehoux JG, Picard S. Effects of gestation on enzymes controlling aldosterone synthesis in the rat adrenal. *Endocrinology*. 1997;**138**:2354-2358. DOI: 10.1210/endo.138.6.5198
- [107] Brochu M, Roy-Clavel E, Picard S, St-Louis J. In vivo regulation of enzymes controlling aldosterone synthesis in pregnant rats. *The Journal of Endocrinology*. 1998;**24**:575-579
- [108] Abou-Samra AB, Pugeat M, Dechaud H. Increased plasma concentration of N terminal beta lipotrophin and unbound cortisol during pregnancy. *Clinical Endocrinology*. 1984;**20**:221-228
- [109] Jensen E, Wood CH, Keller-Wood M. The normal increase in adrenal secretion during pregnancy contributes to maternal volume expansion and fetal homeostasis. *Journal of the Society for Gynecologic Investigation*. 2002;**9**:362-371
- [110] Wilson M, Morganti AA, Zervoudakis I, Letcher RI, Romns BM, Von Oeyon P, et al. Blood pressure, the renin aldosterone system and sex

steroids throughout normal pregnancy.
The American Journal of Medicine.
1980;**68**:97-104

[111] Petit A, Guillon G, Tence M, Jard S, Gallo-Payet N, Bellabarba D, et al. Angiotensin II stimulates both inositol phosphate production and human placental lactogen release from human trophoblastic cells. The Journal of Clinical Endocrinology and Metabolism. 1989;**69**(2):280-286

[112] West CA, Sasser JM, Baylis C. The enigma of continual plasma volume expansion in pregnancy: Critical role of the renin-angiotensin-aldosterone system. American Journal of Physiology. Renal Physiology. 2016;**311**:1125-1134. DOI: 10.1152/ajprenal.00129.2016

[113] Hayslett JP. A new role for progesterone: An agonist for mineralocorticoid receptor activation and pregnancy-related hypertension. The American Journal of Kidney Diseases. 2001;**38**:893-895. DOI: 10.1053/ajkd.2001.27723

[114] Bay W, Ferris T. Factors controlling plasma renin and aldosterone during pregnancy. Hypertension. 1979;**1**:410-415

[115] Smeaton TC, Andersen GJ, Fulton JS. Study of aldosterone levels in plasma during pregnancy. The Journal of Clinical Endocrinology and Metabolism. 1977;**44**:1-7. DOI: 10.1210/jcem-44-1-1