We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Defence against Oxidative Stress and Insecticides in *Musca domestica*

Tan Yong Hao, Siti Nasuha Hamzah and Zazali Alias

Abstract

This review is looking at the way *Musca domestica* defends itself against harmful molecules. One of the most notable enemies is against oxidative stress. Over the years there were reports that indicated the development of resistance on range of pesticides that are used against the flies. Researches have demonstrated that there are several functional protein molecules which contribute directly or indirectly as a response to oxidative stress and resistance against insecticides. As currently, the whole genome sequencing of the organisms has enabled future study to be conducted in evaluating the behaviour of the targeted protein/enzyme in response to oxidative stress and intake of insecticides in the flies.

Keywords: *Musca domestica*, oxidative stress, insecticide resistance

1. Introduction

An estimated 150,000 of species of Diptera have been described [1], and houseflies (*Musca domestica*) are one of the wonderfully evolved organism. A notorious vector, houseflies are associated with more than 100 pathogens [2], and resistance towards insecticides of houseflies have been reported all over the world. According to Scott et al., *Musca domestica* is suitable as a model organism for resistance studies and development of new insecticides. The knowledge on cellular metabolism in recent years has been expanded to understand the metabolic aspect of oxidative stress. In *Musca domestica* alone, a few families of proteins have been more or less associated with oxidative stress response: glutathione S-transferases (GST) [3–5], superoxide dismutase (SOD) [6] and glutathione peroxidases [7–9].

2. *M. domestica* response towards insecticide

Naturally houseflies' main ecosystem role is to decompose and recycle organic material. Houseflies are synanthropic insect in urban areas where high densities of human waste are their food source [10, 11]. It has been known to be vectors of various diseases of over 30 bacteria, protozoan, viruses and helminth eggs [12]. It also transfers viruses such as polioviruses [13] and *Coxsackie* viruses, as well as numerous bacteria such as *Campylobacter jejuni*, *Helicobacter pylori* [14], *Salmonella* sp. [14], *Listeria* sp., *Yersinia pseudotuberculosis* [15], *Shigella* sp. [16], *Escherichia coli* [17], and *Vibrio* sp. [13]. Flies may also be vectors of protozoan flies such as *Giardia* and

Entamoeba [16] and eggs of several tapeworms [18]. In 2010, there were further proof on transmission of Newcastle disease virus (NDV—*Paramyxoviridae*), a highly infectious virus shed in the faeces in infectious birds [19] with *Musca domestica* as vector in both field and laboratory. More recently, *Musca domestica* were also reported to carry antibiotic-resistant bacteria such as methicillin-resistant *Staphylococcus* and ticarcillin-resistant *Pseudomonas* [20], which possess threat in hospitals and health-care facilities [18]. Flies are causing 6 million cases of childhood blindness each year (<http://www.who.int/topics/trachoma/en/>). *Musca domestica* also create implications in economical ways, and costs of pesticides were estimated at more than US\$200 million yearly in the United States [21] and US\$1.6 million in 1998 [22].

The types of insecticides used to control houseflies on field are adulticides and larvicides (www.flycontrol.novartis.com). Adulticides are carbamates (e.g. propoxur and methomyl), organophosphates (e.g. fenitrothion, azamethiphos and dimethoate), pyrethroids (e.g. cyfluthrin, deltamethrin and permethrin) and recently neonicotinoids (e.g. imidacloprid, thiamethoxam). Larvicides are insect growth regulators (IGRs) (e.g. triflumuron, diflubenzuron, cyromazine [23], and novaluron and juvenile hormone synthetic analogues (e.g. methoprene, fenoxycarb, pyriproxyfen [23] (www.flycontrol.novartis.com)). Since the first case of DDT resistance is reported on the housefly [24], resistance of adult *Musca domestica* towards various insecticides in various sites (agricultural, wild and urban) is a fast-growing global issue. There has been an increasing resistance profile report from various places in the world.

In the United Kingdom, a resistance risk assessment done by [25] showed that although farmers claimed they had reduced using insecticides (a measure to reduce selective stress on field housefly strains), there was no sign of decrease of housefly resistance towards piperonyl butoxide synergized pyrethrins. Flies with high fenitrothion and dimethoate resistance were also discovered in Denmark [26]. In 1997, an increase in pyrethroid-resistant strains and widespread of azamethiphos-resistant strains in 21 different farms all over Denmark were confirmed [27]. In Argentina, a first insecticide survey was reported [28]. Several *Musca domestica* populations were found to be permethrin-, dichlorovinyl dimethyl phosphate (an organophosphate)- and cyromazine-resistant. In the neighbouring Brazil, [29] led a first evaluation of cyromazine resistance of houseflies in five different sites, and three out of the five sites indicated cyromazine resistance. There was a report suggesting the occurrence of insecticide tolerance in tsunami-hit villages in India [30]. With hygiene at minimum provision, immediate fly control was imposed by spraying 76% dichlorvos, and LD₉₀ of adult housefly was 3.5–3.9 times higher than the flies from control sites. As in the United States, in a study tested against nine insecticides, the fly strains showed high resistances in tetrachlorvinphos, permethrin and cyfluthrin [31], while in southeastern Nebraska, houseflies are shown to be moderately resistant to permethrin yet extremely resistant to stirofos and methoxychlor [32]. Deltamethrin-resistant flies were discovered in urban garbage dump of cities of Beijing, Tianjin and Zhangjiakou [33].

In Malaysia, [34] resistance of housefly from a garbage dump, poultry farm and agricultural farm was evaluated. It was shown that garbage dump and poultry farm fly samples were more resistant than agricultural farm. It was also shown that two poultry farms in the state of Penang against malathion, propoxur and DDT, with resistance ratio, have been found with strong correlations against relative humidity, which is a first in field discovery [35]. However, on housefly larvae, resistance assessment has been relatively scarce with only a handful of feeding and toxicity tests done. A report on an increase in diflubenzuron resistance and new-found cyromazine resistant strain was also obtained [36]. A dip test-emergence test of *Musca domestica* third instar larvae on eucalyptol extracts has been done [37] with LD₅₀ values of 118 mg/fly and 177 mg/fly on male and female flies, respectively.

3. Impact of oxidative stress-induced resistance

In an oxidative stress-induced insecticide resistance research, rats [38–40], humans [41], fresh water fish *Brycon cephalus* [42] and black tiger shrimp *Penaeus monodon* [43] have been used as models to investigate insecticide inflicted oxidative stress. Insecticides including pyrethroids [44, 45], organophosphates [46–48] and organochlorines [49] have known to be inducing oxidative stress. It was reported that there were changes in activities of the antioxidative enzymes such as superoxide dismutase, catalase, glutathione peroxidase and glutathione reductase and in GSH level changes both in the liver and erythrocyte homogenate [39]. Molecular resistances are consisted of target site resistance and metabolic-based resistance [50]. Yet, most of the works, as far as *Musca domestica* is concerned, have been more in top-down approach. While genome sequencing was still ongoing for that time being, specific gene family is identified and sequenced before getting into expression studies. With other fly species such as the dipteran *Drosophila melanogaster* and *Anopheles* genome as comparable reference database, it was also concluded three groups of gene superfamilies are involved in metabolic-based resistance [51], i.e. glutathione S-transferases, cytochrome P450 and acetylcholinesterase.

In cytochrome P450, [52] it was revealed that three P450 genes, CYP4D4v2, CYP4G2, and CYP 6A38, were up-regulated in response to permethrin treatment on permethrin-resistant ALHF strains. By using PCR technology, constant overexpression of CYP 6A1, CYP 6D1 and CYP 6D3 in neocotinoid-resistant strains in Denmark during thiomethoxam challenge was demonstrated [53]. CYP6D1 was also found to be implicating more than 5000-fold of cypermethrin resistance in Learn pyrethroid-resistant strain found in New York [54]. Significant increase in non-specific esterases and glutathione S-transferases activities were also evaluated [34]. A remarkable drop on GST activity has been reported on a DDT-resistant strain 698ab [27]. Point mutation was reported as the cause of insecticide sensitivity in the case of acetylcholinesterases (E.C 3.1.1.7) [55]. As far as metabolic-based resistance is concerned, there are still much more questions to be addressed. A study [31] stated that there is very little knowledge about the mechanism of the pyrethroid resistance (monooxygenase/CYP450), although pathways have been elucidated via genomic means. There was a significant correlation between *kdr* allele (i.e. genes reducing the sensitivity of the nervous system to pyrethroids) frequencies and the levels of knockdown resistance by deltamethrin via a PCR-based assay [33]. It was also demonstrated that a behavioural resistance might be playing a role in contributing such resistance and such traits are still being inherited in the field [25]. The upregulation mediated by changes to transacting factors reveals that these mechanisms were underlying in some cases of resistances of P450, GSTs, and acetylcholinesterases [56, 57].

3.1 Enzymatic removal of cellular hydrogen peroxide

It was suggested that aerobic organisms survive due to their evolved antioxidant capability [58]. Catalase (EC 1.11.1.6) was discovered in tobacco extracts [59]. Catalase detoxifies H_2O_2 into water and oxygen [60]. Catalase is one of the well-described enzymes, and it is a class of enzyme including the iron-heme enzyme, catalase-peroxidases and a small group of manganese enzymes [61]. Superoxide dismutase (EC 1.15.1.1) is a well-known enzyme against oxidative stress. SOD1, the first superoxide dismutase to be identified, uses free radical as a substrate [62]. A metalloenzyme, superoxide dismutase catalyses the dismutation of superoxide anion ($O_2^{\cdot-}$) to hydrogen peroxide and oxygen, as the first defence line against oxidative stress [63]. They are also known to exhibit additional peroxidase activity when hydrogen peroxide level is at its large. It has been suggested that the removal of superoxide

anion will reduce SOD alternate toxic behaviour [6]. Copper-zinc and manganese SODs scavenge and dismutate superoxide anion in mitochondrial electron transport systems. It was demonstrated that a manganese superoxide dismutase-deficient yeast thrived in hyperoxia conditions (95% oxygen, 5% carbon dioxide) under the removal of electron transport system [64]. A copper-zinc SOD1 in baker's yeast was characterized at the intermembrane space of mitochondria [65].

Glutathione peroxidase (EC 1.11.1.9) utilizes reduced glutathione (GSH) to decompose hydrogen peroxide [66–68]. This enzyme was discovered [66] and identified as selenocysteine enzymes at first [69], better known as GPx1. Later, more selenocysteines were identified such as GPxs-GPx2, GPx3, and GPx4 [70]. It was also found in mammals [68, 71]. Later, a catalytic cysteine residue on rat was discovered [72], known as GPx5, and followed by GPx6 [73] which is a selenocysteine proteins in humans but not in rats or mice [74]. Mammalians GPx7 and GPx8 were the last to be elucidated but have a low GPx activity [75].

Peroxiredoxins (EC 1.11.1.15) are another group of enzymes worth mentioning when discussing about oxidative stress in cellular organisms. Peroxiredoxins are a family of antioxidant enzymes [76]. Highly specific in reducing hydrogen peroxide [77], its cysteine residue makes up the active site of peroxiredoxins, which in turn are being oxidized into sulfenic acid and recycled back to thiol, via sulfiredoxins [78]. They also control cytokine-induced peroxide levels which, in turn, mediate signal transduction in mammalian cells [79].

4. Oxidative stress-related proteins in *Musca domestica*

There are several possible candidates of oxidative stress defence proteins. Those are superoxide dismutase, catalase, glutathione peroxidase, glutathione S-transferases, GSSG reductase, thiol transferases, gamma-glutamylcysteine synthetase, and glucose-6-phosphate dehydrogenase. Oxidative stress hypothesis is evident on aging and has always been raising questions from researchers. *Musca domestica* [80], *Drosophila melanogaster* [81, 82] and *Caenorhabditis elegans* [83] are made as model tested on hyperoxia conditions. Aging is resulted from oxidative damage from cellular macromolecules [81]. It was stated that the main prediction of this hypothesis is that the rate of aging cannot be slowed down without corresponding attenuation of oxidative damage/stress [84].

GST gene family and their isoforms have been discovered to participate in oxidative stress pathway. Overexpression and peroxidase activity of GSTs in peroxide treatment were observed [85]. Other than oxidative stress resistance, GSTs detoxify xenobiotics, protect from tissue damage, participate in Jun-kinase signaling pathway and act as non-catalytic carrier proteins (ligandins) in the intracellular transport of hydrophobic compounds [3–5]. Glutathione synthetase (GSHs) are responsible in the antioxidant defence as the dominant non-protein sulphhydryls in the cell [86] forming conjugates non-enzymatically or more by the catalysis and mediation of GSTs. H_2O_2 oxidizes thiolate group in cysteine residues (-S-) into thiols (-SOH), which is present in the exposing active site. Reaction against peroxidants is also energy-consuming due to the inhibition of oxidative phosphorylation [87] and deprives energy to maintain the recycling of NADPH during pentose phosphate pathway and glucose 6-phosphate dehydrogenase, making cells hyperglycaemic [88] and able to topple the condition of cell redox levels in levels of lactate/pyruvate ratio [89]. Most of the cases above were investigated towards organophosphates and pyrethroids. In cadmium ion treatment, concentration ranging from 0.2 to 5 mM in the medium, widely known to enhance reactive oxygen species in cell, increases the levels of superoxide dismutase [90]. Lowering the intake of selenium

via diet increases the events of a peroxidative injury. The group further purified the selenium-independent glutathione peroxidase [8] and suggested this enzyme and the related pathways should be in the picture during the investigation of insect antioxidant defence system. There was no direct research work on peroxiredoxins with relation to houseflies, and its mechanisms and activities *in vivo* are not much of knowledge. However, it was discovered that there was no increase in catalase activity even though the diet of selenium in *Musca domestica* was lowered [7]. Another investigation [9] in houseflies revealed that the total inhibition of catalase also did not affect the survival of the flies, although slight increase in the level of SOD activity was observed.

5. Conclusion

Despite such remarkable immunity and rising insecticide tolerance exhibited by *Musca domestica*, and being such prominence as model for biochemistry and insect physiology, no genome project has been launched till 2009 [2]. More importantly, to the best of our knowledge, only a handful of *Musca domestica*-related proteomic work has been reported. However, in this last 5 years, there is an increasing interest unravelling the inner molecular workings of this insect. A genome project was launched [2], and a full genome of *Musca domestica* was successfully sequenced [91]. The sequenced genome is 691 MB, and some gene sequences notably 771 putative immune-related, 86 CYP450-related, and 33 glutathione S-transferase and 92 are predicted to have esterase activities. In comparison, this genome contained a plethora of shared and novel sequences than its *Drosophila* counterparts, supporting the fact of an exemplary ability of *Musca domestica* of associating closely with numerous amounts of pathogens living in a septic environment. Pioneering transcriptomic works have been done on *Musca domestica* larvae, by massive cDNA parallel pyrosequencing [92]. Thus with the help of recent advancement, a better insight on the mechanisms that are associated with oxidative stress and resistance against insecticides in *Musca domestica* is better understood.

Acknowledgements

The work has been funded by the University of Malaya Postgraduate Research Fund (PPP: PV091/2011A) and Ministry of Higher Education under the Fundamental Research Grant (FRGS: FP052-2014A).

IntechOpen

Author details

Tan Yong Hao¹, Siti Nasuha Hamzah² and Zazali Alias^{1,3*}

1 Faculty of Science, Institute of Biological Sciences, University of Malaya, Kuala Lumpur, Malaysia

2 School of Biological Sciences, Universiti Sains Malaysia, Pulau Pinang, Malaysia

3 University of Malaya Halal Research Centre (UMHRC), University of Malaya, Kuala Lumpur, Malaysia

*Address all correspondence to: alias@um.edu.my

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Thompson FC. Biosystematic Database of World Diptera. 2004. Retrieved from: www.diptera.org
- [2] Scott JG, Liu N, Kristensen M, Clark AG. A case for sequencing the genome of *Musca domestica* (Diptera: Muscidae). *Journal of Medical Entomology*. 2009;**46**(2):175-182
- [3] Ketterer B. Glutathione S-transferases and prevention of cellular free radical damage. *Free Radical Research*. 1998;**28**:647-668
- [4] Strange RC, Jones PW, Fryer AA. Glutathione S-transferase: Genetics and role in toxicology. *Toxicology Letters*. 2000;**15**:357-363
- [5] Yin Z, Ivanov VN, Habelhah H, Tew K, Ronai Z. Glutathione S-transferase elicits protection against H₂O₂-induced cell death via coordinated regulation of stress kinases. *Cancer Research*. 2000;**60**:4053-4057
- [6] den Hartog GJM, Haenen GRMM, Vegt E, van der Vijgh WJF, Bast A. Superoxide dismutase: The balance between prevention and induction of oxidative damage. *Chemico-Biological Interactions*. 2003;**145**:33-39
- [7] Simmons TW, Jamail IS, Lockshin RA. The effect of selenium deficiency on peroxidative injury in the house fly, *Musca domestica*: A role for glutathione peroxidase. *FEBS Letters*. 1987;**218**(2):251-254
- [8] Simmons TW, Jamail IS, Lockshin RA. Selenium-independent glutathione peroxidase activity associated with glutathione S-transferase from the housefly. *Musca domestica*. *Comparative Biochemistry and Physiology*. Part B: *Comparative Biochemistry*. 1989;**94**(2):323-332
- [9] Allen RG, Farmer KJ, Sohal RS. Effect of catalase inactivation on levels of inorganic peroxides, superoxide dismutase, glutathione, oxygen consumption and life span in adult houseflies (*Musca domestica*). *Biochemical Journal*. 1983;**216**(2):503-506
- [10] Dahlem G. House fly (*Musca domestica*). In: Resh V, Carde R, editors. *Encyclopedia on Insects*. 1st ed. Vol. 1. San Diego: CA Academic Press; 2003. pp. 532-534
- [11] Marshall S. *Insects: Their Natural History and Diversity*. Buffalo, New York: Firefly Books Ltd; 2006
- [12] Greenberg B, Kowalski JA, Klowden MJ. Factors affecting the transmission of salmonella by flies: Natural resistance to colonization and bacterial interference. *Infection and Immunity*. 1970;**2**:800-809
- [13] Kettle DS. Muscidae (houseflies, stableflies). In: Kettle DS, editor. *Medical and Veterinary Entomology*. Wallingford: CAB International; 1990. pp. 223-240
- [14] Grubel P, Hoffman JS, Chong FK, Burstein NA, Mepani C, Cave D. Vector potential of houseflies (*Musca domestica*) for *Helicobacter pylori*. *Journal of Clinical Microbiology*. 1997;**35**:1300-1303
- [15] Zurek L, Denning SS, Schal C, Watson DW. Vector competence of *Musca domestica* (Diptera: Muscidae) for *Yersinia pseudotuberculosis*. *Journal of Medical Entomology*. 2001;**38**:333-335
- [16] Ugbogu OC, Nwachukwu NC, Ogbuagu MN. Isolation of *Salmonella* and *Shigella* species from houseflies

(*Musca domestica* L.) in Utru, Nigeria. The African Journal of Biotechnology. 2006;5:1090-1091

[17] Szalanski AL, Owens CB, Mckay T, Steelman CD. Detection of *Campylobacter* and *Escherichia coli* O157:H7 from filth flies by polymerase chain reaction. Medical and Veterinary Entomology. 2004;18:241-246

[18] Graczyk TK, Knight R, Gilman RH, Cranfield MR. The role of non biting flies in the epidemiology of human infectious diseases. Microbes and Infection. 2001;3:231-235

[19] Barin A, Arabkhazaeli F, Rahbari S, Madani SA. The housefly, *Musca domestica*, as a possible mechanical vector of Newcastle disease virus in the laboratory and field. Medical and Veterinary Entomology. 2010;24:88-90

[20] Boulesteix G, Le Dantec P, Chevalier B, Dieng M, Niang B, Diatta B. Role of *Musca domestica* in the transmission of multiresistant bacteria in the centres of intensive care setting in a sub-Saharan Africa. Annales Francaises d'Anesthésie et de Réanimation. 2005;24:361-365

[21] Geden CJ, Arends JJ, Axtell RC, Barnard DR, Gaydon DM, Hickie LA, Hogstette JA, Jones WF, Mullens BA, Nolan Jr. MP, Petersen JJ, and Sheppard DC. Economic significance of poultry. In: Geden CJ, and Hogstette JA, editors, Research and extension needs for integrated pest management for arthropods of veterinary importance: Proceedings of a Workshop in Lincoln; 12-14 April 1994; Nebraska Gainesville, Florida: Center for Medical, Agricultural, and Veterinary Entomology USDA-ARS; pp. 1-328

[22] Crespo DC, Lecuona RE, Hogstette JA. Biological control: An important component in integrated management of *Musca domestica* (Diptera: Muscidae) in caged layer poultry houses in Buenos Aires. Biological Control. 1998;13:16-24

[23] Kočiřová A, Petrovský M, Toporčák J, Novák P. The potential of some insect growth regulators in housefly (*Musca domestica*) control. Biologia. 2004;59:661-668

[24] Saccà GS. Estenza di *mosche domestiche* resistenti al DDT. Rivista di Parassitologia. 1947;8:127-128

[25] Learmount J, Chapman P, Macnicoll A. Impact of an insecticide resistance strategy for house fly (Diptera: Muscidae) control in intensive animal units in the United Kingdom. Journal of Economic Entomology. 2002;95(6):1245-1250

[26] Keiding J. The House Fly—Biology and Control. Geneva: World Health Organization; 1986

[27] Kristensen M. Glutathione S-transferase and insecticide resistance in laboratory strains and field populations of *Musca domestica*. Journal of Economic Entomology. 2005;98(4):1341-1348

[28] Acevedo GR, Zapater M, Toloza AC. Insecticide resistance of house fly, *Musca domestica* (L.) from Argentina. Parasitology Research. 2009;105:489-493

[29] Pinto MC, Prado APD. Resistance of *Musca domestica* L. populations to cyromazine (insect growth regulator) in Brazil. Memórias do Instituto Oswaldo Cruz. 2001;96(5):729-732

[30] Srinivasan R, Jambulingam P, Gunasekaran K, Boopathidoss P. Tolerance of housefly, *Musca domestica* L. (Diptera: Muscidae) to dichlorvos (76% EC) an insecticide used for fly control in the tsunami-hit coastal villages of southern India. Acta Tropica. 2008;105:187-190

[31] Scott JG, Alefantis TG, Kaufman PE, Rutz DA. Insecticide resistance in house flies from caged-layer poultry

facilities. Pest Management Science. 2000;**56**:147-153

[32] Marçon PCRG, Thomas GD, Siegfried BD, Campbell JB, Skoda SR. Resistance status of house flies (Diptera: Muscidae) from Southeastern Nebraska beef cattle feedlots to selected insecticides. Journal of Economic Entomology. 2003;**96**(3):1016-1020

[33] Cao XM, Song FL, Zhao TY, Dong YD, Sun CHX, Lui BL. Survey of deltamethrin resistance in house flies (*Musca domestica*) from urban garbage dumps in northern China. Environmental Entomology. 2006;**35**(1):1-9

[34] Nazni WA, Ursula MP, Lee HL, Sadiyah I. Susceptibility of *Musca domestica* L. (Diptera: Muscidae) from various breeding sites to commonly used insecticides. Journal of Vector Ecology. 1999;**24**(1):54-60

[35] Bong LJ, Zairi J. Temporal fluctuations of insecticides resistance in *Musca domestica* Linn. (Diptera: Muscidae) in Malaysia. Tropical Biomedicine. 2010;**27**(2):317-325

[36] Kristensen M, Jespersen JB. Larvicide resistance in *Musca domestica* (Diptera: Muscidae) populations in Denmark and establishment of resistant laboratory strains. Journal of Economic Entomology. 2003;**96**(4):1300-1306

[37] Sukontason KL, Boonchu N, Sukontason K, Choochote W. Effects of eucalyptol on housefly (Diptera: Muscidae) and blow fly (Diptera: Calliphoridae). Revista do Instituto de Medicina Tropical de São Paulo. 2004;**46**(2):97-101

[38] Lukaszewicz-Hussain A, Moniuszko-Jakoniuk J. Activities of superoxide dismutase and catalase in erythrocytes and concentration of malondialdehyde in serum of rats

intoxicated with chlorfenvinphos in low doses. Polish Journal of Environmental Studies. 1999;**8**:234-236

[39] Lukaszewicz-Hussain A, Moniuszko-Jakoniuk J. Organophosphate insecticide chlorfenvinphos affects enzymatic and nonenzymatic antioxidants in erythrocytes and serum of rats. Polish Journal of Environmental Studies. 2003;**12**:417-423

[40] El-Demerdash FM. Lipid peroxidation, oxidative stress and acetylcholinesterase in rat brain exposed to organophosphate and pyrethroid insecticides. Food and Chemical Toxicology. 2011;**49**:1346-1352

[41] Ranjbar A, Solhi H, Mashayekhi FJ, Susanabdi A, Rezaie A, Abdollahi M. Oxidative stress in acute human poisoning with organophosphorus insecticides; a case control study. Environmental Toxicology and Pharmacology. 2005;**20**:88-91

[42] Monteiro DA, Rantin FT, Kalinin AL. The effects of selenium on oxidative stress biomarkers in the freshwater characid fish matrinxã, *Brycon cephalus* (Günther, 1869) exposed to organophosphate insecticide Folisuper 600 BR® (methyl parathion). Comparative Biochemistry and Physiology (Part C). 2009;**149**:40-49

[43] Dorts J, Silvestrea F, Huynh TT, Tybergheina AE, Nguyen TP, Kestemonta P. Oxidative stress, protein carbonylation and heat shock proteins in the black tiger shrimp, *Penaeus monodon*, following exposure to endosulfan and deltamethrin. Environmental Toxicology and Pharmacology. 2009;**28**:302-310

[44] Giray B, Gürbay A, Hincal F. Cypermethrin-induced oxidative stress in rat brain and liver is prevented by vitamin E or allopurinol. Toxicology Letters. 2001;**118**:139-146

- [45] Gupta A, Nigam D, Gupta A, Shukla GS, Agarwal AK. Effect of pyrethroid-based liquid mosquito repellent inhalation on the blood-brain barrier function and oxidative damage in selected organs of developing rats. *Journal of Applied Toxicology*. 1999;**19**:67-72
- [46] Akhgari M, Abdollahi M, Kebryaezadeh A, Hosseini R, Sabzevari O. Biochemical evidence for free radical-induced lipid peroxidation as a mechanism for subchronic toxicity of malathion in blood and liver of rats. *Human and Experimental Toxicology*. 2003;**22**:205-211
- [47] Fortunato JJ, Agostinho FR, Réus GZ, Petronilho FC, Dal-Pizzol F, Quevedo J. Lipid peroxidative damage on malathion exposure in rats. *Neurotoxicity Research*. 2006;**9**(1):23-28
- [48] Lukaszewicz-Hussain A. Role of oxidative stress in organophosphate insecticide toxicity—Short review. *Pesticide Biochemistry and Physiology*. 2010;**98**(2):145-150
- [49] Koner BC, Banerjee BD, Ray A. Organochlorine pesticide-induced oxidative stress and immune suppression in rats. *Indian Journal of Experimental Biology*. 1998;**36**(4):395-398
- [50] Perry T, Batterham P, Daborn PJ. The biology of insecticidal activity and resistance. *Insect Biochemistry and Molecular Biology*. 2011;**41**:411-422
- [51] Ranson H, Claudinos C, Ortelli F, Abgrall C, Hemingway J, Sharakova MV, et al. Evolution of supergene families associated with insecticide resistance. *Science*. 2002;**298**:179-181
- [52] Zhu F, Li T, Zhang L, Liu N. Co-up-regulation of three P450 genes in response to permethrin exposure in permethrin resistant house flies, *Musca domestica*. *BMC Physiology*. 2008;**8**:18
- [53] Kristensen M, Spencer AG, Jespersen JB. The status and development of insecticide resistance in Danish populations of the housefly *Musca domestica* L. *Pest Management Science*. 2001;**57**(1):82-89
- [54] Scott JG, Georgiou GP. Rapid development of high-level permethrin resistance in a field-collected strain of housefly (Diptera: Muscidae) under laboratory selection. *Journal of Economic Entomology*. 1985;**78**:316-319
- [55] Fournier D. Mutations of acetylcholinesterase which confer insecticide resistance in insect populations. *Chemico-Biological Interactions*. 2001;**157-158**:257-261
- [56] Feyereisen R. Insect P450 enzymes. *Annual Review of Entomology*. 1999;**44**:507-533
- [57] Hemingway J. The molecular basis of two contrasting metabolic mechanisms of insecticide resistance. *Insect Biochemistry and Molecular Biology*. 2000;**30**:1009-1015
- [58] Halliwell B, Gutteridge JMC. *Free Radicals in Biology and Medicine*. 4th ed. Oxford: Clarendon Press; 2006
- [59] Loew O. A new enzyme of general occurrence in organisms. *Science*. 1900;**11**(279):701-702
- [60] Keilin D, Hartree EF. On the mechanism of the decomposition of hydrogen peroxide by catalase. *Proceedings of the Royal Society of London - Series B: Biological Sciences*. 1938;**124**(837):397-405
- [61] Nicholls P. Classical catalase: Ancient and modern. *Archives of Biochemistry and Biophysics*. 2012;**525**:95-101
- [62] McCord JM, Fridovich I. Superoxide dismutase: An enzymic function for erythrocuprein. The

Journal of Biological Chemistry.
 1969;**224**(22):6049-6055

[63] Fridovich I. Superoxide radical and superoxide dismutases. Annual Review of Biochemistry. 1995;**64**:97-112

[64] Guidot DM, McCord JM, Wright RM, Repine JE. Absence of electron transport (Rho 0 state) restores growth of a manganese-superoxide dismutase-deficient *Saccharomyces cerevisiae* in hyperoxia. Evidence for electron transport as a major source of superoxide generation in vivo. The Journal of Biological Chemistry. 1993;**268**:26699-26703

[65] Sturtz LA, Diekert K, Jensen LT, Lill R, Culotta VC. A fraction of yeast Cu, Zn-superoxide dismutase and its metallochaperone, CCS, localize to the intermembrane space of mitochondria: A physiological role for SOD1 in guarding against mitochondrial oxidative damage. The Journal of Biological Chemistry. 2001;**276**:38084-38089

[66] Mills GC. Hemoglobin catabolism. Glutathione peroxidase, an erythrocyte enzyme which protects hemoglobin from oxidative breakdown. The Journal of Biological Chemistry. 1957;**266**:20752-20760

[67] Forstrom JW, Zakowski JJ, Tappel AL. Identification of the catalytic site of rat liver glutathione peroxidase as selenocysteine. Biochemistry. 1978;**17**(13):2639-2644

[68] Ursini F, Maiorino M, Gregolin C. The selenoenzyme phospholipid hydroperoxide glutathione peroxidase. Biochimica et Biophysica Acta. 1985;**839**:62-70

[69] Chance B, Sies H, Boveris A. Hydroperoxide metabolism in mammalian organs. Physiological Reviews. 1979;**59**:527-605

[70] Chu FF, Doroshov JH, Esworthy RS. Expression characterization, and tissue-distribution of a new cellular selenium-dependent glutathione peroxidase, GSHPx-GI. The Journal of Biological Chemistry. 1993;**268**:2571-2576

[71] Takahashi K, Avissar N, Whittin J, Cohen H. Purification and characterization of human plasma glutathione peroxidase: A selenoglycoprotein distinct from the known cellular enzyme. Archives of Biochemistry and Biophysics. 1987;**256**:677-686

[72] Ghyselinck NB, Dufaure JP. A mouse cDNA sequence for epididymal androgen regulated proteins related to glutathione peroxidase. Nucleic Acids Research. 1990;**18**:7144

[73] Dear TN, Campbell K, Rabbitts TH. Molecular cloning of putative odorant-binding and odorant metabolizing proteins. Biochemistry. 1991;**30**:10376-10382

[74] Kryukov GV, Castellano S, Novoselov SV, Lobanov AV, Zehtab O, Guigó R, et al. Characterization of mammalian selenoproteomes. Science. 2003;**300**:1439-1443

[75] Brigelius-Flohe R, Maiorino M. Glutathione peroxidases. Biochimica et Biophysica Acta. 1980;**2012**:3289-3303

[76] Kim K, Kim IH, Lee KY, Rhee SG, Stadtman ER. The isolation and purification of a specific "protector" protein which inhibits enzyme inactivation by a thiol/Fe(III)/O₂ mixed-function oxidation system. The Journal of Biological Chemistry. 1988;**263**:4704-4711

[77] Peskin AV, Low FM, Paton LN, Maghzal GJ, Hampton MB, Winterbourn CC. The high reactivity of peroxiredoxin 2 with H₂O₂ is not

reflected in its reaction with other oxidants and thiol reagents. The Journal of Biological Chemistry. 2007;**282**(16):11885-11892

[78] Woo HA, Jeong W, Chang TS, Park KJ, Park SJ, Yang JS, et al. Reduction of cysteine sulfinic acid by sulfiredoxin is specific to 2-cys peroxiredoxins. The Journal of Biological Chemistry. 2005;**280**:3125-3128

[79] Hofmann B, Hecht HJ, Flohé L. Peroxiredoxins. Biological Chemistry. 2002;**383**(3-4):347

[80] Yan LJ, Levine RL, Sohal R. Effects of aging and hyperoxia on oxidative damage to cytochrome c in the housefly, *Musca domestica*. Free Radical Biology and Medicine. 2000;**29**(1):90-97

[81] Magwere T, West M, Riyahi K, Murphy MP, Smith RAJ, Partridge L. The effects of exogenous antioxidants on lifespan and oxidative stress resistance in *Drosophila melanogaster*. Mechanisms of Ageing and Development. 2005;**127**:356-370

[82] Rebrin I, Sohal R. Comparison between the effects of aging and hyperoxia on glutathione redox state and protein mixed disulfides in *Drosophila melanogaster*. Mechanisms of Ageing and Development. 2006;**127**:869-874

[83] Leiers B, Kampkötter A, Grevelding CG, Link CD, Johnson TE, Henkle Dührsen K. A stress-responsive glutathione S-transferase confers resistance to oxidative stress in *Caenorhabditis elegans*. Free Radical Biology and Medicine. 2003;**34**(11):1405-1415

[84] Sohal RS, Weindruch R. Oxidative stress, caloric restriction, and aging. Science. 1996;**273**:59-63

[85] Veal EA, Toone WM, Jones N, Morgan BA. Distinct roles

for glutathione S-transferases in the oxidative stress response in *Schizosaccharomyces pombe*. The Journal of Biological Chemistry. 2002;**277**(38):35523-35531

[86] Ketterer B. The role of nonenzymatic reactions of glutathione in xenobiotic mechanism. Drug Metabolism Reviews. 1982;**13**:161-187

[87] Milatovic D, Gupta RC, Aschner M. Anticholinesterase toxicity, oxidative stress. The Scientific World Journal. 2006;**6**:295-310

[88] Rahimi R, Abdollahi M. A review on the mechanisms involved in hyperglycemia induced by organophosphorus pesticides. Pesticide Biochemistry and Physiology. 2007;**88**:115-121

[89] Lukaszewicz-Hussain A, Moniuszko-Jakoniuk J. Procesy glikolityczne w warunkach szkodliwego działania chlorfenwinfosu. Medycyna Pracy. 1997;**48**:580-583

[90] Dabas A, Nagpure NS, Kumar R, Kushwaha B, Kumar P, Lakra WS. Assessment of tissue-specific effect of cadmium on antioxidant defense system and lipid peroxidation in freshwater murrel, *Channa punctatus*. Fish Physiology and Biochemistry. 2012;**38**(2):469-482

[91] Scott JG, Warren WC, Beukeborn LW, Bopp D, Clark AG, Giers SD, et al. Genome of the house fly, *Musca domestica* L., a global vector of diseases with adaptations to a septic environment. Genome Biology. 2014;**15**:466

[92] Liu F, Tang T, Sun L, Priya TAJ. Transcriptomic analysis of the housefly (*Musca domestica*) larva using massively parallel pyrosequencing. Molecular Biology Reports. 2012;**39**:1927-1934