We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Introductory Chapter: Characterization and Improvement of Legume Crops

Mohamed A. El-Esawi

1. Introduction

Legumes are agriculturally grown flowering plants that are found in most of the archaeological record of plants [1]. Various ecosystems, including rain forests, arctic/alpine regions, and deserts have been colonized by legumes [1, 2]. The most two popular flowering plants are Asteraceae and Orchidaceae [1]. The third in terms of popularity is Leguminosae or Fabaceae with 670–750 genera and 18,000–19,000 species, respectively [1]. Legumes are utilized efficiently as (a) food crops for humans and animals; (b) pulps for paper, wood, and timber manufacturing; (c) sources for fuel and oil production; (d) ornamental plants used as living barriers and firebreaks, among others [3]; and (e) cover crops such as cereals and other staple foods [1]. Additionally, they can be utilized for other purposes including production of massive amounts of organic nitrogen. This is because legumes can be intercropped with rhizobia resulting in high yield productivity, soil organic matter improvement, modification of soil osmosis and texture, nutrient reuse, decrease of soil pH and soil pressure, microorganism differentiation, and alleviation of disease problems [1, 4]. Furthermore, legumes can produce amounts of organic nitrogen at a slow rate when rotated with cereals. Such nitrogen produced can be utilized in prospective cropping technologies for improving the production of these crops, recognizing their potential role in promoting better human nutrition and soil health [1, 5].

2. Main legumes

Forage legumes such as alfalfa (*Medicago sativa*), clover (*Trifolium* spp.), bird's-foot trefoil (*Lotus corniculatus*), and vetch (*Vicia* spp.) are utilized as main sources for dairy and meat which are used for protein, fiber, and energy production [1]. Global production of alfalfa was approximately 436 tons in 2006 suggesting that it is the most essential forage crop. The highest amount of alfalfa was produced in the United States, being produced around 15 million tons in 2010 [1, 6]. Grain legumes or pulses are crops harvested massively for the dry seeds. They are found containing high amounts of protein in their seeds. Therefore, they represent a major food source for population consumption. They are considered as the main protein suppliers especially for people from developing countries [1]. Additionally, their high amino acid content is of nutritional value during utilization of cereals and tubers as food sources [1, 7]. The soybean (*Glycine max*), a native plant of Eastern Asia, is an annual

summer legume of great agricultural possibilities due to its fundamental role in the nutrition of many people and livestock besides its industrial possibilities [1, 8].

3. Enhancing legume productivity

Legumes are highly diversified, so they are utilized for several economic and cultural purposes including their role as vegetables to tolerate various ecological conditions, their source for producing large quantities of proteins, their utilization in grazing domains, and their function in increasing worldwide productivity of food and other commodities [1]. Therefore, recent findings have directed towards developing new biological and environment-friendly techniques to enhance the growth efficiency of legumes [1]. Scientists have derived several economic and ecological uses when legumes form symbiotic associations with nitrogen-fixing fungi and bacteria [9]. Biological nitrogen fixation (BNF) within legumes occurs through their association with microorganisms [1]. These microorganisms, which are also needed for the Earth's nitrogen cycle, are utilized for developing agricultural production of plants. Furthermore, they participate in soil colonization and plant growth promotion when utilized in live formulations or biofertilizers applied to seed, root, soil, or the interior of the plant as they can supply large amount of proteins to host cell and enhance soil protection [1]. The need of agroecosystems for nitrogen is assessed through a cost-effective, prospective, and eco-friendly process of biological nitrogen fixation rather than chemical nitrogen fixation. There are several benefits to the process of biological nitrogen fixation. It meets the needs of legumes and intercropped or succeeding crops for nitrogen [1]. This, in turns, avoids or even restricts the application of nitrogen fertilization. Additionally, nitrogen-fixing organisms play an essential role when the amount of nitrogen in the soil is low. They introduce ammonium into the legume biomass to allow faster growing than their plant competitors, but if the protein content is high, nitrogen-fixing microorganisms become alternative to non-fixing species due to high bioenergy cost of nitrogen fixation process [1, 10]. Thus, it can be concluded that nitrogen fixation in legume systems occurs through a variety of physiological and ecological possibilities including the plant's need for nitrogen and the C:N stoichiometry of the ecosystem [1]. It has proven experimentally and theoretically the hypothesis of a feedback control between legume's need for nitrogen and BNF in a specific ecosystem [11].

To enhance the efficiency of the nitrogen fixation process, the most suitable microorganisms for such purpose are selected, and/or genetic engineering of plant species are involved to guarantee high legume crop productivity [1]. Farmers are familiar with the application of commercially available microorganisms (inoculants) that are of great efficiency to nodulate plants and fix nitrogen in the soil [1]. These microorganisms such as rhizobia form associations with legumes in a situation called symbiosis that introduces benefits for both parts [1]. In this scenario, leguminous plants represent the source of energy and photosynthetic products to rhizobia, while rhizobia supplies the legumes with nitrogen in form of ammonium [1, 12]. The symbiosis begins when the roots of leguminous plants are inoculated the rhizobia, which, in turn, form root nodules where BNF occurs with the help of nitrogenase enzyme [1, 13]. In conclusions, several techniques have been developed genetically and biochemically to enhance plant development and crop productivity, suggesting their marvelous importance in improving legumes and other crops [1, 14–37].

IntechOpen

IntechOpen

Author details

Mohamed A. El-Esawi

Botany Department, Faculty of Science, Tanta University, Tanta, Egypt

*Address all correspondence to: mohamed.elesawi@science.tanta.edu.eg

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Morel MA, Braña V, Castro-Sowinski S. Legume Crops, Importance and Use of Bacterial Inoculation to Increase Production, Crop Plant, Aakash Goyal. IntechOpen; 2012. DOI: 10.5772/37413. Available from: <https://www.intechopen.com/books/crop-plant/legume-crops-importance-and-use-of-bacterial-inoculation-to-increase-production>
- [2] Schrire BD, Lewis GP, Lavin M. Biogeography of the Leguminosae. In: Lewis G, Schrire G, Mackinder B, Lock M, editors. Legumes of the World. Kew, UK: Royal Botanic Gardens. ISBN: 8773043044; 2005. pp. 21-54
- [3] Lewis G, Schrire B, MacKinder B, Lock M. Legumes of the World. UK: Royal Botanical Gardens, Kew Publishing; 2005. ISBN: 1900347806
- [4] USDA. Soil Quality-Agronomy Technical Note. 1998. Available from: http://soils.usda.gov/sqi/management/files/sq_atn_6.pdf [Accessed: 30 August 2011]
- [5] Popelka C, Terryn N, Higgins T. Gene technology for grain legumes: Can it contribute to the food challenge in developing countries? Plant Science. 2004;167:195-206. ISSN: 01689452
- [6] FAO STAT. Food and Agriculture Organization Statistical Database. 2011. Available from: <http://faostat.fao.org> [Accessed: 30 August]
- [7] Graham PH, Vance CP. Legumes: Importance and constraints to greater use. Plant Physiology. 2003;131:872-877. ISSN: 0032-0889
- [8] Ali N. Soybean processing and utilization. In: Singh G, editor. The Soybean: Botany, Production and Uses. UK: CAB International. ISBN: 9781845936440; 2010. pp. 345-362
- [9] Zahran HH. Enhancement of Rhizobia-legumes symbioses and nitrogen fixation for crops productivity improvement. In: Khan MS, Zaidi A, Musarrat J, editors. Microbial Strategies for Crop Improvement. Netherlands: Springer; 2009. pp. 227-254. ISBN: 9783642019784
- [10] Houlton BZ, Wang YP, Vitousek PM, Field CB. A unifying framework for dinitrogen fixation in the terrestrial biosphere. Nature. 2008;454:27-334. ISSN: 0028-0836
- [11] Soussana J-F, Tallec T. Can we understand and predict the regulation of biological N₂ fixation in grassland ecosystems? Nutrient Cycling in Agroecosystems. 2010;88:197-213. ISSN: 1385-1314
- [12] Howard JB, Rees DC. Structural basis of biological nitrogen fixation. Chemical Reviews. 1996;96:2965-2982. ISSN: 0009-2665
- [13] Masson-Boivin C, Giraud E, Perret X, Batut J. Establishing nitrogen-fixing symbiosis with legumes: How many *Rhizobium* recipes? Trends in Microbiology. 2009;17:458-466. ISSN: 0966-842X
- [14] Elansary HO, Yessoufou K, Abdel-Hamid AME, El-Esawi MA, Ali HM, Elshikh MS. Seaweed extracts enhance Salam Turfgrass performance during prolonged irrigation intervals and saline shock. Frontiers in Plant Science. 2017;8:830
- [15] El-Esawi MA. Micropropagation technology and its applications for crop improvement. In: Anis M, Ahmad N, editors. Plant Tissue Culture: Propagation, Conservation and Crop Improvement. Singapore: Springer; 2016. pp. 523-545
- [16] El-Esawi MA. Nonzygotic embryogenesis for plant development.

In: Anis M, Ahmad N, editors.
Plant Tissue Culture: Propagation,
Conservation and Crop Improvement.
Singapore: Springer; 2016. pp. 583-598

[17] El-Esawi MA. Somatic hybridization and microspore culture in brassica improvement. In: Anis M, Ahmad N, editors. Plant Tissue Culture: Propagation, Conservation and Crop Improvement. Singapore: Springer; 2016. pp. 599-609

[18] El-Esawi MA. Genetic diversity and evolution of *Brassica* genetic resources: From morphology to novel genomic technologies—A review. Plant Genetic Resources. 2017;15:388-399

[19] El-Esawi MA. SSR analysis of genetic diversity and structure of the germplasm of faba bean (*Vicia faba* L.). Comptes Rendus Biologies. 2017;340:474-480

[20] El-Esawi MA, Alayafi AA. Overexpression of rice *Rab7* gene improves drought and heat tolerance and increases grain yield in rice (*Oryza sativa* L.). Genes. 2019;10:56

[21] El-Esawi MA, Sammour R. Karyological and phylogenetic studies in the genus *Lactuca* L. (Asteraceae). Cytologia. 2014;79:269-275

[22] El-Esawi MA, Al-Ghamdi AA, Ali HM, Alayafi AA, Witczak J, Ahmad M. Analysis of genetic variation and enhancement of salt tolerance in French pea (*Pisum sativum* L.). International Journal of Molecular Sciences. 2018;19:2433

[23] El-Esawi MA, Alaraidh IA, Alsahli AA, Ali HM, Alayafi AA, Witczak J, et al. Genetic variation and alleviation of salinity stress in barley (*Hordeum vulgare* L.). Molecules. 2018;23:2488

[24] El-Esawi MA, Alaraidh IA, Alsahli AA, Alamri SA, Ali HM,

Alayafi AA. *Bacillus firmus* (SW5) augments salt tolerance in soybean (*Glycine max* L.) by modulating root system architecture, antioxidant defense systems and stress-responsive genes expression. Plant Physiology and Biochemistry. 2018;132:375-384

[25] El-Esawi MA, Alaraidh IA, Alsahli AA, Alzahrani SM, Ali HM, Alayafi AA, et al. *Serratia liquefaciens* KM4 improves salt stress tolerance in maize by regulating redox potential, ion homeostasis, leaf gas exchange and stress-related gene expression. International Journal of Molecular Sciences. 2018;19:3310

[26] El-Esawi MA, Elansary HO, El-Shanhorey NA, Abdel-Hamid AME, Ali HM, Elshikh MS. Salicylic acid-regulated antioxidant mechanisms and gene expression enhance rosemary performance under saline conditions. Frontiers in Physiology. 2017;8:716

[27] El-Esawi M, Arthaut L, Jourdan N, d'Harlingue A, Martino C, Ahmad M. Blue-light induced biosynthesis of ROS contributes to the signaling mechanism of *Arabidopsis* cryptochrome. Scientific Reports. 2017;7:13875

[28] El-Esawi MA, Elkelish A, Elansary HO, Ali HM, Elshikh M, Witczak I, et al. Genetic transformation and hairy root induction enhance the antioxidant potential of *Lactuca serriola* L. Oxidative Medicine and Cellular Longevity. 2017;2017. Article ID: 5604746, 8 pages

[29] El-Esawi MA, Germaine K, Bourke P, Malone R. Genetic diversity and population structure of *Brassica oleracea* germplasm in Ireland using SSR markers. Comptes Rendus Biologies. 2016;339:133-140

[30] El-Esawi MA, Germaine K, Bourke P, Malone R. AFLP analysis of genetic diversity and phylogenetic

relationships of *Brassica oleracea* in Ireland. *Comptes Rendus Biologies*. 2016;**339**:163-170

transcription factor enhances drought stress tolerance in cotton (*Gossypium barbadense* L.). *Genes*. 2019;**10**:142

[31] El-Esawi M, Glascoe A, Engle D, Ritz T, Link J, Ahmad M. Cellular metabolites modulate *in vivo* signaling of *Arabidopsis* cryptochrome-1. *Plant Signaling & Behavior*. 2015;**10**:e1063758

[32] El-Esawi MA, Mustafa A, Badr S, Sammour R. Isozyme analysis of genetic variability and population structure of *Lactuca* L. germplasm. *Biochemical Systematics and Ecology*. 2017;**70**:73-79

[33] El-Esawi MA, Al-Ghamdi AA, Ali HM, Alayafi AA. *Azospirillum lipoferum* FK1 confers improved salt tolerance in chickpea (*Cicer arietinum* L.) by modulating osmolytes, antioxidant machinery and stress-related genes expression. *Environmental and Experimental Botany*. 2019;**159**:55-65

[34] Jourdan N, Martino C, El-Esawi M, Witczak J, Bouchet P-E, d'Harlingue A, et al. Blue light dependent ROS formation by *Arabidopsis* cryptochrome-2 may contribute towards its signaling role. *Plant Signaling & Behavior*. 2015;**10**:e1042647

[35] Vwioko E, Adinkwu O, El-Esawi MA. Comparative physiological, biochemical and genetic responses to prolonged waterlogging stress in okra and maize given exogenous ethylene priming. *Frontiers in Physiology*. 2017;**8**:632

[36] El-Esawi MA, Al-Ghamdi AA, Ali HM, Ahmad M. Overexpression of *AtWRKY30* transcription factor enhances heat and drought stress tolerance in wheat (*Triticum aestivum* L.). *Genes*. 2019;**10**(2):163

[37] El-Esawi MA, Alayafi AA. Overexpression of *StDREB2*