We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Introductory Chapter: Definition of Cognitive Behavioral Therapy and Its Principal Applications

Sandro Misciagna

1. Story of modern cognitive behavioral therapy

Cognitive behavioral therapy (CBT) is a modern form of short-term psychotherapy based on the idea that the way an individual thinks and feels affects the way he behaves.

The core premise of this treatment approach was pioneered by Albert Ellis who in 1957 introduced the term “rational emotive therapy” (RET) to emphasize its focus on emotional outcomes. Successively, Aaron Beck in 1976 created “cognitive therapy” (CT), which served as the bases for the development of CBT.

According to Beck’s formulation, maladaptive cognitions, which consist in general beliefs or schemas about the self, the world and the future, contribute to the maintenance of emotional distress and behavioral problems. According to this model, specific therapeutic strategies that change maladaptive cognitions lead to change emotional distress and problematic behaviors.

In 1995, Ellis created the term “rational emotive behaviour therapy” (REBT) because behavioral factors constitute a fundamental component of this treatment approach [1].

More recently, practitioners and scholars started to call it rational emotive and “cognitive behaviour therapy” to emphasize its role in CBT paradigm.

REBT protocols were applied in domains of clinical psychology, rational emotive education, organizational setting or counseling [2].

Since the late 1980s and 1990s, REBT was investigated in a series of randomized control trials that demonstrated its efficacy in many psychological conditions such as social phobia [3], obsessive–compulsive disorders [4], depression [5], psychotic symptoms [6] and behavioral disorders.

In 2013, on the bases of meta-analysis studies, REBT was included in the category of CBT [7] since its protocols are similar in structure with CBT protocols, while the main difference is that REBT specifically focuses on evaluative beliefs and not descriptive or inferential ones [8].

Since these early models, CBT have developed appropriate protocols to treat subjects of almost every age such as children, adolescent, adults or elderly and for individual, families and couples.

2. Principal applications of CBT

Cognitive behavioral therapy is an empirical and evidence-based psychotherapy that integrates cognitive science and behavioral theories, combined with clinical

psychology, to conclude that the way people perceive a situation determines their reaction more than the actual reality of the situation does [9].

CBT provides useful tools that can be used to induce or facilitate belief revision such as cognitive restructuring or exposure/response prevention; these protocols have been applied both in groups and in individuals, even if the individual format is used more frequently than the group format.

This treatment plan uses patients' collaboration as a motivating factor, in order to generate changes in their behaviour, beliefs and habits that can be self-reinforced. In fact, CBT gets patients actively involved in their treatment so that they understand that the way to improve their lives is to adjust their thinking and their approach to everyday situations.

The overall goal of this treatment is symptom reduction, improvement in functioning and remission of the disorder. The initial sessions of CBT illustrate the close relationship between cognition and emotions. Each typical therapy session begins establishing an agenda of current problems, followed by cognitive restructuring of maladaptive cognitions. At the end of the session, the therapist assigns homework to help the patient to apply specific skills in his real life. Every step of CBT is reasoned and transparent. If the patient suffers from psychomotor retardation, behavioral strategies are implemented with cognitive interventions. The therapist can also use a series of questions to help the patients evaluate the utility and validity of their cognitions [10].

CBT consists in different protocols that are effective in the treatment of a vast variety of mental disorders, such as generalized anxiety disorders, panic disorders (in particular with agoraphobia or social phobia) and obsessive-compulsive disorders [11].

The session content varies for each specific disorder based on the empirically proven cognitive and behavioral model of each disorder. For example, cognitive themes concern about consequences of a panic attack (in phobia disorders), concern about social embarrassment (in social anxiety disorders), concern about dangers of worrying (in generalized anxiety disorders) and concern about consequences of intrusive thoughts (in obsessive-compulsive disorders).

CBT can be used to treat mood and emotional disorders such as depression or dysthymia [12]. With depressive disorders, the general therapeutic process of CBT is to split up into different steps, with an insistence on distinct therapeutic mechanisms, establishing a therapeutic relationship and managing maladaptive behaviors and cognitions.

CBT has been extensively tested for a wide range of neurotic and stress-related disorders such as general stress, post-traumatic stress disorders and somatoform disorders (such as hypochondriasis and body dysmorphic disorder) or medical problems with psychological components.

Several studies have demonstrated CBT utility in problematic gambling, substance use disorder (as nicotine, cannabis, opioid or alcohol dependence), eating disorders (as bulimia nervosa, binge eating disorders) [13] and sleep dysfunction (in particular insomnia) [14] or to approach fatigue, chronic pain conditions [15, 16] and inflammation pathologies [17], especially if associated with distress [18].

CBT is probably effective also for psychotic disorders associated with positive symptoms (i.e. delusions and/or hallucinations) in schizophrenia, personality disorders (including antisocial personality disorder) [19], anger expression (anger, verbal and physical aggression, driving anger, anger suppression and anger difficulties) and bipolar disorders. CBT is particularly promising for schizophrenia in patients who suffer from acute episode of psychosis rather than a more chronic condition [20].

Neurofunctional studies have demonstrated that CBT induces brain activation and functional changes in the amygdala, insula and anterior corticolimbic brain circuits that control cognitive, motivational and emotional aspects of physiology and behaviour [21].

CBT should be used by a health professional with experience and training in cognitive and behavioral therapies, especially when used for the treatment of anxiety and mood disorders.

This book, written by authors that are expertice in CBT, is useful both for clinicians and psychotherapists who wants to understand modern cognitive psychology and develop specific personalized treatment plans of cognitive behavioral therapy.

Author details

Sandro Misciagna

Neurology Department and Stroke Unit, Belcolle Hospital, Viterbo, Italy

*Address all correspondence to: sandromisciagna@yahoo.it

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Ellis A. Changing rational-emotive therapy (RET) to rational emotive behavior therapy (REBT). *Journal of Rational-Emotive & Cognitive-Behavior Therapy*. 1995;13(2):85-89
- [2] David D. *Rational Emotive Behavior Therapy*. New York: Oxford University Press; 2014
- [3] Mersch PPA, Emmelkamp PM, Bögels SM, Van der Sleen J. Social phobia: Individual response patterns and the effects of behavioral and cognitive interventions. *Behaviour Research and Therapy*. 1989;27(4):421-434
- [4] Emmelkamp PM, Beens H. Cognitive therapy with obsessive-compulsive disorder: A comparative evaluation. *Behaviour Research and Therapy*. 1991;29(3):293-300
- [5] David D, Szentagotai A, Lupu V, Cosman D. Rational emotive behavior therapy, cognitive therapy, and medication in the treatment of major depressive disorder: A randomized clinical trial, posttreatment outcomes, and six-month followup. *Journal of Clinical Psychology*. 2008;64(6):728-746
- [6] Meaden A, Keen N, Aston R, Barton K, Bucci S. *Cognitive Therapy for Command Hallucinations: An Advanced Practical Companion*. New York: Routledge; 2013
- [7] Cuijpers P, Berking M, Andersson G, Quigley L, Kleiboer A, Dobson KS. A meta-analysis of cognitive behavioural therapy for adult depression, alone and in comparison with other treatments. *Canadian Journal of Psychiatry*. 2013;58(7):376-385
- [8] David D, Lynn SJ, Ellis A. *Rational and Irrational Beliefs: Research, Theory, and Clinical Practice*. Oxford: Oxford University Press; 2010
- [9] Butler AC, Chapman JE, Forman EM, Beck AT. The empirical status of cognitive-behavioral therapy: A review of meta-analyses. *Clinical Psychology Review*. 2006;26:17-31
- [10] Tarrier N, editor. *Case Formulation in Cognitive Behaviour Therapy: The Treatment of Challenging and Complex Cases*. New York: Routledge/Taylor & Francis Group; 2006
- [11] Clark DM, Ehlers A, Hackmann A, et al. Cognitive therapy versus exposure and applied relaxation in social phobia: A randomized controlled trial. *Journal of Consulting and Clinical Psychology*. 2006;74:568-578
- [12] Stubbings DR, Rees CS, Roberts LD, Kane RT. Comparing in-person to videoconference-based cognitive behavioral therapy for mood and anxiety disorders: Randomized controlled trial. *Journal of Medical Internet Research*. 2013;15(11):e258
- [13] Valenzuela F, Lock J, Le Grange D, Bohon C. Comorbid depressive symptoms and self-esteem improve after either cognitive-behavioural therapy or family-based treatment for adolescent bulimia nervosa. *European Eating Disorders Review*. 2018;26(3):253-258. DOI: 10.1002/erv.2582. Epub: February 15, 2018
- [14] Cheong MJ, Lee GE, Kang HW, Kim S, Kim HK, Jo HI, et al. Clinical effects of mindfulness meditation and cognitive behavioral therapy standardized for insomnia: A protocol for a systematic review and meta-analysis. *Medicine (Baltimore)*. 2018;97(51):1-7
- [15] Bach E, Beissner K, Murtaugh C, Trachtenberg M, Reid MC. Implementing a cognitive-behavioral pain self-management program in home health care. Part 2: Feasibility

and acceptability cohort study. *Journal of Geriatric Physical Therapy* (2001). 2013;**36**(3):130-137

[16] Beissner K, Bach E, Murtaugh C, Parker SJ, Trachtenberg M, Reid MC. Implementing a cognitive-behavioral pain self-management program in home health care. Part 1: Program adaptation. *Journal of Geriatric Physical Therapy* (2001). 2013;**36**(3):123-129

[17] Lopresti AL. Cognitive behaviour therapy and inflammation: A systematic review of its relationship and the potential implications for the treatment of depression. *The Australian and New Zealand Journal of Psychiatry*. 2017;**51**(6):565-582

[18] Hofmann S, Reinecke M, editors. *Cognitive-Behavioral Therapy with Adults: A Guide to Empirically Informed Assessment and Intervention*. Cambridge, UK: Cambridge University Press; 2010

[19] Sanatinia R, Wang D, Tyrer P, Tyrer H, Crawford M, Cooper S, et al. Impact of personality status on the outcomes and cost of cognitive-behavioural therapy for health anxiety. *The British Journal of Psychiatry*. 2016;**209**(3):244-250. DOI: 10.1192/bjp.bp.115.173526. Epub: July 21, 2016

[20] Morrison AP, Law H, Carter L, Sellers R, Emsley R, Pyle M, et al. Antipsychotic drugs versus cognitive behavioural therapy versus a combination of both in people with psychosis: A randomised controlled pilot and feasibility study. *Lancet Psychiatry*. 2018;**5**:411-423

[21] Lueken U, Straube B, Konrad C, Wittchen HU, Ströhle A, Wittmann A, et al. Neural substrates of treatment response to cognitive-behavioral therapy in panic disorder with agoraphobia. *The American Journal of Psychiatry*. 2013;**170**(11):1345-1355