We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Numerical Solutions to Some Families of Fractional Order Differential Equations by Laguerre Polynomials

Adnan Khan, Kamal Shah and Danfeng Luo

Abstract

This article is devoted to compute numerical solutions of some classes and families of fractional order differential equations (FODEs). For the required numerical analysis, we utilize Laguerre polynomials and establish some operational matrices regarding to fractional order derivatives and integrals without discretizing the data. Further corresponding to boundary value problems (BVPs), we establish a new operational matrix which is used to compute numerical solutions of boundary value problems (BVPs) of FODEs. Based on these operational matrices (OMs), we convert the proposed (FODEs) or their system to corresponding algebraic equation of Sylvester type or system of Sylvester type. The resulting algebraic equations are solved by MATLAB® using Gauss elimination method for the unknown coefficient matrix. To demonstrate the suggested scheme for numerical solution, many suitable examples are provided.

Keywords: FODEs, numerical solution, Laguerre polynomials, operational matrices

1. Introduction

The theory of integrals as well as derivatives of arbitrary order is known by the special name “fractional calculus.” It has an old history just like classical calculus. The chronicle of fractional calculus and encyclopedic book can be studied in [1, 2]. Researchers have now necessitated the use of fractional calculus due to its diverse applications in different fields, specially in electrical networks, signal and image processing and optics, etc. For conspicuous work on FODEs in the fields of dynamical systems, electrochemistry, advanced techniques of microorganisms culturing, weather forecasting, as well as statistics, we refer to peruse [3, 4]. Fractional derivatives show valid results in most cases where ordinary derivatives do not. Also annotating that fractional order derivatives as well as fractional integrals are global operators, while ordinary derivatives are local operators. Fractional order derivative provides greater degree of freedom. Therefore from different aspects, the aforesaid areas were investigated. For instance, many researchers have provide understanding to existence and uniqueness results about FODEs, for few results, we refer [5–7], and many others have actualized the instinctive framework of fractional differential equations in various problems [8–19] with many references included in them.

Often it is very difficult to obtain the exact solution due to global nature of fractional derivatives in differential equations. Contrarily approximate solutions are obtained by numerical methods assorted in [20–22]. Various new numerical methods have been developed, among them is one famous method called “spectral method” which is used to solve problems in various realms [23]. In this method operational matrices are obtained by using orthogonal polynomials [24]. Many authors have successfully developed operational matrices by using Legendre, Jacobi, and various other polynomials [25, 26]. For delay differential and various other related equations, Laguerre spectral methods have been used [27–32]. Bernstein polynomials and various classes of other polynomials were also used to obtain operational matrices corresponding to fractional integrals and derivatives [33–40]. Apart from them, operational matrices were also developed with the collocation method (see Refs. [41–43]). Since spectral methods are powerful tools to compute numerical solutions of both ODEs and FODEs. Therefore, we bring out numerical analysis via using Laguerre polynomials of some families and coupled systems of FODEs under initial as well as boundary conditions. In this regard we investigate the numerical solutions to the given families under initial conditions

$$\begin{cases} {}^c_0D_t^\gamma z(t) \pm z(t) = 0, & 0 < \gamma \leq 1, \\ z(0) = z_0, & z_0 \in R, \end{cases} \quad (1)$$

and subject to boundary conditions

$$\begin{cases} {}^c_0D_t^\gamma z(t) \pm z(t) = 0, & 1 < \gamma \leq 2, \\ z(0) = z_0, z(1) = z_1, & z_0, z_1 \in R. \end{cases} \quad (2)$$

By similar numerical techniques, we also investigate the numerical solutions to the following systems with fractional order derivatives under initial and boundary conditions as

$$\begin{cases} {}^c_0D_t^\gamma z(t) + az(t) + by(t) = f(t), \\ {}^c_0D_t^\gamma y(t) + cy(t) + dz(t) = g(t), \\ z(0) = z_0, y(0) = y_0 \end{cases} \quad (3)$$

for $0 < \gamma \leq 1$ and

$$\begin{cases} {}^c_0D_t^\gamma z(t) + az(t) + by(t) = f(t), \\ {}^c_0D_t^\gamma y(t) + cy(t) + dz(t) = g(t), \\ z(0) = z_0, y(0) = y_0, & z(1) = z_1, y(1) = y_1, \end{cases} \quad (4)$$

for $1 < \gamma \leq 2$ where $f, g : [0, 1] \times R^2 \rightarrow R$ and $z_0, y_0, z_1, y_1 \in R$. We first obtain OM for fractional derivatives and integrals by using Laguerre polynomials. Also corresponding to boundary conditions, we construct an operational matrix which is needed in numerical analysis of BVPs. With the help of the OM we convert the considered problem of FODEs under initial/boundary conditions to Sylvester-type algebraic equations. Solving the mentioned matrix equations by using MATLAB®, we compute the numerical solutions of the considered problems.

2. Preliminaries

Here we recall some basic definition results that are needed in this work onward, keeping in mind that throughout the paper we use fractional derivative in Caputo sense.

Definition 1. The fractional integral of order $\gamma > 0$ of a function $z : (0, \infty) \rightarrow R$ is defined by

$${}_0I_t^\gamma z(t) = \frac{1}{\Gamma(\gamma)} \int_0^t \frac{z(s)}{(t-s)^{1-\gamma}} ds,$$

provided the integral converges at the right sides. Further a simple and important property of ${}_0I_t^\gamma$ is given by

$${}_0I_t^\gamma t^\delta = \frac{\Gamma(\delta+1)}{\Gamma(\delta+\gamma+1)} t^{\gamma+\delta}.$$

Definition 2. Caputo fractional derivative is defined as

$${}_0^cD_t^\gamma f(t) = \frac{1}{\Gamma(n-\gamma)} \int_0^t (t-s)^{n-\gamma-1} f^{(n)}(s) ds,$$

where n is a positive integer with the property that $n-1 < \gamma \leq n$. For example, if $0 < \gamma \leq 1$, then Caputo fractional derivative becomes

$${}_0^cD_t^\gamma f(t) = \frac{1}{\Gamma(1-\gamma)} \int_0^t (t-s)^{-\gamma-1} f'(s) ds.$$

Theorem 1. The FODE given by

$${}_0^cD_t^\gamma f(t) = 0$$

has a unique solution, such that

$$f(t) = d_0 + d_1 t + d_2 t^2 + \dots + d_{n-1} t^{n-1}, \quad n = [\gamma] + 1.$$

Lemma 1. Therefore in view of this result, if $h \in L^n[0, T]$, then the unique solution of nonhomogenous FODE

$${}_0^cD_t^\gamma f(t) = h(t), \quad n-1 < \gamma \leq n$$

is written as

$$f(t) = d_0 + d_1 t + d_2 t^2 + \dots + d_{n-1} t^{n-1} + {}_0I_t^\gamma h(t),$$

where d_i for $i = 0, 1, 2, 3 \dots n-1$ are real constants.

The above lemma is also stated as

$$f(t) = {}_0I_t^\gamma h(t) + \sum_{i=0}^{n-1} \frac{f^{(i)}(0)}{i!} t^i.$$

Definition 3. The famous Laguerre polynomials are represented by $L_i^\gamma(t)$ and defined as

$$L_i^\gamma(t) = \sum_{k=0}^i \frac{(-1)^k \Gamma(i+\gamma+1)}{\Gamma(k+1+\gamma) \Gamma(i-k+1) \Gamma(k+1)} t^k.$$

They are orthogonal on $[0, \infty]$. If $L_i^\gamma(t)$ and $L_j^\gamma(t)$ are Laguerre polynomials, then the orthogonality condition is given as

$$\int_0^\infty L_i^\gamma(t) L_j^\gamma(t) W^\gamma(t) dt = \delta_{ij} U_k,$$

where

$$W^\gamma(t) = t^\gamma e^{-t},$$

is the weight function and

$$U_k = \begin{cases} \frac{\Gamma(1 + \gamma + k)}{\Gamma(1 + k)}, & i = j \\ 0 & i \neq j. \end{cases}$$

Now let $Z(t)$ be any function, defined on the interval $[0, \infty]$. We express the function in terms of Laguerre polynomials as

$$\begin{aligned} Z(t) &= \sum_{i=0}^n c_i L_i^\gamma(t). \\ &= c_0 L_0^\gamma(t) + c_1 L_1^\gamma(t) + \dots + c_N L_N^\gamma(t) \\ &= [c_0 \ c_1 \ \dots \ c_N] \begin{bmatrix} L_0^\gamma(t) \\ \vdots \\ L_n^\gamma(t) \end{bmatrix}. \end{aligned} \quad (5)$$

We set the above two vectors into their inner product and represent the column matrix by $\Psi(t)$, so that

$$Z(t) = c^t \Psi(t).$$

Again as

$$\begin{aligned} Z(t) &= \sum_{i=0}^n c_i L_i^\gamma(t), \\ \int_0^L Z(t) W^\gamma(t) L_j^\gamma(t) dt &= \int_0^L \sum_{i=0}^n c_i L_i^\gamma(t) L_j^\gamma(t) W^\gamma(t) dt, \end{aligned}$$

which is written as

$$\sum_{i=0}^n c_i \int_0^L L_i^\gamma(t) L_j^\gamma(t) W^\gamma(t) dt.$$

We call h_i to the general term of integration

$$\int_0^L Z(t) W^\gamma(t) L_j^\gamma(t) dt = \sum_{i=0}^n c_i h_i.$$

Hence the coefficient c_i is

$$c_i = \frac{1}{h_i} \int_0^L Z(t) W^T(t) L_j^T(t) dt.$$

In vector form we can write Eq. (5) as

$$Z(t) = c_M^T \Psi_M(t).$$

where $M = m + 1$, c_M is the M terms coefficient vector and $\Psi_M(t)$ is the M terms function vector.

2.1 Representation of Laguerre polynomial with Caputo fractional order derivative

If the Caputo fractional order derivative is applied to Laguerre polynomial, by considering whole function constant except t^k . We use the definition of Caputo fractional order derivative for t^k to obtain (6) as

$${}_0^C D_t^\gamma L_i^\gamma(t) = \sum_{k=0}^i (t^{k-\gamma}) \frac{(-1)^k \Gamma(i + \gamma + 1)}{\Gamma(k + 1 + \gamma) \Gamma(i - k + 1) \Gamma(1 + k - \gamma)}. \quad (6)$$

2.2 Error analysis

The proof of the following results can be found with details in [20].

Lemma 2. Let $L_i^\beta(t)$ be given; then

$${}_0^C D_t^\gamma L_i^\beta(t) = 0, \quad i = 0, 1, 2, \dots, [\beta] - 1, \gamma > 0.$$

Theorem 2. For error analysis, we state the theorem such that, a be any integer and $0 \leq s \leq a$, and then

$$\|P_{M,az} - z(t)\|_{A_\alpha^s, \Lambda} \leq c M^{\frac{s-a}{2}} |z(t)|_{A_\alpha^a, \Lambda}, \quad \forall z(t) \in A_\alpha^a(\Lambda),$$

where $A_\alpha^a = \{z/z \text{ is measurable on } \Lambda \text{ and } \|z\|_{A_\alpha^a, (\Lambda)} < \infty\}$ and

$$|z|_{A_\alpha^a, (\Lambda)} = \|\partial_p^a z\|_{w\alpha+a, \Lambda},$$

$$\|z\|_{A_\alpha^a, (\Lambda)} = \left(\sum_{k=0}^a |z|_{A_\alpha^k, (\Lambda)}^2 \right)^{\frac{1}{2}}.$$

Now let $\Lambda = \mathcal{Q}/0 < \mathcal{Q} < \infty$ with $\chi(\mathcal{Q})$ be a weight function. Then

$$L_\chi^2(\Lambda) = \{\kappa / \kappa \text{ is measurable on } \Lambda \text{ and } \|u\|_{L_\chi^2, \Lambda} < \infty\}.$$

with the following inner product and norm

$$(u, v)_{\chi, \Lambda} = \int_\Lambda u(\mathcal{Q}) v(\mathcal{Q}) d\mathcal{Q}, \quad \|v\|_{\chi, \Lambda} = \sqrt{\langle u, v \rangle_{\chi, \Lambda}}.$$

3. Operational matrices corresponding to fractional derivatives and integrals

Here in this section, we provide the required OM's via Laguerre polynomials of fractional derivatives and integrals.

Lemma 3. Let $\Psi_M(t)$ be a function vector; the fractional integral of order γ for the function $\Psi_M(t)$ can be generalized as

$${}_0I_t^\gamma \Psi_M(t) \approx G_{N \times N}^\gamma \Psi_M(t),$$

where $G_{N \times N}^\gamma$ is the OM of integration of fractional order γ and given by

$$\begin{bmatrix} \Upsilon_{0,0,k,r}^\gamma & \Upsilon_{0,1,k,r}^\gamma & \cdots & \Upsilon_{0,j,k,r}^\gamma & \cdots & \Upsilon_{0,m,k,r}^\gamma \\ \Upsilon_{1,0,k,r}^\gamma & \Upsilon_{1,1,k,r}^\gamma & \cdots & \Upsilon_{1,j,k,r}^\gamma & \cdots & \Upsilon_{1,m,k,r}^\gamma \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \Upsilon_{i,0,k,r}^\gamma & \Upsilon_{i,1,k,r}^\gamma & \cdots & \Upsilon_{i,j,k,r}^\gamma & \vdots & \Upsilon_{i,m,k,r}^\gamma \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \Upsilon_{m,0,k,r}^\gamma & \Upsilon_{m,1,k,r}^\gamma & \cdots & \Upsilon_{m,j,k,r}^\gamma & \cdots & \Upsilon_{m,m,k,r}^\gamma \end{bmatrix},$$

where

$$\Upsilon_{ij,k,r}^\gamma = \sum_{k=0}^i \sum_{r=0}^i \frac{(-1)^{k+r} \Gamma(j+1) \Gamma(i+\gamma+1) \Gamma(k+\gamma+\alpha+r+1)}{\Gamma(j-r+1) \Gamma(i-k+1) \Gamma(r+1) \Gamma(k+\gamma+1) \Gamma(k+\alpha+1) \Gamma(\gamma+r+1)}.$$

Proof. We apply the fractional order integral of order γ to the Laguerre polynomials

$${}_0^c I_t^\gamma L_i^\gamma(t) = \sum_{k=0}^i \frac{\Gamma(i+\gamma+1)}{\Gamma(i-k+1) \Gamma(k+\gamma+1) \Gamma(k+1)} {}_0^c I_t^\gamma t^k. \quad (7)$$

Since from (7), we have

$${}_0^c I_t^\gamma t^k = \frac{\Gamma(k+1)}{\Gamma(1+k+\alpha)} t^{k+\gamma}.$$

Therefore Eq. (7) implies that

$${}_0^c I_t^\gamma L_i^\gamma(t) = \sum_{k=0}^i t^{k+\gamma} \frac{\Gamma(i+\gamma+1)}{\Gamma(i-k+1) \Gamma(k+\gamma+1) \Gamma(k+1)} \frac{\Gamma(k+1)}{\Gamma(1+k+\alpha)},$$

which is equal to

$${}_0^c I_t^\gamma L_i^\gamma(t) = \sum_{k=0}^i (-1)^k \frac{\Gamma(i+\gamma+1)}{\Gamma(i-k+1) \Gamma(k+\gamma+1) \Gamma(1+k-\gamma)} t^{k+\gamma}. \quad (8)$$

We approximate $t^{k+\gamma}$ in (8) with Laguerre polynomials, i.e.

$$t^{k+\gamma} \approx \sum_{j=0}^n H_j L_j^\gamma(t).$$

By using the relation of orthogonality, we can find coefficients

$$H_j = \sum_{r=0}^j (-1)^k \frac{\Gamma(j+1) \Gamma(k+\alpha+r+\gamma+1)}{\Gamma(1+j-r) \Gamma(1+r) \Gamma(1+r+\gamma)}.$$

So Eq. (8) implies

$$\begin{aligned}
 {}^c_0I_t^\gamma L_i^\gamma(t) &= \sum_{k=0}^i (-1)^k \frac{\Gamma(i+\gamma+1)}{\Gamma(i-k+1)\Gamma(k+\gamma+1)\Gamma(1+k-\gamma)} \\
 &\quad \times \sum_{r=0}^j (-1)^r \frac{\Gamma(j+1)\Gamma(k+\alpha+r+\gamma+1)}{\Gamma(j-r+1)\Gamma(r+1)\Gamma(r+\gamma+1)}. \\
 {}^c_0I_t^\gamma L_i^\gamma(t) &= \sum_{k=0}^i \sum_{r=0}^j (-1)^{k+r} \frac{\Gamma(j+1)\Gamma(i+\gamma+1)\Gamma(k+\alpha+r+\gamma+1)}{\Gamma(1-k+i)\Gamma(j-\gamma+1)\Gamma(\gamma+1)\Gamma(k+\gamma+1)\Gamma(k+\alpha+1)\Gamma(\gamma+r+1)}.
 \end{aligned}$$

which is the desired result.

Lemma 4. Let $\Psi_M(t)$ be a function vector; then the fractional derivative of order γ for $\Psi_M(t)$ is generalized as

$${}^cD_t^\gamma \Psi_M(t) \approx \mathbf{W}_{M \times M}^\gamma \Psi_M(t),$$

where $\mathbf{W}_{M \times M}^\gamma$ is the OM of derivative of order γ , defined as in (9)

$$\mathbf{W}_{M \times M}^\gamma = \begin{bmatrix} 0 & 0 & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \Theta_{[\gamma],0,k,\alpha}^\gamma & \Theta_{[\gamma],1,k,\alpha}^\gamma & \dots \Theta_{[\gamma],j,k,\alpha}^\gamma & \dots & \dots & \Theta_{[\gamma],n,k,\alpha}^\gamma \\ \Theta_{i,0,k,\alpha}^\gamma & \Theta_{i,1,k,\alpha}^\gamma & \Theta_{i,j,k,\alpha}^\gamma & \dots & \dots & \Theta_{i,n,k,\alpha}^\gamma \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \Theta_{n,0,k,\alpha}^\gamma & \Theta_{n,1,k,\alpha}^\gamma & \Theta_{n,j,k,\alpha}^\gamma & \dots & \dots & \Theta_{n,n,k,\alpha}^\gamma \end{bmatrix}, \quad (9)$$

where

$$\Theta_{ij,k,\alpha}^\gamma = \sum_{k=\gamma}^i \sum_{r=0}^i \frac{(-1)^{\gamma+k} \Gamma(j+1) \Gamma(i+\alpha+1) \Gamma(k+\alpha-r+\gamma+1)}{\Gamma(j-r+1) \Gamma(i-k+1) \Gamma(r+1) \Gamma(k+\alpha+1) \Gamma(k-\gamma+1) \Gamma(\alpha+\gamma+1)}.$$

Proof. Leaving the proof as it is very similar to the proof of the above lemma.

Lemma 5. We consider a function $Z(t)$ defined on $[0, \infty]$ and $y(t) = K_M \Psi_M^T(t)$; then

$$Z(t)[{}_0I_t^\gamma y(t)] = K_M Q_{M \times M}^\gamma \Psi_M(t),$$

where $Q_{M \times M}^\gamma$ is the operational matrix, given by

$$\begin{bmatrix} C_{0,0} & C_{0,1} & \dots & C_{0,j} & \dots & C_{0,m} \\ C_{1,0} & C_{1,1} & \dots & C_{1,j} & \dots & C_{1,m} \\ \vdots & \vdots & \vdots & \vdots & \dots & \vdots \\ C_{i,0} & C_{i,1} & \dots & C_{i,j} & \vdots & C_{i,m} \\ \vdots & \vdots & \vdots & \vdots & \dots & \vdots \\ C_{m,0} & C_{m,1} & \dots & C_{m,j} & \dots & C_{m,m} \end{bmatrix},$$

where

$$C_{ij} = \frac{1}{h_i} \int_0^1 \Delta_{i,\gamma,k} Z(t) L_j^\gamma(t) dt,$$

with

$$w_i = \sum_{k=0}^i \frac{(-1)^{i+1} \Gamma(i+1+\gamma)}{\Gamma(k+\gamma+1) \Gamma(1-k+i) \Gamma(k+\gamma)}.$$

Proof. By considering the general term of $\Psi_M(t)$

$${}_0I_1^\gamma L_i(t) = \frac{1}{\Gamma(\gamma)} \int_0^1 (1-s)^{\gamma-1} L_i(s) ds.$$

$${}_0I_1^\gamma L_i(t) = \frac{1}{\Gamma(\gamma)} \int_0^1 (1-s)^{\gamma-1} \sum_{k=0}^i (s)^k \frac{(-1)^k \Gamma(i+1+\gamma)}{\Gamma(-k+1+i) \Gamma(k+1+\gamma) \Gamma(1+k)} ds.$$

$${}_0I_1^\gamma L_i(t) = \sum_{k=0}^i \frac{(-1)^k \Gamma(i+1+\gamma)}{\Gamma(\gamma) \Gamma(-k+1+i) \Gamma(k+1+\gamma) \Gamma(1+k)} \int_0^1 (1-s)^{\gamma-1} (s)^k ds. \quad (10)$$

Using the famous Laplace transform, we have from (10)

$$\begin{aligned} \mathcal{L}\left(\int_0^1 (1-s)^{\gamma-1} s^k ds\right) &= \frac{\Gamma(\gamma) \Gamma(k+1)}{\Gamma(\gamma+k)}. \\ {}_0I_1^\gamma L_i(t) &= \sum_{k=0}^i \frac{(-1)^k \Gamma(i+1+\gamma)}{\Gamma(\gamma) \Gamma(-k+1+i) \Gamma(k+1+\gamma) \Gamma(1+k)} \frac{\Gamma(\gamma) \Gamma(k+1)}{\Gamma(\gamma+k)}. \\ &= \sum_{k=0}^i \frac{(-1)^k \Gamma(i+1+\gamma)}{\Gamma(-k+1+i) \Gamma(k+\gamma+1) \Gamma(1+k)} = \Delta_{i,\gamma,k}. \end{aligned}$$

Now using Laguerre polynomials, we have

$$\Delta_{i,\gamma,k} Z(t) = \sum_{j=0}^m C_{ij} L_j(t),$$

where C_{ij} is calculated by using orthogonality as

$$C_{ij} = \frac{1}{h_i} \int_0^1 \Delta_{i,\gamma,k} Z(t) L_j^\gamma(t) dt. \quad (11)$$

To get the desired result, we evaluate the above (11) relation for $i = 0, 1, \dots, m$ and $j = 0, 1, \dots, m$.

4. Main result

In this section, we discuss some cases of FODEs with initial condition as well as boundary conditions. The approximate solution obtained through desired

method is compared with the exact solution. Similarly we investigate numerical solutions to various coupled systems under some initial conditions as well as boundary conditions.

4.1 Treatment of FODEs under initial and boundary conditions

Here we discuss different cases.

Case 1. In the first case, we consider the fractional order differential equation

$$\begin{cases} {}^c_0D_t^\gamma z(t) \pm z(t) = 0, & 0 < \gamma \leq 1, \\ z(0) = z_0, & z_0 \in R \end{cases} \quad (12)$$

we see that

$${}^c_0D_t^\gamma z(t) = \mathbb{L}_M \psi_M^T(t).$$

and applying ${}_0I_t^\gamma$ by the Lemma 1, on (12) we write

$$z(t) = e_0 + {}_0I_t^\gamma [\mathbb{L}_M \psi_M^T(t)],$$

Using the initial condition to get $e_0 = z_0$ and approximate z_0 as $z_0 \approx F_M \psi_M^T(t)$, Eq. (12) implies

$$\mathbb{L}_M \psi_M^T(t) + \mathbb{L}_M G_{M \times M}^\gamma \psi_M^T(t) + F_M \psi_M^T(t) = 0.$$

Finally the Sylvester-type algebraic equation is obtained as

$$\mathbb{L}_M + \mathbb{L}_M G_{M \times M}^\gamma \psi_M^T(t) + F_M = 0.$$

Solving the Sylvester matrix for $\mathbb{L}_M$, we get the numerical value for $z(t)$.

Example 1.

$$\begin{cases} {}^c_0D_t^\gamma z(t) \pm z(t) = 0, & 0 < \gamma \leq 1, \\ z(0) = 1, & z_0 \in R. \end{cases}$$

Since the exact solution is given by

$$z(t) = E_\gamma(-t^\gamma),$$

where E_γ is the Mittag-Leffler representation, and at $\gamma = 1$, $z(t) = e^{-t}$.

Approximating the solution through the proposed method and plotting the exact as well as numerical solution by using scale $M = 8$ corresponding to $\gamma = 1$ in

Figure 1, we see that the proposed method works very well.

Case 2.

$$\begin{cases} {}^c_0D_t^\gamma z(t) + z(t) = 0, & 1 < \gamma \leq 2, \\ z(0) = z_0, z(1) = z_1, & z_0, z_1 \in R. \end{cases} \quad (13)$$

We take

$${}^c_0D_t^\gamma z(t) = K_M \psi_M^T(t). \quad (14)$$

Figure 1.
Plots of both approximate and exact solution for the Example 1 for Case 1.

Applying Lemma 1 to Eq. (14), we get

$$z(t) = e_0 + e_1(t) + {}_0I_t^\gamma K_M \psi_M^T(t). \quad (15)$$

Using the conditions by putting $t = 0$ and $t = 1$ to get $e_0 = z_0$ and

$$e_1 = z_1 - z_0 - K_M {}_0I_1^\gamma \psi_M^T(t)/_{t=1}.$$

Equation (15) implies

$$z(t) = z_0 + (z_1 - z_0)t - tK_M {}_0I_1^\gamma \psi_M^T(t)/_{t=1} + {}_0I_t^\gamma K_M \psi_M^T(t),$$

where $z_0 + (z_1 - z_0)t$ is the smooth function of t and constants; we approximate it as

$$z_0 + (z_1 - z_0)t \approx G_{M \times M}^\gamma \psi_M^T(t)$$

and

$$tK_M {}_0I_1^\gamma \psi_M^T(1) \approx K_M Q_{M \times M}^\gamma \psi_M^T(t).$$

Hence

$$z(t) = G_{M \times M}^\gamma \psi_M^T(t) - K_M Q_{M \times M}^\gamma \psi_M^T(t) + K_M G_{M \times M}^\gamma \psi_M^T(t)$$

So Eq. (13) implies

$$K_M \psi_M^T(t) + G_{M \times M}^\gamma \psi_M^T(t) - K_M Q_{M \times M}^\gamma \psi_M^T(t) + K_M G_{M \times M}^\gamma \psi_M^T(t) = 0$$

which is further solved for K_M to get the required numerical solution.

For Case 2, we give the following example.

Example 2.

$$\begin{cases} {}^c_0D_t^\gamma z(t) + z(t) = 0, & 0 < \gamma \leq 2, \\ z(0) = -1, \quad z(1) = 1. \end{cases} \quad (16)$$

At $\gamma = 2$, we get the exact solution as of (16) as given by (17)

$$z(t) = 114.58 \sin(x) - \cos(x) \quad (17)$$

Figure 2.
 The plot of exact and approximate solution for Example 2 for Case 2.

Upon using the suggested method, we see from the subplot at the left of **Figure 2** that exact and numerical solutions are very close to each other for very low scale level. Also, the absolute error is given in subplot at the right of **Figure 2**.

4.2 Coupled systems of linear FODEs under initial and boundary conditions

In this subsection, we consider different forms of coupled systems of FODEs with the initials as well as boundary conditions.

Case 1. First we take the coupled system of FODEs as

$$\begin{cases} {}^c_0D_t^\gamma z(t) + az(t) + by(t) = f(t) \\ {}^c_0D_t^\gamma y(t) + cy(t) + dz(t) = g(t), \end{cases} \quad (18)$$

with the conditions

$$z(0) = z_0, \quad y(0) = y_0, z_0, y_0 \in R. \quad (19)$$

Let

$${}^c_0D_t^\gamma z(t) = \mathbf{L}_M \psi_M^T(t), \quad {}^c_0D_t^\gamma y(t) = \mathbf{K}_M \psi_M^T(t). \quad (20)$$

Applying Lemma 1 to Eq. (20), we get

$$\begin{cases} z(t) = e_0 + \mathbf{L}_M G_{M \times M}^\gamma \psi_M^T(t), \\ y(t) = d_0 + \mathbf{K}_M G_{M \times M}^\gamma \psi_M^T(t). \end{cases} \quad (21)$$

Using the initial conditions given in Eq. (19), from Eq. (21), we get

$$\begin{cases} z(t) = F_M^1 \psi_M^T(t) + \mathbf{L}_M G_{M \times M}^\gamma \psi_M^T(t), \\ y(t) = y_0 \approx F_M^2 \psi_M^T(t) + \mathbf{K}_M G_{M \times M}^\gamma \psi_M^T(t). \end{cases} \quad (22)$$

We take approximation as

$$z_0 \approx F_M^1 \psi_M^T(t),$$

and

$$y_0 \approx F_M^2 \psi_M^T(t),$$

while source functions are approximated as

$$f(t) \approx F_M^3 \Psi_M^T(t),$$

and

$$g(t) \approx F_M^4 \Psi_M^T(t).$$

Therefore the consider system on using (19)–(22), (18) becomes

$$\begin{cases} \mathbb{L}_M \psi_M^T + a(F_M^1 \psi_M^T(t) + \mathbb{L}_M G_{M \times M}^\gamma \psi_M^T(t)) \\ + b(F_M^2 \psi_M^T(t) + K_M G_{M \times M}^\gamma \psi_M^T(t)) = F_M^3 \psi_M^T(t). \\ K_M \psi_M^T + c(F_M^2 \psi_M^T(t) + K_M G_{M \times M}^\gamma \psi_M^T(t)) \\ + d(F_M^1 \psi_M^T(t) + \mathbb{L}_M G_{M \times M}^\gamma \psi_M^T(t)) = F_M^4 \psi_M^T(t). \end{cases}$$

On further rearrangement we have

$$\begin{cases} \mathbb{L}_M + a(F_M^1 + \mathbb{L}_M G_{M \times M}^\gamma) + b(F_M^2 + K_M G_{M \times M}^\gamma) = F_M^3 \\ K_M + c(F_M^2 + K_M G_{M \times M}^\gamma) + d(F_M^1 + \mathbb{L}_M G_{M \times M}^\gamma) = F_M^4. \end{cases}$$

which further can be written as

$$\begin{cases} \mathbb{L}_M(I_{M \times M} + aG_{M \times M}^\gamma) + K_M(bG_{M \times M}^\gamma) + (aF_M^1 + bF_M^2 - F_M^3) = 0 \\ K_M(I_{M \times M} + cG_{M \times M}^\gamma) + \mathbb{L}_M(dG_{M \times M}^\gamma) + (cF_M^2 + dF_M^1 - F_M^4) = 0. \end{cases}$$

In matrix form we write as

$$\begin{bmatrix} \mathbb{L}_M & K_M \end{bmatrix} \begin{bmatrix} I_{M \times M} + aG_{M \times M}^\gamma & 0 \\ 0 & I_{M \times M} + cG_{M \times M}^\gamma \end{bmatrix} + \begin{bmatrix} \mathbb{L}_M & K_M \end{bmatrix} \begin{bmatrix} 0 & dG_{M \times M}^\gamma \\ bG_{M \times M}^\gamma & 0 \end{bmatrix} \\ + \begin{bmatrix} aF_M^1 + bF_M^2 - F_M^3 \\ cF_M^2 + dF_M^1 - F_M^4 \end{bmatrix} = 0.$$

We solve this system of matrix equation for $[\mathbb{L}_M \ K_M]$ by using Gaussian's elimination method. The considered system is in the form of $X\bar{A} + X\bar{B} + \bar{C} = 0, .$

$$\text{where } X = [\mathbb{L}_M \ K_M] \bar{A} = \begin{bmatrix} I_{M \times M} + aG_{M \times M}^\gamma & 0 \\ 0 & I_{M \times M} + cG_{M \times M}^\gamma \end{bmatrix}, .$$

$$\bar{B} = \begin{bmatrix} 0 & dG_{M \times M}^\gamma \\ bG_{M \times M}^\gamma & 0 \end{bmatrix} \text{ and } \bar{C} = \begin{bmatrix} aF_M^1 + bF_M^2 - F_M^3 \\ cF_M^2 + dF_M^1 - F_M^4 \end{bmatrix}.$$

Upon computation of matrices $\mathbb{L}_M, K_M$ by using MATLAB®, we put these matrices in Eq. (22) to find z_{app} and y_{app} , respectively.

Example 3. We now provide its example by considering the system of FODEs:

$$\begin{cases} {}^c_0D_t^\gamma z(t) + z(t) + y(t) = f(t) \\ {}^c_0D_t^\gamma y(t) + y(t) + z(t) = g(t), \\ z(0) = 2, \ y(0) = 1. \end{cases}$$

By taking $\gamma = 1$, the exact solution is obtained as

$$z(t) = \cos(t) + e^t, \quad y = \sin(t) + e^{-t},$$

where the external source functions are given by $f(t) = \cos(t) + e^{-t} + 2e^t$ and $g(t) = e^{-t} + \sin(t) + 2\cos(t)$. The exact solution z_{ex}, y_{ex} can be computed by any method of ODEs. Approximating the problem by the considered method, we see that the computed numerical and exact solutions have close agreement at very small-scale level. The corresponding accuracy has been recorded in **Table 1**. Further the comparison between exact and numerical solution and the results about absolute error have been demonstrated in **Figures 3** and **4**, respectively. In **Figure 3** we are given the comparison between exact solution and approximate solutions by using proposed method. Similarly the absolute errors have been described in **Figure 4**.

By comparing the exact and numerical solution through the proposed method, we observe that our numerical solution does not show any disagreement with the exact solution as can be seen in **Figure 3**. The absolute errors $\|z_{app} - z_{ex}\|$ and $\|y_{app} - y_{ex}\|$ plotted at the scale $M = 5$ are very low as given in **Figure 4**, which describes the efficiency of the proposed method.

Case 2. Similarly for the coupled system of FODEs with boundary conditions, we consider

$$\begin{cases} {}^c_0D_t^\gamma z(t) + az(t) + by(t) = f(t), \\ {}^c_0D_t^\gamma y(t) + cy(t) + dz(t) = g(t), \end{cases} \quad (23)$$
$$z(0) = z_0, y(0) = y_0, z(1) = z_1, y(1) = y_1.$$

t	CPU time (s)	Absolute error $\ z_{app} - z_{ex}\ $	Absolute error $\ y_{app} - y_{ex}\ $	CPU time (s)
0	30.5	0.00003	0.000006	32.5
0.15	32.7	0.000016	0.000034	33.3
0.35	35.8	0.000013	0.00003	33.9
0.65	33.6	0.000012	0.00003	35.6
0.87	34.8	0.000018	0.000036	36.5
1	35.9	0.00003	0.000006	36.8

Table 1.
Absolute error at $M = 5$, $\gamma = 0.9$, for different values of t in Example 3.

Figure 3.
Plots of exact and approximate solution of Example 3.

Figure 4.
Plots of absolute error of Example 3.

Let us assume

$$\begin{cases} {}^c_0D_t^\gamma z(t) = \mathbf{L}_M \Psi_M^T(t), \\ {}^c_0D_t^\gamma y(t) = K_M \Psi_M^T(t). \end{cases} \quad (24)$$

Applying Lemma 1 to Eq. (24), we get

$$\begin{cases} z(t) = e_0 + e_1(t) + \mathbf{L}_M G_{M \times M}^\gamma \Psi_M^T(t) \\ y(t) = d_0 + d_1(t) + K_M G_{M \times M}^{\star \gamma} \Psi_M^T(t), \end{cases} \quad (25)$$

where $d_0, d_1, e_0, e_1 \in \mathbb{R}$. Using the initial conditions in Eq. (25), we have $e_0 = z_0$, $d_0 = y_0$. On using boundary conditions, we have from Eq. (25)

$$\begin{aligned} z(1) &= z_0 + e_1 + \mathbf{L}_M G_{M \times M}^\gamma \Psi_M^T(t)|_{t=1}, \\ z(1) - z_0 - \mathbf{L}_M G_{M \times M}^\gamma \Psi_M^T(t)|_{t=1} &= e_1. \end{aligned}$$

Similarly

$$\begin{aligned} y(1) &= y_0 + d_1 + K_M G_{M \times M}^{\star \gamma} \Psi_M^T(t)|_{t=1}, \\ y(1) - y_0 - K_M G_{M \times M}^{\star \gamma} \Psi_M^T(t)|_{t=1} &= d_1. \end{aligned}$$

Equation (25) implies that

$$\begin{cases} z(t) = z_0 + t(z_1 - z_0) - t(\mathbf{L}_M G_{M \times M}^\gamma \Psi_M^T(t)|_{t=1}) + \mathbf{L}_M G_{M \times M}^\gamma \Psi_M^T(t) \\ y(t) = y_0 + t(y_1 - y_0) - t(K_M G_{M \times M}^{\star \gamma} \Psi_M^T(t)|_{t=1}) + K_M G_{M \times M}^{\star \gamma} \Psi_M^T(t). \end{cases} \quad (26)$$

Let $z_0 + t(z_1 - z_0) \approx F_M^1 \Psi_M^T(t)$ and $y_0 + t(y_1 - y_0) \approx F_M^2 \Psi_M^T(t)$, with

$$\begin{aligned} \mathbf{L}_M G_{M \times M}^\gamma \Psi_M^T(t) &= \mathbf{L}_M Q_{M \times M}^{\gamma, z} \Psi_M^T(t) \\ t K_M G_{M \times M}^{\star \gamma} \Psi_M^T(t) &= K_M Q_{M \times M}^{\gamma, y} \Psi_M^T(t). \end{aligned} \quad (27)$$

Hence Eq. (26) implies

$$\begin{cases} z(t) = F_M^1 \Psi_M^T(t) - \mathbf{L}_M Q_{M \times M}^{\gamma, z} \Psi_M^T(t) + \mathbf{L}_M G_{M \times M}^\gamma \Psi_M^T(t) \\ y(t) = F_M^2 \Psi_M^T(t) - K_M Q_{M \times M}^{\gamma, y} \Psi_M^T(t) + K_M G_{M \times M}^{\star \gamma} \Psi_M^T(t). \end{cases} \quad (28)$$

approximating $f(t)$ and $g(t)$ such that

$$\begin{cases} f(t) \approx F_M^3 \Psi_M^T(t) \\ g(t) \approx F_M^4 \Psi_M^T(t). \end{cases} \quad (29)$$

On using (24)–(29), system (23) can be written as

$$\begin{cases} L_M \Psi_M^T(t) + a(F_M^1 \Psi_M^T(t) - L_M Q_{M \times M}^{\gamma, z} \Psi_M^T(t) + L_M G_{M \times M}^{\gamma} \Psi_M^T(t)) \\ + b(F_M^2 \Psi_M^T(t) - K_M Q_{M \times M}^{\gamma, y} \Psi_M^T(t) + K_M G_{M \times M}^{\gamma} \Psi_M^T(t)) - F_M^3 \Psi_M^T(t) = 0 \\ K_M \Psi_M^T(t) + c(F_M^2 \Psi_M^T(t) - K_M Q_{M \times M}^{\gamma, y} \Psi_M^T(t) + K_M G_{M \times M}^{\gamma} \Psi_M^T(t)) \\ + d(F_M^1 \Psi_M^T(t) - L_M Q_{M \times M}^{\gamma, z} \Psi_M^T(t) + L_M G_{M \times M}^{\gamma} \Psi_M^T(t)) - F_M^4 \Psi_M^T(t) = 0. \end{cases}$$

On rearrangement of terms, the above equations give

$$\begin{cases} L_M(I_{M \times M} - aQ_{M \times M}^{\gamma, z} + aG_{M \times M}^{\gamma}) + K_M(I_{M \times M} - bQ_{M \times M}^{\gamma, y} + bG_{M \times M}^{\gamma}) \\ + aF_M^1 + bF_M^2 - F_M^3 = 0 \\ K_M(I_{M \times M} - cQ_{M \times M}^{\gamma, y} + cG_{M \times M}^{\gamma}) + L_M(I_{M \times M} - dQ_{M \times M}^{\gamma, z} + dG_{M \times M}^{\gamma}) \\ + cF_M^2 + dF_M^1 - F_M^4 = 0. \end{cases}$$

In matrix form, we can write

$$\begin{aligned} & \begin{bmatrix} L_M & K_M \end{bmatrix} \begin{bmatrix} I_{M \times M} - aQ_{M \times M}^{\gamma, z} + aG_{M \times M}^{\gamma} & 0 \\ 0 & I_{M \times M} - cQ_{M \times M}^{\gamma, y} + cG_{M \times M}^{\gamma} \end{bmatrix} \\ & + \begin{bmatrix} L_M K_M \end{bmatrix} \begin{bmatrix} 0 & I_{M \times M} - dQ_{M \times M}^{\gamma, z} + dG_{M \times M}^{\gamma} \\ I_{M \times M} - bQ_{M \times M}^{\gamma, y} + bG_{M \times M}^{\gamma} & 0 \end{bmatrix} \\ & + \begin{bmatrix} aF_M^1 + bF_M^2 - F_M^3 \\ cF_M^2 + dF_M^1 - F_M^4 \end{bmatrix} = 0. \end{aligned}$$

We convert the system to algebraic equation by considering

$$\begin{aligned} \bar{L} &= \begin{bmatrix} I_{M \times M} - aQ_{M \times M}^{\gamma, z} + aG_{M \times M}^{\gamma} & 0 \\ 0 & I_{M \times M} - cQ_{M \times M}^{\gamma, y} + cG_{M \times M}^{\gamma} \end{bmatrix} \\ \bar{M} &= \begin{bmatrix} 0 & I_{M \times M} - dQ_{M \times M}^{\gamma, z} + dG_{M \times M}^{\gamma} \\ I_{M \times M} - bQ_{M \times M}^{\gamma, y} + bG_{M \times M}^{\gamma} & 0 \end{bmatrix} \\ \text{and } \bar{N} &= \begin{bmatrix} aF_M^1 + bF_M^2 - F_M^3 \\ cF_M^2 + dF_M^1 - F_M^4 \end{bmatrix}. \end{aligned}$$

so that the system is of the form

$$X\bar{L} + X\bar{M} + \bar{N} = 0,$$

and solving the given equation for the unknown matrix $X = [L_M K_M]$, we get the required solution.

Example 4. As an example, we consider the Caputo fractional differential equation for the coupled system with the boundary conditions as

$$\begin{cases} {}^c_0D_t^\gamma z(t) + 2z(t) - 2y(t) - f(t) = 0, \\ {}^c_0D_t^\gamma y(t) - 3y(t) + 2z(t) - g(t) = 0, \\ z(0) = 4 \quad z(1) = -4, \\ y(0) = 2, \quad y(1) = -2. \end{cases}$$

At $\gamma = 2$, the exact solutions are

$$\begin{cases} z(t) = t^6 + t^5 + t^4 - t^3 + t + 1, \\ y(t) = t^7 - t^6 + t^5 + t^4 + t^3 - t^2 - t + 1. \end{cases}$$

where the source functions are given by

$$\begin{cases} f(t) = -2t^7 + 4t^6 + 30t^4 + 16t^3 + 12t^2 - 2t + 2 \\ g(t) = -3t^7 + 12t^6 + 35t^5 - 27t^4 - 19t^3 + 20t^2 + 9t - 4. \end{cases}$$

We approximate the solution at the considered method by taking scale level $M = 5$. One can see that numerical plot and exact solution plot coincide very well as shown in **Figure 5**. Similarly the absolute error has been plotted at the given scale $M = 5$ in **Figure 6**, which is very low. The lowest value of absolute error $\|z_{app} - z_{ex}\|$ and $\|y_{app} - y_{ex}\|$ indicates efficiency of the proposed method. The table shows the

Figure 5.
Plots of exact and approximate solution for Case 4, boundary value problem.

Figure 6.
Plots of absolute error for Case 4, boundary value problem.

t	Absolute error $\ z_{app} - z_{ex}\ $	CPU time (s)	Absolute error $\ y_{app} - y_{ex}\ $	CPU time (s)
0	0.011	49.4	0.010	50.0
0.15	0.0062	50.3	0.0052	52.5
0.35	0.0058	51.2	0.0047	54.6
0.65	0.006	51.5	0.005	55.5
0.85	0.0075	52.6	0.007	56.4
1	0.011	53.8	0.010	56.2

Table 2.
Absolute error at different values of t for Example 4.

comparison of errors for exact and approximate solutions for fixed scale level $M = 5$ and order $\gamma = 1.9$. Further the absolute error has been recorded at different values of space variable in **Table 2** which provides the information about efficiency of the proposed method.

5. Conclusion

We have successfully used the class of orthogonal polynomials of Laguerre polynomials to establish a numerical method to compute the numerical solution of FODEs and their coupled systems under some initial and boundary conditions. By using these polynomials, we have obtained some operational matrices corresponding to fractional order derivatives and integration. Also we have computed a new matrix corresponding to boundary conditions for boundary value problems of FODEs. Using the aforementioned matrices, we have converted the considered problem of FODEs to Sylvester-type algebraic equations. To obtain the numerical solution, we easily solved the desired algebraic equations by taking help from MATLAB®. Corresponding to the established procedure, we have provided numbers of examples to demonstrate our results. Also some error analyses have been provided along with graphical representations. By increasing the scale level, the accuracy is increased and vice versa. On the other hand, when the fractional order is approaching to integer value, the solutions tend to the exact solutions of the considered FODE. Therefore in each example, we have compared the exact and approximate solution and found that both the solutions were in closure contact with each other. Hence the established method can be very helpful in solving many classes and systems of FODEs under both initial and boundary conditions. In future the shifted Laguerre polynomials can be used to compute numerical solutions of partial differential equations of fractional order.

Author contribution

All authors equally contributed this paper and approved the final version.

Competing interests

We declare that no competing interests exist regarding this manuscript.

IntechOpen

Author details

Adnan Khan¹, Kamal Shah^{1,2*} and Danfeng Luo³

1 Department of Mathematics, University of Malakand, Dir(L),
Khyber Pakhtunkhwa, Pakistan

2 Department of Mathematics and General Sciences, Prince Sultan University,
Riyadh, Saudi Arabia

3 Key Laboratory of Computing and Stochastic Mathematics (Ministry of
Education), School of Mathematics and Statistics, Hunan Normal University,
Changsha, Hunan, P.R. China

*Address all correspondence to: kamalshah408@gmail.com

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Butzer PL, Westphal U. An Introduction to Fractional Calculus. Singapore: World Scientific; 2000
- [2] Samko SG, Kilbas AA, Marichev OI. Fractional Integrals and Derivatives. Switzerland: Gordon and Breach; 1993
- [3] Scalas E, Raberto M, Mainardi F. Fractional calculus and continuous time finance. *Physica A: Statistical Mechanics and its Applications*. 2000; **284**:376-384
- [4] Hilfer R. Applications of Fractional Calculus in Physics. Singapore: World Scientific; 2000
- [5] Amairi M, Aoun M, Najar S, Abdelkrim MN. A constant enclosure method for validating existence and uniqueness of the solution of an initial value problem for a fractional differential equation. *Applied Mathematics and Computation*. 2010; **217**(5):2162-2168
- [6] Deng J, Ma L. Existence and uniqueness of solutions of initial value problems for nonlinear fractional differential equations. *Applied Mathematics Letters*. 2000; **23**:676-680
- [7] Girejko E, Mozyrska D, Wyrwas M. A sufficient condition of viability for fractional differential equations with the Caputo derivative. *Journal of Mathematical Analysis and Applications*. 2011; **38**:146-154
- [8] Baleanu D, Diethelm K, Scalas E, Trujillo JJ. Fractional Calculus Models and Numerical Methods. Singapore: World Scientific; 2009
- [9] Guy J. Modeling fractional stochastic systems as non-random fractional dynamics driven by Brownian motions. *Applied Mathematical Modelling*. 2008; **32**:836-859
- [10] Sabatier JATMJ, Agrawal OP, Machado JT. Advances in Fractional Calculus. Dordrecht: Springer; 2007
- [11] Kilbas AA, Srivastava HM, Trujillo JJ. Theory and Applications of Fractional Differential Equations. Amsterdam: Elsevier; 2006
- [12] Lakshmikantham I, Leela S. Theory of Fractional Dynamical Systems. Cambridge, UK: Cambridge Scientific Publishing; 2009
- [13] Li CP, Deng WH. Remarks on fractional derivatives. *Applied Mathematics and Computation*. 2007; **187**:777-784
- [14] Li CP, Dao XH, Guo P. Fractional derivatives in complex planes. *Nonlinear Analysis: Theory Methods & Applications*. 2009; **71**:5-6
- [15] Li C, Gong Z, Qian D, Chen Y. On the bound of the Lyapunov exponents for the fractional differential systems. *Chaos: An Interdisciplinary Journal of Nonlinear Science*. 2010; **20**(1):013127
- [16] Oldham KB, Spanier J. The Fractional Calculus. New York: Acad. Press; 1974
- [17] Ortigueira MD. Comments on modeling fractional stochastic systems as non-random fractional dynamics driven Brownian motions. *Applied Mathematical Modelling*. 2009; **33**:2534-2537
- [18] Qian DL, Li CP, Agarwal RP, Wong PJY. Stability analysis of fractional differential system with Riemann-Liouville derivative. *Mathematical and Computer Modelling*. 2010; **52**:862-874
- [19] West BJ, Bologna M, Grigolini P. Physics of Fractional Operators. New York: Springer; 2003

- [20] Yang S, Xiao A, Su H. Convergence of the variational iteration method for solving multi-order fractional differential equations. *Computers & Mathematics with Applications*. 2010; **60**:2871-2879
- [21] Ray SS, Bera RK. Solution of an extraordinary differential equation by adomian decomposition method. *Journal of Applied Mathematics*. 2004; **4**:331338
- [22] Hashim I, Abdulaziz O, Momani S. Homotopy analysis method for fractional IVPs. *Communications in Nonlinear Science and Numerical Simulation*. 2009; **14**:674-684
- [23] Bengochea G. Operational solution of fractional differential equations. *Applied Mathematics Letters*. 2014; **32**: 48-52
- [24] Khalil H, Khan RA. The use of Jacobi polynomials in the numerical solution of coupled system of fractional differential equations. *International Journal of Computer Mathematics*. 2015; **92**(7): 1452-1472
- [25] Doha EH, Bhrawy AH, Ezz-Eldien SS. Efficient Chebyshev spectral methods for solving multi-term fractional orders differential equations. *Applied Mathematical Modelling*. 2011; **35**:5662-5672
- [26] Esmaeili S, Shamsi M, Luchko Y. Numerical solution of fractional differential equations with a collocation method based on Muntz polynomials. *Computers & Mathematics with Applications*. 2011; **62**:918-929
- [27] Odibat Z, Momani S, Erturk VS. Generalized differential transform method an application to differential equations of fractional order. *Applied Mathematics and Computation*. 2008; **197**:467-477
- [28] Baleanu D, Mustafa OG, Agarwal RP. An existence result for a superlinear fractional differential equation. *Applied Mathematics Letters*. 2010; **23**:1129-1132
- [29] Baleanu D, Mustafa OG, Agarwal RP. On the solution set for a class of sequential fractional differential equations. *Journal of Physics A*. 2010; **43**:385-209
- [30] Doha EH, Abd-Elhameed WM. Efficient solutions of multidimensional sixth-order boundary F value problems using symmetric generalized Jacobi-Galerkin method. *Abstract and Applied Analysis*. 2012; **2012**:12
- [31] Bhrawy AH, Al-Shomrani MM. A Jacobi dual-Petrov Galerkin-Jacobi collocation method for solving Korteweg-de Vries equations. *Abstract and Applied Analysis*. 2012; **2012**:14
- [32] Singh AK, Singh VK, Singh VK. The Bernstein operational matrix of integration. *Applied Mathematical Sciences*. 2009; **3**:2427-2436
- [33] Bhrawy AH, Alofi AS, Ezz-Eldien SS. A quadrature tau method for fractional differential equations with variable coefficients. *Applied Mathematics Letters*. 2011; **24**:2146-2152
- [34] Bhrawy AH, Mohammed MA. A shifted Legendre spectral method for fractional-order multi-point boundary value problems. *Advances in Difference Equations*. 2012; **2012**:8
- [35] Khalil H, Khan RA. New operational matrix of integration and coupled system of Fredholm integral equations. *Chinese Journal of Mathematics*. 2014; **16**:12
- [36] Khan RA, Khalil H. A new method based on Legendre polynomials for solution of system of fractional order partial differential equations.

International Journal of Computer
Mathematics. 2014;**91**(12):2554-2567

[37] Khalil H, Khan RA. A new method
based on Legendre polynomials for
solutions of the fractional two-
dimensional heat conduction equation.
Computers & Mathematics with
Applications. 2014;**67**:1938-1953

[38] Guo BY, Wang LL. Modified
Laguerre pseudospectral method refined
by multidomain Legendre
pseudospectral approximation. Journal
of Computational and Applied
Mathematics. 2006;**190**:304-324

[39] Gulsu M, Gurbuz B, Ozturk Y,
Sezer M. Laguerre polynomial approach
for solving linear delay difference
equations. Applied Mathematics and
Computation. 2011;**217**:6765-6776

[40] Bhrawy AH, Taha TM,
Machado JAT. A review of operational
matrices and spectral techniques for
fractional calculus. Nonlinear
Dynamics. 2015;**81**(3):1023-1052

[41] Diethelm K, Ford NJ. Numerical
solution of the Bagley Torvik equation.
BIT Numerical Mathematics. 2002;
42(1):490-500

[42] Akyuz-Dascioglu A, Isler N.
Bernstein collocation method for solving
nonlinear differential equations.
Mathematical and Computational
Applications. 2013;**18**:293-300

[43] Shah K. Using a numerical method
by omitting discretization of data to
study numerical solutions for boundary
value problems of fractional order
differential equations. Mathematical
Methods in the Applied Sciences. 2019;
42:6944-6959. DOI: 10.1002/mma.5800