We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Microorganisms as Biocatalysts and Enzyme Sources

Arturo Cano-Flores, Javier Gómez,

Iker S. Escalona-Torres and Benjamín Velasco-Bejarano

Abstract

Microbial-catalyzed biotransformations have considerable potential for the generation of an enormous variety of structurally diversified organic compounds, especially natural products with complex structures like triterpenoids, flavonoids, steroids, steroidal saponins, and sesquiterpenoids. They offer efficient and economical ways to produce semisynthetic analogues and novel lead molecules. Microorganisms such as bacteria and fungi could catalyze *chemo*-, *regio*-, and *stereospecific* hydroxylations of diverse substrates that are extremely difficult to produce by chemical routes. During recent years, considerable research has been performed on the microbial transformation of bioactive compounds, in order to obtain biologically active molecules with diverse structural features. In green chemistry, biotransformations are an important chemical methodology toward more sustainable industrial processes.

Keywords: microorganisms, fungi, bacteria, microbial transformation, natural products, enzymes

1. Introduction

Microbial transformation is regarded as an enzymatic reaction by using the metabolic activities of microorganisms to modify the chemical structures of bioactive substrates for finding the new chemical derivatives with the potent bioactivities and physical-chemical characteristics. It has a number of advantages over chemical synthesis such as higher *stereo*- and *regio*selectivity but is also enantiospecific, allowing the production of chiral products from racemic mixtures. The conditions for biotransformations are mild, and in the majority of cases, they do not require the protection of pre-existing functional groups. Furthermore, some reactions that do not occur when using chemical approaches are easily carried out by microbial transformation. Microbial factories show advantages, for instance, growing rapidly and ease of large-scale production [1–3].

The use of microorganisms may be a highly efficient method of production of these compounds. The reactions involved in biotransformation of organic compounds by whole cells of various microorganisms include oxidation, reduction, hydroxylation, esterification, methylation, demethylation, isomerization, hydrolysis, glycosylation, and hydrogenation [4, 5].

Biotransformation may be carried out with isolated enzyme systems or with intact organism. Although isolated enzyme systems may be more specific and

efficient for certain biotransformation, these reactions may involve isolating the enzyme system, and, for some classes of enzyme-catalyzed reaction, a recycling sequence may be required to regenerate the enzyme [6].

Fungi are playing a prominent role in the catalysis of organic compounds and in the production of commercially and industrially important compounds, because of their ability to catalyze novel reactions [7]. Fungi are commonly used in the industry for production of fermented beverages, foods, physiologically active substances, solvents, organic acids, polysaccharides, antibiotics, etc. Of the zygomycota, *Mucor* and *Rhizopus* are commonly used in the industry. *Rhizopus* strains are important in citric acid production. *Mucor* strains make a significant number of important lipases and catalyze the hydroxylation of a wide range of chemical compounds [2–4].

The use of the microbial model offers a number of advantages over the use of animals in metabolism studies, mainly: (1) simple, easy, and can be prepared at low cost; (2) screening for a large number of strains is a simple repetitive process; (3) the large number of metabolites formed allows easier detection, isolation, and structural identification; (4) newer metabolites can be isolated; (5) utilized for synthetic reactions involving many steps; (6) useful in cases where *regio*- and *stereospecificity* is required; (7) maintenance of stock cultures of microorganisms is simpler and cheaper than the maintenance of cell or tissue cultures or laboratory animals; (8) ease of setup and manipulation; and (9) more reliable and reproducible [8, 9].

The objective of this review is to highlight the importance of microorganisms or enzymes isolated from them in the biotransformation process of natural products or xenobiotic compounds, according to green chemistry or white biotechnology.

2. Microbiological transformations of some selected natural products with different microorganisms

2.1 Sesquiterpene lactone

Artemisinin (**1**), a sesquiterpene lactone endoperoxide and an antimalarial drug, is effective against chloroquine-resistant parasites; but its toxicities and low solubility in water hamper its therapeutical use. Studies on modification of **1** through biological and chemical methodologies have been reported to yield more effective and water-soluble derivatives. A wide array of microbial transformations of **1** involve oxidation, reduction, and degradation reactions by different microorganisms, such as *Aspergillus niger*, *A. flavus*, *A. adametzi* (ATCC 10407), *Cunninghamella echinulata*, *Caenorhabditis elegans*, *Mucor polymorphous*, *M. rammanianus*, *Streptomyces griseus*, *Penicillium simplicissimum*, *P. chrysogenum*, *P. purpuresceus*, *Pestalotiopsis guepini* (P-8), *Eurotium amstelodami*, *Trichoderma viride* (T-58), *Saccharomyces cerevisiae*, and *Pichia pastoris*. Biotransformation of **1** usually includes the processes such as hydroxylation of methyl, methine and methylene groups, deoxidation reactions, hydration and acetylation reactions, epimerization, and breakdown of heterocyclic rings (**Table 1**).

2.2 Triterpene

Ursolic acid (3 β -hydroxy-urs-12-en-28-oic acid, UA, **2**), a natural pentacyclic triterpene, is broadly used in food, cosmetics, and biomedical industries. As a ubiquitous constituent in the plant kingdom and the major component of many traditional medicine herbs, ursolic acid remarkably exhibits a lot of biological

Microorganism	Products	Action	Reference
<i>A. niger</i>	3 β -hydroxy-4,12-epoxy-1-deoxyartemisinin Artemisinin G 3,13-epoxyartemisinin 4 α -hydroxy-1-deoxyartemisinin	Epoxidation, hydroxylation C-3 β site Endoperoxide function reduction Breakdown of heterocyclic rings Epoxidation C-3 and C-13 Hydroxylation C-4, endoperoxide function reduction	[10]
<i>A. flavus</i> (MTCC 9167)	14-hydroxyartemisinin Artemisinin G 4 α -hydroxydeoxyartemisinin Deoxyartemisinin	Hydroxylation of C-14 site Breakdown of heterocyclic rings Hydroxylation of C-4 α site Endoperoxide function reduction	[11, 12]
<i>C. elegans</i> (ATCC 9245)	7 β -hydroxy-9 α -artemisinin 4 α -hydroxy-1-deoxyartemisinin 7 β -hydroxyartemisinin 6 β -hydroxyartemisinin 7 α -hydroxyartemisinin 6 β ,7 α -dihydroxyartemisinin	Hydroxylation C-7 β site Epimerization C-9 Hydroxylation C-4 α site Hydroxylation C-7 β site Hydroxylation C-6 β site Hydroxylation C-7 α site Hydroxylation C-6 β and C-7 α sites	[13, 14]
<i>P. simplicissimum</i>	9 β -acetoxyartimisinin 9 α -hydroxyartemisinin	Acetylation of C-9 β site Hydroxylation of C-9 α site	[15]
<i>R. stolonifer</i>	Deoxyartemisinin 1 α -hydroxyartemisinin 10 β -hydroxyartemisinin	Endoperoxide function reduction Hydroxylation of C-1 site	[16]
<i>S. griseus</i> (ATCC 13273)	9-artemisitone 9 α -hydroxyartemisinin 9 β -hydroxyartemisinin 3 α -hydroxyartemisinin	Oxidation of C-9 site Hydroxylation of C-9 α site Hydroxylation of C-9 β site Hydroxylation of C-3 α site	[17]
<i>N. corallina</i>	Deoxyartemisinin	Endoperoxide function reduction	[11]

Table 1.
Products obtained from the biotransformation of artemisinin (1) by different microorganisms.

activities, such as antibacterial, anti-allergic, antioxidative, anti-inflammation, and antitumor activities [18].

Microbial transformation of ursolic acid (2) by *Bacillus megaterium* CGMCC 1.1741 yielded five metabolites identified as 3-oxo-urs-12-en-28-oic acid (3, 6.2%); 1 β ,11 α -dihydroxy-3-oxo-urs-12-en-28-oic acid (4, 13.5%); 1 β -hydroxy-3-oxo-urs-12-en-28,13-lactone (5, 5.0%); 1 β ,3 β ,11 α -trihydroxy-urs-12-en-28-oic acid (6, 26.9%); and 1 β ,11 α -dihydroxy-3-oxo-urs-12-en-28-O- β -D-glucopyranoside (7, 8.6%) [19]. The biotransformation studies of 2 by *Alternaria longipes* AS 3.2875 have led to the isolation of six products of hydroxylation or glycosylation. Their structures were identified as 3-carbonyl-ursolic acid-28-O- β -D-glucopyranosyl ester (8), ursolic acid-3-O- β -D-glucopyranoside (9), ursolic acid-28-O- β -D-glucopyranosyl ester (10), 2 α ,3 β -dihydroxy-ursolic acid-28-O- β -D-glucopyranosyl ester (11), 3 β ,21 β -dihydroxy-ursolic acid-28-O- β -D-glucopyranosyl ester (12), and 3-O-(β -D-glucopyranosyl)-ursolic acid-28-O-(β -D-glucopyranosyl) ester (13). Glycosylation reaction on pentacyclic triterpenoid fulfilled with difficulty in the process of chemical synthesis is facile by microbial transformation [20]. Biotransformation of 2 by *A. alternata* eight metabolites were found to be 2 α ,3 β -dihydroxyurs-12-en-28-oic acid (corosolic acid, 14), urs-12-en-2 α ,3 β ,28-triol (15), 3 β ,23-dihydroxyurs-12-en-28-oic acid (16), 2 α ,3 β ,23-trihydroxyurs-12-en-28-oic acid (17), 2 α ,3 β ,23,24-tetrahydroxyurs-12-en-28-oic acid (18), 3 β ,28-dihydroxy-12-ursene (19), urs-12-en-3 β -ol (20), and urs-12-en-2 α ,3 β -diol (21). The reduction of the C-28 carboxyl group and hydroxylation at C-2, 23, and 24 are steps in the metabolic pathway of 2 [21].

Biotransformation of UA by *S. racemosum* (3.2500) yielded five metabolites 3 β ,7 β ,21 β -trihydroxy-urs-12-en-28-oic acid (22); 3 β ,21 β -dihydroxy-urs-11-en-28-oic acid-13-lactone (23); 1 β ,3 β ,21 β -trihydroxy-urs-12-en-28-oic acid (24); 3 β ,7 β ,21 β -trihydroxy-urs-1-en-28-oic acid-13-lactone (25); and 1 β ,3 β -dihydroxy-urs-12-en-21-oxo-28-oic acid (26) which were afforded [22]. Additionally, of the biotransformation of 2 with by *S. racemosum* compounds 27–30 and 11,26-epoxy-3 β -21 β -dihydroxy-urs-12-en-28-oic acid were obtained (31), (Figure 1) [23].

The endophytic fungi *Pestalotiopsis microspora* isolated from medical plant *Huperzia serrata* can transform 1 to afforded 3-oxo-15 β ,30-dihydroxy-urs-12-en-28-oic acid (32), 3 β ,15 β -dihydroxy-urs-12-en-28-oic acid (33), 3 β ,15 β ,30-trihydroxy-urs-12-en-28-oic acid (34), and 30 [24].

Microbial transformation of ursolic acid by *Mucor spinosus* AS 3.3450 were isolated and their structures were identified as 9, 22 and 3 β ,7 β -dihydroxy-ursolic acid-28-ethanone (35) (Figure 1) [25].

The gum resin *Boswellia serrata* has been used for the treatment of inflammatory and arthritic diseases. Its major active constituents are ursane triterpenoids, which include 11-keto- β -boswellic acid (KBA, 36), β -boswellic acid (BA), and acetyl- β -boswellic acid (ABA). Microbial transformation 36 by *Cunninghamella blakesleeana* (AS 3.970) yielded ten regioselective transformed products: 7 β -hydroxy-11-keto- β -boswellic acid (37), 7 β ,15 α -dihydroxy-11-keto- β -boswellic acid (38), 7 β ,16 β -dihydroxy-11-keto- β -boswellic acid (39), 7 β ,16 α -dihydroxy-11-keto- β -boswellic acid (40), 7 β ,22 β -dihydroxy-11-keto- β -boswellic acid (41), 7 β ,21 β -dihydroxy-11-keto- β -boswellic acid (42), 7 β ,20 β -dihydroxy-11-keto- β -boswellic acid (43), 7 β ,30-dihydroxy-11-keto- β -boswellic acid (44), 3 α ,7 β -dihydroxy-11-oxours-12-en,24,30-dioic acid (45), and 3 α ,7 β -dihydroxy-30-(2-hydroxypropanoyloxy)-11-oxours-12-en, 24-oic acid (46). Bioconversion of 36 with *Bacillus megaterium* based on a recombinant cytochrome P450 system yielded *regio*- and *stereoselective* 15 α -hydroxylation (47) of substrate (Figure 2) [26].

Figure 1.
Biotransformation products of ursolic acid (2).

Figure 2.
Biotransformation products of 11-keto-b-boswellic acid (36).

18 β -glycyrrhetic acid (48) is the active form of glycyrrhizin which is the major pentacyclic triterpene found in licorice (*Glycyrrhiza glabra* L.). Glycyrrhetic acid has been shown to possess several pharmacological activities, such as antiulcerative, anti-inflammatory, immunomodulating, antitumor, antiviral, antihepatitis effects, and anticancer. Biotransformation 48 with a fungus *C. blakesleeana* (AS 3.970) yielded 3-oxo-7 β -hydroxyglycyrrhetic acid (49) and 7 β -hydroxyglycyrrhetic acid (50) [27], while of 48 using *Absidia pseudocylindrospora* (ATCC 24169), *Gliocladium viride* (ATCC 10097) and *Cunninghamella echinulata* (ATCC 8688a) afforded seven derivatives: 51, 52, 7 β ,15 α -dihydroxy-18 β -glycyrrhetic acid (53), 15 α -hydroxy-18 β -glycyrrhetic acid (54), 1 α -hydroxy-18 β -glycyrrhetic acid (55) and 13 β -hydroxy-7 α ,27-oxy-12-dihydro-18 β -glycyrrhetic acid (56), and the epimer of compound 53 on C-17 (Figure 3) [28].

Ginsenoside Rb1 (61) is the most predominant protopanaxadiol-type ginsenoside in *Panax* species (ginseng). Several microbial transformations of this substrate (Ginsenoside Rb1) have been accomplished with an ample and varied group of microorganisms, all of these having β -glucosidase activities.

Figure 3.
Biotransformation products of 18 β -glycyrrhetic acid (48).

Figure 4.

Triterpenic acid: ginsenoside Rb1 (**61**), oleanolic acid (**62**), betulinic (**63**) and betulonic acid (**64**).

Deglycosylation appears to be the major transformation pathway, and the intermediate and the final hydrolysis products of **61** depended on the microorganisms used. The biotransformation of various triterpenes, such as **61–64**, has been described in the literature. For each triterpenoid, the transforming microorganism together with the type and site of the reaction catalyzed is given in **Table 2** (**Figure 4**) [29].

Biotransformation of oleanolic acid (**62**) with *Bacillus subtilis* (ATCC 6633) resulted in five more polar metabolites as 28-O- β -D-glucopyranosyl oleanic acid (**63**), 3-O- β -D-glucopyranosyl oleanic acid (**64**), 3-O-(β -D-glucopyranosyl)-oleanic acid-28-O- β -D-glucopyranoside (**61**), 24-hydroxyl-oleanolic acid (**62**), and 3 β -24-dihydroxy-olean-12-en-28-O- β -D-glucopyranosyl-oic acid (**63**), while echinocystic acid (**64**, 250 mg) was metabolized to three more polar metabolites as 28-O- β -D-glucopyranosyl echinocystic acid (**65**), 3-O-(β -D-glucopyranosyl)-echinocystic acid-28-O- β -D-glucopyranoside (**66**), and 24-hydroxyl-28-O- β -glucopyranosyl echinocystic acid (**67**), and then biotransformation of betulinic acid (**68**) contributed four metabolites as 28-O- β -D-glucopyranosyl betulinic acid (**69**), 3-O-(β -D-glucopyranosyl)-betulinic acid-28-O- β -D-glucopyranoside (**70**), 23-hydroxy-betulinic acid (**71**), and 23-hydroxy-28-O- β -D-glucopyranosyl betulinic acid (**72**). In this way there were two types of reactions in the biotransformation of triterpenic acids **58**, **64**, and **68**: hydroxylation and glycosylation [41]. Biotransformation of **58** by *C. muscae* yielded nine hydroxylated and glycosylated metabolites. The specific hydroxylation (7 β , 15 α , and 21 β) was main reaction type. In addition, the selective glycosylation at C-28 was another main reaction type. It was also observed that the 3 β -OH group was selectively dehydrogenated into carbonyl group [42].

A C-3 oxidized derivative of oleanolic acid **73** (3-oxoolean-12-en-28-oic acid) was transformed by the *Chaetomium longirostre* (RF-1095) into 4-hydroxy-3,4-seco-olean-12-ene-3,28-dioic acid (**74**) and the corresponding 21-hydroxylated derivative (**75**). Analogous ring-A cleavage oxidation reactions have been observed in the biotransformation of triterpenoid substrates with the fungi *Septomyxa affinis* ATCC 6737 and *Glomerella fusarioides* ATCC 9552. (**Figure 5**) [4, 43].

2.3 Steroidal saponins

Diosgenin [(25R)-spirost-5-en-3 β -ol, **76**] is an important natural starting material in the pharmaceutical industry to produce steroid drugs and hormones since the last century. In recent years, a wide array of new biological activities of **76** has been disclosed. Diosgenin was subjected to several structural modification studies to secure new derivatives via microbial transformation. Several microorganisms have been found to be capable of degrading **76**, *Bacillus megaterium*, *Corynebacterium mediolanum*, *Mycobacterium fortuitum*, *M. phlei*, *Nocardia rhodochrous*, and *F. solani*.

Triterpenoid	Microorganism	Reaction	Reference
Ginsenoside Rb ₁ (61)	<i>A. niger</i> (KTC 6909)	Deglycosylation at the C-3 and C-20 sites	[30]
	<i>A. niger</i> (AS 3.1858)	Deglycosylation at the C-3 and C-20 sites	[30]
	<i>A. usamii</i> (KTC 6956)	Deglycosylation at the C-3 and C-20 sites	[30]
	<i>F. sacchari</i>	Deglycosylation at the C-3 and C-20 sites	[31]
	<i>P. oxalicum</i>	Deglycosylation at the C-3 site	[32]
	<i>C. lunata</i> (AS 3.1109)	Deglycosylation at the C-20 site, hydration $\Delta^{24(25)}$ Formation of tertiary alcohol	[33]
	<i>R. stolonifer</i> (AS 3.822)	Deglycosylation at the C-3 and C-20 sites	[33]
Oleanolic acid (62)	<i>C. blakesleeana</i>	Diverse hydroxylation at the C-1 β , C-7 β , C-13 β sites	[4]
	<i>F. lini</i>	Dehydrogenation C-13 and C-18. oleanderolide formation	[4]
	<i>P. chrysogenum</i>	Hydroxylation on C-21. Oxidation of the hydroxyl group in C-3	[4]
	<i>C. phomoides</i>	Hydroxylation in C-6 β	[4]
	<i>A. ochraceus</i> (NG 1203)	Hydroxylation in C-11 α	[4]
	<i>Chaetomium longirostre</i>	Oxidative ring A cleavage, hydroxylation at the C-21 β sites	[34]
	<i>Nocardia</i> sp. (NRRL 5646)	Methyl esterification of the C-28 carboxyl group	[4]
	<i>R. miehei</i> (CECT 2749)	Hydroxylation of the C-7 β , C-15 α and C-30 sites Deshydrogenation $\Delta^{9(11)}$	[35]
Betulinic acid (63)	<i>B. megaterium</i> (ATCC 14581)	Dehydrogenation of the C-3 secondary alcohol group, hydroxylation at the C-6 α and C-7 β sites	[36]
	<i>B. megaterium</i> (ATCC 13368)	Dehydrogenation of the C-3 secondary alcohol group, hydroxylation at the C-7 β and C-15 α sites	[37]
	<i>C. elegans</i> (ATCC 9244)	Hydroxylation at the C-1 β and C-7 β sites	[4]
	<i>Cunninghamella</i> sp.	Introduction of a β -glucopyranosyl at the C-28 carboxylic acid group	[38]
Betulonic acid (64)	<i>B. megaterium</i> (ATCC 13368)	Ketone α -hydroxylation at the C-2 site	[37]
	<i>Ch. longirostris</i>	Oxidative ring A cleavage, hydroxylation, decarboxylation	[39]
	<i>C. lunata</i> (ATCC 13432)	Hydroxylations at the C-7 β and (or) C-15 β sites	[4, 40]

Table 2.
Examples of biotransformed triterpenes (61–64) with different microorganisms.

Three major products were accumulated, diosgenone (77), 1-dehydrodiosgenone (78), androst-4-en-3,17-dione (AD, 79), and androsta-1,4-diene-3,17-dione (ADD, 80) (Table 3) [44, 45]. In addition, two side-chain cleavage intermediates of 76 were produced by *C. elegans* and *Aspergillus nidulans*. Microbial transformation

Figure 5.
Biotransformation products of oleanolic acid (62), echnocystic acid (68) and betulinic acid (63).

Fungi	Diosgenona (77)	AD (79)	ADD (80)	Progesterone (102)	16-AD
<i>A. nidulans</i>	++	++	++	++	
<i>C. elegans</i>	++			++	
<i>F. solani</i>	++				++
<i>Rhizopus</i> sp.	++	++	++		

AD androst-4-ene-3,17-dione; ADD androsta1,4-diene-3,17-dione; 16-AD androst-4-ene-3,16-dione.

Table 3.
The ability of different fungi to transform diosgenin (76).

of 76 using white-rot fungus *Coriolus versicolor* afforded eight polyhydroxylated steroids, 7 β -hydroxydiosgenin (81), (25R)-spirost-5-en-3 β ,7 β ,21-triol (82), (25R)-spirost-5-en-3 β ,7 β ,12 β -triol (83), (25R)-spirost-5-en-3 β ,7 α ,15 α ,21-tetraol (84), (25R)-spirost-5-en-3 β ,7 β ,12 β ,21-tetraol (85), (25R)-spirost-5-en-3 β ,7 α ,12 β ,21-tetraol (86), and (25R)-spirost-5-en-3 β ,7 β ,11 α ,21-tetraol (87). The 3 β -hydroxyl group and double bond in the B-ring of 76 were found to be important structural determinants for their activity [46].

Microbial transformation of 76 using *Cunninghamella blakesleeana* AS 3.970 afforded polyhydroxylated derivatives, such as (25R)-spirost-5-en-3 β ,7 α ,12 β -triol (88), (25R)-spirost-5-en-3 β ,7 α ,12 β ,15 α ,21-pentaol (89), (25R)-spirost-5-en-3 β ,7 α ,12 β ,18-tetraol (90), (25R)-spirost-5-en-3 β ,7 α ,12 β ,15 α -tetraol (91), (25R)-spirost-5-en-3 β ,7 α ,11 α ,21-tetraol (92), (25R)-spirost-5-en-3 β ,7 β ,15 α ,21-tetraol (93), and (25R)-spirost-5-en-3 β ,7 β ,12 β ,18-tetraol (94) [47], specifically, the hydroxylation, ketonization, and methoxylation by *Cunninghamella blakesleeana*, *C. elegans*, *Helicostylum piriforme*, and *Streptomyces virginiae*, at C-7, C-9, C-11, C-12, and C-25 positions of 76. Biotransformation of 76 by *Syncephalastrum racemosum* afforded (25R)-spirost-5-en-3 β ,7 α ,9 α -triol (95, 1%), (25R)-spirost-5-en-3 β ,9 α ,12 α -triol-7-one (96, 2%), (25R)-spirost-5-en-3 β ,9 α -diol-7,12-dione (97, 1.5%), (25R)-spirost-4-en-9 α ,12 β ,14 α -triol-3-one (98, 0.66%), and (25S)-spirost-4-en-9 α ,14 α ,25 β -triol-3-one (99, 0.66%) [48]. *C. echinulata* (CGMCC3.2716) metabolized 76 to afford 81 (0.9%), 83 (7.7%), (25R)-spirost-5-en-3 β ,7 β -diol-11-one (100, 7, 1.5%), and (25R)-spirost-5-en-3 β ,7 β ,11 α -triol (101, 6.2%) (Figure 6) [49].

2.4 Steroids

Microorganisms are able to hydroxylate steroids in different positions C-1 to C-21. These represent the most widespread type of steroid bioconversion carried

Figure 6.
 Biotransformation products of diosgenin (76).

out by fungi. The commercialized microbial process in the steroid field was in the production of 11 α -hydroxyprogesterone. This process was realized for the first time by Peterson and Murray (1952), which patented this process of 11 α -hydroxylation of progesterone (102) by *Rhizopus* species [50]. Microbial hydroxylation of 102 by *A. griseola* produced two hydroxylated pregnane identified as 6 β ,14 α -dihydroxyprogesterone (103) and 7 α ,14 α -dihydroxyprogesterone (104). *R. pusillus* produced 6 β ,11 α -dihydroxyprogesterone (105) with excellent yield (65.5%) and 7 α ,14 α -dihydroxyprogesterone (106) (Figure 7) [51].

Industry, which is carried by different microorganisms, such as different species of *Curvularia* spp., *Cunninghamella* spp. and fungi *Trichoderma hamatum*, *Cochliobolus lunatus*. Structural transformation of steroidal compounds through microorganisms has emerged as an important application in the steroidal drug industry. Microbial conversions of steroids generally involve dehydrogenation, esterification, halogenation, isomerization, methoxylation, and side-chain modification of steroidal skeleton. Recently, *Mucor circinelloides lusitanicus* transformed 5-en-3 β -ol steroids (108 and 109) into di- and trihydroxy products. The compound

Figure 7.
 Biotransformation products of progesterone (102).

Figure 8.
 Biotransformation products of 5-en-3 β -ol steroids.

108 yielded 3 β ,7 α ,11 α -trihydroxypregna-5-en-20-one (**110**, 46.4%), and **109** afforded **111** (3 β ,7 α -dihydroxyandrost-5-en-17-one, 43.6%) (**Figure 8**) [52].

Microbial transformation of (20*S*)-20-hydroxymethylpregna-1,4-dien-3-one (**112**) is by four filamentous fungi, *Cunninghamella elegans* (**113–119**), *Macrophomina phaseolina* (**115**, **117**, **120–122**), *Rhizopus stolonifer* (**113**, **123**), and *Gibberella fujikuroi* (**115–117**, **123**). These metabolites were obtained as a result of biohydroxylation of **112** at C-6 β , 7 β , 11 α , 14 α , 15 β , 16 β , and 17 α positions (**Figure 9**) [53].

The 11 α -, 11 β -, 15 α , and 16 α -hydroxylations are currently established processes in the steroid industry mainly for the production of adrenal cortex hormones and their analogues. 11 α -, 11 β -, and 16 α -hydroxylations are usually performed using *Rhizopus* spp. or *Aspergillus* spp., *Curvularia* spp. or *Cunninghamella* spp. and *Streptomyces* spp., respectively (**Figure 10**) (**Table 4**) [54].

Boldenone (**124**) is an important steroid hormone drug which is the derivative of testosterone. Biotransformation of **124** by *Arthrobacter simplex* and recombinant *Pichia pastoris* with 17 β -hydroxysteroid dehydrogenase from *Saccharomyces cerevisiae* produces BD (**124**) from androst-4-ene-3,17-dione (**79**, AD) efficiently [65]. Many microorganisms such as *Mucor racemosus*, *Nostoc muscorum*, and *Arthrobacter oxydans* can utilize androst-1,4-diene-3,17-dione (**80**, ADD) as substrate to produce testosterone through 17 β -carbonyl reduction reactions (**Table 5**). The ability of

Figure 9.
 Biotransformation products of (20*S*)-20-hydroxymethylpregna-1, 4-dien-3-one (**112**).

Figure 10.
The ability of different microorganism to transform progesterone (102).

Hydroxylation sites	Microorganisms	Applications	Reference
C-7α	<i>Fusarium</i> sp., <i>Gibberella</i> sp., <i>Nigrospora</i> sp., <i>Acremonium</i> sp., <i>Phycomyces</i> sp.	Production of bile acids and drugs for neuropsychiatry and immunology	[55, 56]
C-7β	<i>Mortierella</i> sp.	Obtaining drugs for prostate cancer	[56, 57]
11β	<i>Curvularia</i> sp., <i>Absidia</i> sp., <i>Cunninghamella</i> sp., <i>Trichoderma</i> sp., <i>Cochliobolus</i> sp.	Obtaining anti-inflammatory drugs, like hydrocortisone, prednisone acetate, dexamethasone	[56–60]
11α	<i>Aspergillus</i> sp., <i>Rhizopus</i> sp.	Obtaining of anti-inflammatory, immunosuppressive, anti-allergic drugs, and production of contraceptive drugs	[56, 61]
14α	<i>Mucor</i> sp.		[56]
15β	<i>Bacillus</i> sp.		[56, 62]
16α	<i>Streptomyces</i> sp.		[63, 64]

Table 4.
Some examples of steroid hydroxylation reactions promoted by microorganisms and their applications.

microorganisms to reduce 17-keto- to 17β-hydroxysteroids was evidenced for a wide variety of substrates and microorganisms of different taxonomy: bacteria, fungi, and yeast [54, 56, 57, 66].

Fungi		Yeast	Bacteria
<i>Actinomucor elegans</i>	<i>Fusarium</i>	<i>Candida albicans</i>	<i>B. stearothermophilus</i>
<i>Agaricus silvaticus</i>	<i>culmorum</i>	<i>C. pelliculosa</i>	<i>Bacteroides fragilis</i>
<i>A. pantherina</i>	<i>F. oxysporum</i> var.	<i>C. pseudotropicalis</i>	<i>Brevibacterium sterolicum</i>
<i>A. spissa</i>	<i>cubense</i>	<i>C. robusta</i>	<i>Clostridium paraputrificum</i>
<i>Armillaria mellea</i>	<i>F. solani</i>	<i>C. tropicalis</i>	<i>Comamonas testosteroni</i>
<i>Corticium centrifugum</i>	<i>Mucor piriformis</i>	<i>C. utilis</i>	<i>Lactobacillus bulgaricus</i>
<i>Fusarium</i> spp.	<i>M. spinosus</i>	<i>Cryptococcus</i>	<i>Mycobacterium</i> spp.
<i>Gibberella saubineti</i>	<i>P. chrysogenum</i>	<i>albidus</i>	<i>B. stearothermophilus</i>
<i>Mucor</i> spp.	<i>P. crustosum</i>	<i>C. laurentii</i>	<i>Bacteroides fragilis</i>
<i>Penicillium</i> spp.	<i>P. blakesleeanus</i>	<i>C. tsukubaensis</i>	<i>Brevibacterium sterolicum</i>
<i>Aphanocladium album</i>	<i>R. stolonifer</i>	<i>Debaryomyces</i>	<i>Clostridium paraputrificum</i>
<i>Aspergillus chevalieri</i>	<i>Septomyxa affinis</i>	<i>hansenii</i>	<i>Comamonas testosteroni</i> (syn.
<i>A. flavus</i>	<i>T. piriforme</i>	<i>D. klockeri</i>	<i>Pseudomonas testosteroni</i>)
<i>A. oryzae</i>	<i>Trichoderma</i>	<i>D. nicotianae</i>	<i>Lactobacillus bulgaricus</i>
<i>A. tamarii</i>	<i>viride</i>	<i>D. subglobosus</i>	<i>Pediococcus cerevisiae</i>
<i>B. obtusa</i>	<i>Zygodessmus</i> sp.	<i>D. vini</i>	<i>Sarcina lutea</i>
<i>C. aphidicola</i>		<i>Hansenula</i>	<i>Staphylococcus aureus</i>
<i>Ceratocystis paradoxa</i>		<i>anomala</i>	<i>Streptomyces globisporus</i>
<i>C. lunatus</i>		<i>H. californica</i>	<i>S. sphaeroides</i>
<i>Colletotrichum musae</i>		<i>H. schnegii</i>	<i>S. viridochromogenes</i>
<i>C. radicola</i>		<i>H. suaveolens</i>	<i>S. hydrogenans</i>
<i>Exophiala jeanselmei</i> var.		<i>Kloeckera jensenii</i>	<i>S. lavendulae</i>
<i>lecaniicorni</i>		<i>Saccharomyces</i>	
		<i>carlsbergensis</i>	
		<i>S. cerevisiae</i>	
		<i>S. fragilis</i>	
		<i>S. lactis</i>	
		<i>S. oviformis</i>	
		<i>S. turbidans</i>	
		<i>S. validus</i>	
		<i>Pichia farinosa</i>	
		<i>P.</i>	
		<i>membranaefaciens</i>	
		<i>Torulopsis</i> spp.	
		<i>Hortaea werneckii</i>	
		<i>Phaeotheca</i>	
		<i>triangularis</i>	
		<i>P. herbarum</i>	
		<i>P. ostreatus</i>	
		<i>Rhodotorula</i>	
		<i>aurantiaca</i>	
		<i>R. mucilaginoso</i>	

Table 5.
Reduction of the C-17 carbonyl group of steroids by (17βHSDs) different microorganisms.

The oxidation of 17β-hydroxyl group was observed along with hydroxylation of steroids at C₅ (*Penicillium crustosum*, *P. chrysogenum*), C₆ (*Bacillus stearothermophilus*, *B. obtusa*, *P. blakesleeanus*), C₇ (α/β) (*A. coerulea*, *Botrytis cinerea*, *B. obtusa*, *P. blakesleeanus*, *Rhizopus stolonifer*), C₁₀ (*Absidia glauca*), C₁₁ (α/β) (*A. coerulea*, *B. obtusa*, *Cephalosporium aphidicola*, *R. stolonifer*), C₁₂ (*A. glauca*, *B. obtusa*), C₁₄ (*Bacillus* sp.), and C₁₅ (*A. glauca*, *Aspergillus fumigatus*, *B. obtusa*) [54, 56, 57, 67–69]. The biotransformation of 79 with different microorganisms is shown. Compound 79 is an endogenous weak androgen steroid hormone and intermediate in the biosynthesis of estrone and of testosterone from dehydroepiandrosterone (DHEA) [70]. DHEA is an endogenous steroid hormone. It functions as a metabolic intermediate in the biosynthesis of the androgen and estrogen sex steroids. Various microorganisms have had the ability to biotransform steroidal compounds such as AD (79) [54], DHEA (125)

Figure 11. The ability of different fungi to transform DHEA (**125**), testosterone, cortisolone (**126**) and prednisone (**127**). (a) Hydroxylation of 3 β -hydroxy-5-androsten-17-one (DHEA) by various microorganisms. (b) Reduction of C-17 and hydroxylation of testosterone by various microorganisms. (c) Hydroxylation of cortisolone (**123**) by various microorganisms. (d) Reduction and hydroxylation of prednisone (**126**) microorganisms.

[54, 55, 70–73, 76, 77, 80, 81], testosterone [54, 55, 74–76, 81], cortisolone (**126**) [78, 79], and prednisone (**127**) (**Figure 11a–d**) [54, 55, 82].

2.5 Diterpene

Sclareolide (**128**) is a natural product isolated from several plant species which displays phytotoxic and cytotoxic activities against several human tumor cells lines. This compound has also been used as starting material for the synthesis of various bioactive products. Regarding the biotransformation of the **128** with different microorganisms, mono- (**130**, **131**, **135**, **140–142**) and dihydroxylation (**132–134**, **136**, **139**, **143**, **146**), oxidation (**129**, **144**), hydroxylation/oxidation (**145**), epimerization (**137**), and cyclization (**138**) products have been obtained [83]. The microbial transformation of **128** by *Curvularia lunata* yielded 3-ketoesclareolide (**129**), 1 β -hydroxysclareolide (**130**), 3 β -hydroxysclareolide (**131**), 1 α ,3 β -dihydroxysclareolide (**133**), and 1 β ,3 β -dihydroxysclareolide (**134**) [84]. The incubation of **128** with *Cunninghamella elegans* afforded **129**, **131**, **133**, and **135–137** [85]. *C. blakesleeana* metabolized **128** to afford **129**, **135**, **134**, and **138–140**. Biotransformation of **128** with *C. echinulata* yielded 5-hydroxysclareolide (**141**) and 7 β -hydroxysclareolide (**142**) [86]. Fermentation of **148** with *A. niger* using a nutrient-rich culture medium yielded **141** and **144–146** (**Figure 12**) [83].

2.6 Flavonoids

As most important phytochemicals in food, the dietary flavonoids exert a wide range of benefits for human health. Recent researches have explored diverse biological and pharmacological activities of natural flavonoids—antioxidant activity, anti-inflammatory activity, anti-Alzheimer's disease, antibacterial activity, antifungal activity, anti-HIV activity, anticoagulant activity, antileishmanial activity, and

Figure 12.

Biotransformation products of sclareolide (128).

anti-obesity activity [87–91]. Microbial biotransformation strategies for production of flavonoids have attracted considerable interest because they allow yielding novel flavonoids, which do not exist in nature.

The main reactions during microbial biotransformation are hydroxylation, dehydroxylation, O-methylation, O-demethylation, glycosylation, deglycosylation, dehydrogenation, hydrogenation, C ring cleavage of the benzo- γ -pyrone system, cyclization, and carbonyl reduction. *Cunninghamella*, *Penicillium*, and *Aspergillus* strains are very popular to biotransform flavonoids, and they can perform almost all the reactions with excellent yields (Figure 13). Isoflavones are usually hydroxylated at the C-3' position of the B ring by microorganisms. Chalcones 147–152 were regioselectively cyclized to flavanones (Figure 14). Hydrogenation of flavonoids was only reported on transformation of chalcones to dihydrochalcones (Figure 14) [92, 93].

Aspergillus niger is one of the most applied microorganisms in the flavonoids' biotransformation; for example, *A. niger* can transfer flavanone to flavan-4-ol, 2'-hydroxydihydrochalcone, flavone, 3-hydroxyflavone, 6-hydroxyflavanone, and 4'-hydroxyflavanone. The hydroxylation of flavones by microbes usually happens on the ortho position of the hydroxyl group on the A ring and C-4' position of the B ring, and microbes commonly hydroxylate flavonols at the C-8 position. Natural flavonoids, such as naringenin (166), hesperetin (167), chrysin (168), apigenin (169), and luteolin (170) were subjected to microbiological transformations by *Rhodotorula glutinis* (KCh 735). Yeast was able to regioselectively C-8 hydroxylate 167, 168, 169, and 170 to generate 171 (17%), 172 (31%), 173 (12.9%), and 174 (25%), respectively. Naringenin (166) was transformed to carthamidin (175) and isocarthamidin (176) in a ratio of 1:19, respectively (Figure 15) [94].

Figure 13.
 The main reactions during biotransformation of chalcone with microorganisms.

The microorganisms tend to hydroxylate flavanones at the C-5, 6, and 4' positions; however, for prenylated flavanones, dihydroxylation often takes place on the $\Delta^{4(5)}$ double bond on the prenyl group (the side chain of A ring), although cyclization of the prenyl group to dihydrofuran derivatives is rather common biotransformation pathway of prenylated flavonoids. Prenylated flavanones are a unique class of naturally occurring flavonoids characterized by the presence of a prenylated side chain (prenyl, geranyl) in the flavonoid skeleton [95]. The prenyl chain generally refers to the 3,3-dimethylallyl substituent (3,3-DMA), geranyl and lavandulyl. It is proposed that the prenyl-moiety makes the backbone compound more lipophilic, which leads to its high affinity with cell membranes. The prenylation brings the flavonoids with enhancement of antibacterial, anti-inflammatory, antioxidant, cytotoxicity, larvicidal, as well as estrogenic activities. **Figure 16** demonstrated

Figure 14.
 Biotransformation products obtained from biotransformation of chalcones 147-152 with *A. niger*.

Figure 15.

Biotransformation products of flavanone (**166**, **167**) and flavone (**168**–**170**).

Figure 16.

Microbial biotransformation of kurarinone (**177**) using *C. echinulate* and *C. militaris*.

the microbial biotransformation of kurarinone (**177**) using *C. echinulata* and *C. militaris* [96, 97].

Incubation of *Absidia coerulea* (AM93) with prenylnaringenin (**178**) led to metabolite **179** (8-prenylnaringenin 7-O- β -D-glucopyranoside, 49.3%), while *B. bassiana* transformed **178** into **180** (8-prenylnaringenin 7-O- β -D-4''-O-methyl-glucopyranoside, 32.9%); the metabolites **179** and **180** originated in Sabouraud medium. In the absence of glucose in the culture of *A. coerulea*, the sulfation of substrate **178** (8-prenylnaringenin-7-sulfate, **181**, 31.1%) occurs, while *B. bassiana* into the same product (**180**). The capacity of some fungi—*Cunninghamella elegans*, *Streptomyces fulvissimus*, *Mucor ramannianus*, and *B. bassiana*—in the sulfation of certain phenolic compounds has been reported (**Figure 17**) [98].

Regioselective glycosylation of biologically active flavonoid aglycones catalyzed by microorganisms is an interesting and desired reaction, which significantly increases the water solubility of the compound and, therefore, may improve bio-availability of flavonoids. *Absidia glauca* AM177, *A. coerulea* AM93, *Rhizopus nigricans* UPF701, *Beauveria bassiana* AM278, and *B. bassiana* AM446 are able to conjugate

Figure 17.
Biotransformation products of prenylnaringenin (178).

Figure 18.
Biotransformation products of bavachinin (182).

sugar moiety to chalcones, flavanones, and isoflavanones with high regioselectivity. Therefore, it is possible to use *Beauveria* and *Absidia* for the microbial transformation of simple or prenylated flavonoids by glycosidation reactions [97, 99].

Bavachinin (182) is one kind compound of flavanones and isolated from the aerial parts and dried fruits of *Psoralea corylifolia*, and bavachinin displays a broad range of biological activities, such as antioxidant, antibacterial, antifungal, anti-inflammatory, antitumor, anti-pyretic, and analgesic properties [100, 101]. Bavachinin (182) was subject to biotransformation by cultured cells of *A. flavus* (ATCC 30899); *C. elegans* (CICC 40250) afforded the same product 183 [(S)-6-((R)-2,3-dihydroxy-3-methylbutyl)-2-(4-hydroxyphenyl)-7-methoxychromen-4-one]. On the other hand, one major product 184 [(2S,4R)-2-(4-hydroxyphenyl)-7-methoxy-6-(3-methylbut-2-en-1-yl)-chromen-4-ol] was obtained by *P. raistrickii* (ATCC 10490) by the reduction at the position of ketone group of the C-ring (Figure 18) [102].

The biotransformation of xanthohumol (185), a prenylated chalcone isolated from hops by selected fungi, *Absidia coerulea* (AM93), *Rhizopus nigricans* (UPF701), *Mortierella mutabilis* (AM404), and *Beauveria bassiana* (AM446), was investigated. The incubation of *A. coerulea* with 185 resulted in the isolation of xanthohumol 4'-O-β-D-glucopyranoside (186, 29%). This metabolite was also produced by *R. nigricans* (186, 14.2%). Biotransformation of 185 with *B. bassiana* and *M. mutabilis* yielded xanthohumol 7-O-β-D(4'''-O-methyl)-glucopyranoside (187, 23%) and isoxanthohumol 7-O-β-glucopyranoside (188, 49%), respectively (Figure 19) [103]. The compounds 188 (9.3%) and 186 (12%) were also observed as products of 185 transformation by *Cunninghamella elegans* [104]. Another way to obtain 188 is by the transformation of isoxanthohumol (189, 61.6%) with *Absidia glauca*; although

Figure 19.
Biotransformation products of xanthohumol (185).

the efficiency of this process was high (61.6% yield), it required the chemical isomerization of **185** to **189**, prior to biotransformation [105].

2''-(2''-hydroxyisopropyl)-dihydrofurano-[4'',5'':3',4']-4,2'-dihydroxy-6'-methoxychalcone (**190**), mixture of diastereoisomers of (2S, 2''S) and (2S, 2''R) 2''-(2''-hydroxyisopropyl)-dihydrofurano-[4'',5'':7,8]-4'-hydroxy-5-methoxyflavanone (**191**), and (Z)-2''-(2''-hydroxyisopropyl)-dihydrofurano-[4'',5''-6,7]-3',4'-dihydroxy-4-methoxyaurone (**192**) were obtained by transformation of **185** in *Aspergillus ochraceus* (AM 465) culture (**Figure 19**) [106].

Incubation of xanthohumol (**185**) both with *Fusarium avenaceum* (AM11) and *F. oxysporum* (AM727) gave a single metabolite 2''-(2''-hydroxyisopropyl)-dihydrofurano-[4'',5'':3',4']-4',2-dihydroxy-6'-methoxy- α,β -dihydrochalcone (**193**), which turned out to be the product of the prenyl group cyclization and α,β -double bond reduction. *F. tricinatum* reduced α,β -double bond of **185** to give 4,2',4'-trihydroxy-6'-methoxy-3-prenyl- α,β -dihydrochalcone (**194**). *Penicillium albidum* (AM79) oxidized **185** at the double bond of prenyl group to xanthohumol H (**195**) [107]. The culture of the yeast, *Rhodotorula marina* (AM 77), converted **185** and 4-methoxychalcone (**196**) to α,β -dihydroxanthohumol (**197**) and 4-methoxydihydrochalcone (**198**) with the yields of 18% and 20%, respectively [108]. *Penicillium albidum* (AM79) dihydroxylated the $\Delta^{2''(3'')}$ double bond of xanthohumol to produce 3'-[3''-hydroxy-3''-methylbutyl]-4,2',4'-trihydroxy-6'-methoxychalcone (**199**, 22.84%) (**Figure 19**).

B. bassiana AM278 and *Absidia glauca* AM177 converted isoxanthohumol (**189**) into glucoside derivatives (**200**, **201**), whereas *Fusarium equiseti* AM15 transformed it into (2R)-2-(2-hydroxyisopropyl)-dihydrofurano-[2,3:7,8]-4-hydroxy-5-methoxyflavanone (**202**) (**Figure 20**) [95, 106].

C. echinulata (ATCC 9244) sulfated silybin (**203**) to silybin-7-sulfate (**204**) and 2,3-dehydrosilybin-7-sulfate (**205**). Sulfonation at the C-7 position of silybin

Figure 20.
 Biotransformation products of isoxanthohumol (189).

significantly decreased the DPPH free radical scavenging potential; however, further dehydrogenation $\Delta^{2(3)}$ to 2,3-dehydrosilbyn-7-sulfate (206) drastically enhanced the DPPH free radical scavenging potential activity [109] (Figure 21).

2.7 Enzymes isolated from microorganisms and their application

Enzymes are the most proficient catalysts, offering much more competitive processes than chemical catalysts. A number of enzyme-based processes have been commercialized for producing several valuable products. During the 1980s and 1990s, engineering of enzymes based on structural information allowed extension of their substrate ranges, enabling the synthesis of unusual intermediates. Accordingly, the use of enzymes has been expanded to the manufacture of pharmaceutical intermediates and fine chemicals [110]. Microorganisms and enzymes (biocatalysts) are highly enantio-, chemo-, and regioselective in a wide range of reaction conditions. Selectivity is extremely desirable in the synthesis of different synthesis products, since it offers advantages such as minimizing the side reactions that do not require protection and deprotection steps, which allows for shorter synthesis. Biocatalysis provides a technology that is environmentally safer, and it effectively reduces the level of waste and even eliminates the waste generation rather than remediation and disposal of wastes at the end of the process. In addition

Figure 21.
 Biotransformation products of sylbin (204).

to, biocatalysts have many attractive features in the context of green chemistry and sustainable development. Various enzymes used in different industrial processes have been described in the literature. **Table 6** indicates some enzymes, their source, and some applications [111–113].

2.8 Extremophiles

A very interesting research area in biology and biotechnology is the of extremophile microorganisms. Extremophiles can be divided into group according to (i) temperature tolerance, (ii) salt concentration, (iii) pH range, or (iv) pressure conditions. Enzymes from extremophilic microorganisms offer versatile tools for

Microbial enzymes	Microorganism	Application
α -Amylase	<i>Bacillus</i>	Baking, brewing, starch liquefaction
	<i>amyloliquefaciens</i>	Clarification of fruit juice
	<i>B. stearothermophilus</i>	Textile industry
	<i>B. licheniformis</i>	Paper industry
Glucoamylase	<i>Aspergillus niger</i>	Beer production
	<i>A. awamori</i>	High glucose and high fructose syrups
	<i>Rhizopus oryzae</i>	
Proteases	<i>A. usami</i>	
Lactase (β -galactosidase)	<i>Kluyveromyces lactis</i>	Lactose intolerance reduction in people
	<i>K. fragilis</i>	Prebiotic food ingredients
Lipase	<i>Candida antarctica</i>	Cheese flavor development
	<i>C. cylindraceae</i> Ay30	Textile indutry
	<i>Helvina lanuginosa</i>	Medicinal applications
	<i>Pseudomonas</i> sp.	Use in cosmetics
	<i>Geotrichum candidum</i>	Use as biosensors Use in biodegradation
Phospholipases	<i>Fusarium oxysporum</i>	Cheese flavor development
Esterases	<i>Bacillus licheniformis</i>	Enhancement of flavor and fragrance in fruit juice
Xylanases	<i>Streptomyces</i> sp.	Clarification of fruit juice
	<i>Bacillus</i> sp.	Beer quality improvement
	<i>Pseudomonas</i> sp.	
Glucose oxidase	<i>A. niger</i>	Food shelf life important
	<i>Penicillium glaucum</i>	Food flavor improvement
	<i>P. adametzzii</i>	
Laccase	<i>Funalia trogii</i>	Polyphenol removal from wine
	<i>Bacillus licheniformis</i>	baking
	<i>Bacillus vallismortis</i>	
Pectinases	<i>A. niger</i>	Clarification of fruit juice
	<i>A. wentii</i>	
	<i>Rhizopus</i> sp.	
Catalase	<i>A. niger</i>	Food preservation
	<i>Metarhizium</i>	Removal of H ₂ O ₂ from milk prior to cheese
	<i>anisopliae</i>	production
	<i>Psychrobacter piscatorri</i>	
Peroxidase	<i>Streptomyces viridosporus</i>	Development of flavor, color and nutritional quality of food

Table 6.
Enzymes, source, and some applications.

sustainable developments in a variety of industrial applications as they show important environmental benefits due to their biodegradability, specific stability under extreme conditions, improved use of raw materials, and decreased amount of waste products. Although major advances have been made in the last decade, our knowledge of the physiology, metabolism, enzymology, and genetics of this fascinating group of extremophilic microorganisms and their related enzymes is still limited [114–116].

The outstanding properties of thermozymes are suited to industries that employ elevated temperatures, such as the pulp and paper, food, brewing, and feed processing industries. Thermophiles are often highly resistant to harsh conditions such as chemical denaturing agents, wide pH ranges, and/or nonaqueous solvents. Examples of such enzymes are cellulases, xylanases, pectinases, chitinases, amylases, pullulanases, proteases, lipases, glucose isomerases, alcohol dehydrogenases, and esterases. Thermophilic enzymes have played important roles not only at the industrial level but also in pharmaceutical applications requiring use of specific aldolases for the synthesis of enantiopure compounds (Table 7) [118].

Source	Enzyme	Activity	Bioprocess/industry	Reference
<i>Sulfolobus solfataricus</i> <i>S. acidocaldarius</i> <i>Thermoproteus texas</i> <i>Hyperthermus butylicus</i>	Aldolase	Stereoselective C-C bond formation	Pharmaceutical industry	[117]
<i>Pyrococcus furiosus</i>	Hydrogenase	Final stage of glucose oxidation by oxidative pentose phosphate cycle	Enhanced production of biohydrogen	[119]
<i>Geobacillus thermoleovorans</i>	Carboxylesterase	Carboxyl ester hydrolysis	Agriculture, food, and pharmaceutical industries	[120]
<i>Bacillus pumilus</i>	Acidic thermostable lipase	Degradation of palm oil	Treatment of palm oil-containing wastewater	[121]
<i>Geobacillus</i> sp.	Lipase	Hydrolysis of diver's lipid substrates	Biofuel, cosmetics, or perfume production, leather and pulp industries	[122]
Microbial community from solid-state fermentation reactor	Protease	Degradation of hair waste from tannery	Leather industry	[123]
<i>Sulfolobus tokodaii</i>	Chitinase	Hydrolysis of β -(1, 4)-glycosidic bonds in chitin	Biomedical, pharmaceutical, food, and environmental	[124]
<i>Acidothermus cellulolyticus</i>	Endoxylanase	β -(1,4)-xylan cleavage	Biofuel production from lignocellulose	[125]
<i>Thermotoga neapolitana</i>	Pullulanase	Hydrolysis of α -(1, 6)-glucosidic linkages	Biofuel production	[126]

Table 7.
Extremophile microorganisms and some applications of their enzymes.

3. Conclusion

Due to microorganisms' abundant multienzyme systems, microbial transformation possesses advantages against chemosynthesis of environmental friendliness, mild reaction conditions, and high *stereo-*, *regio*-, and *chemo*-selectivities as well as in improving conversion rates and reducing cost. Thus, microbial transformation technique is being increasingly used to structurally modify natural and synthetic compounds.

The hydrolytic and reductive capabilities of microorganisms have been known and are currently used in preparative and industrial reactions. Various classes of bioactive organic compounds have been subjected to enzymatic transformation to obtain more active and less toxic substances or to elucidate their metabolic pathways.

For example, biotransformation-derived steroids are used for a wide range of pharmacotherapeutic purposes, such as anti-inflammatory, immunosuppressive, progestational, diuretic, anabolic, as neurosteroids, and as contraceptive. Researchers continue to discover more useful steroid compounds and to isolate microorganisms that can perform the structural transformations desired. New technologies such as genomics, metagenomics, gene shuffling, and DNA evolution provide valuable tools for improving or adapting enzyme properties to the desired requirements.

An alternative may be extremophilic microorganisms such as biocatalysts for countless future industrial applications that are more environmentally friendly.

Acknowledgements

The authors thank Carrera de Biología, FES-Zaragoza, UNAM, Al Departamento de Química Orgánica, FES-Cuautitlán, UNAM.

Conflict of interest

The authors report no conflicts of interest.

IntechOpen

Author details

Arturo Cano-Flores^{1*}, Javier Gómez¹, Iker S. Escalona-Torres¹
and Benjamín Velasco-Bejarano²

¹ Laboratorio de Biotransformaciones y Química de Productos Naturales, L-314,
Facultad de Estudios Superiores Zaragoza, UNAM, CDMX, Mexico

² Sección de Química Orgánica, Departamento de Ciencias Químicas,
FES-Cuautitlán, UNAM, Edo. de México, Mexico

*Address all correspondence to: aecanomx@unam.mx; acano1750@gmail.com

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Lehman LR, Stewart JD. Filamentous fungi: Potentially useful for the biohydroxylation of non-activated carbón centres. *Current Organic Chemistry*. 2001;5:439-470. DOI: 10.2174/1385272013375490
- [2] Pollard DJ, Woodley JM. Biocatalysis for pharmaceutical intermediates: The future is now. *Trends in Biotechnology*. 2006;25:66-72. DOI: 10.1016/j.tibtech.2006.12.005
- [3] Straathof AJJ. The production of fine chemicals by biotransformation. *Current Opinion in Biotechnology*. 2002;13:548-556. DOI: 10.1016/S0958-1669(02)00360-9
- [4] Parra A, Rivas F, Garcia-Granados A, Martínez A. Microbial transformation of triterpenoids. *Mini-Reviews in Organic Chemistry*. 2009;6:307-320. DOI: 10.2174/157019309789371569
- [5] Schrewe M, Julsing MK, Buhler B, Schmid A. Whole-cell biocatalysis for selective and productive C-O functional group introduction and modification. *Chemical Society Reviews*. 2013;42:6346. DOI: 10.1039/c3cs60011d
- [6] Brenna E, Fugati C, Gatti FG, Serra S. Biocatalytic methods for the synthesis of enantioenriched odor active compounds. *Chemical Reviews*. 2011;111:4036-4072. DOI: 10.1021/cr100289r
- [7] Sultana N, Saeed SZ. Enzymatic biotransformation of terpenes as bioactive agents. *Journal of Enzyme Inhibition and Medicinal Chemistry*. 2013;28:113-1128. DOI: 0.3109/14756366.2012.727411
- [8] Silva EO, Gonçalves NS, Santos RA, Furtado NAJC. Microbial metabolism of atovaquone and cytotoxicity of the produced phase I metabolite. *European Journal of Drug Metabolism and Pharmacokinetics*. 2016;41:645-650
- [9] Pervaiz I, Ahmad S, Madni MA, Ahmad H, Khaliq FH. Microbial biotransformation: A tool for drug designing. *Applied Biochemistry and Microbiology*. 2013;49:437-450. DOI: 10.1134/S0003683813050098
- [10] Zhan Y, Liu H, Wu Y, Wei P, Chen Z, William JS. Biotransformation of artemisinin by *Aspergillus niger*. *Applied Microbiology and Biotechnology*. 2015;99:3443-3446. DOI: 10.1007/s00253-015-6464-x
- [11] Ponnappalli MG, Sura MB, Sudhakar R, Govindarajalu G, Sijwali PS. Biotransformation of Artemisinin to 14-hydroxydeoxyartemisinin: C-14 hydroxylation by *Aspergillus flavus*. *Journal of Agricultural and Food Chemistry*. 2018;66:10490-10495. DOI: 10.1021/acs.jafc.8b03573
- [12] Srivastava S, Luqman S, Fatima A, Dorokar MP, Negi AS, Kumar JK, et al. Biotransformation of artemisinin mediated through fungal strains for obtaining derivatives with novel activities. *Scientia Pharmaceutica*. 2009;77:87-95. DOI: 10.3797/scipharm.0803-15
- [13] Parhikov IA, Muraludharan KM, Avery MA, Williamson JS. Transformation of artemisinin by *Cunninghamella elegans*. *Applied Microbiology and Biotechnology*. 2004;64:782-786. DOI: 10.1007/s00253-003-1524-z
- [14] Zhan Y, Wu Y, Xu F, Bai Y, Guan Y, Williamson JS, et al. A novel dihydroxylated derivative of artemisinin from microbial transformation. *Fitoterapia*. 2017;120:93-97. DOI: 10.1016/j.fitote.2017.05.015
- [15] Goswami A, Saikia PP, Barua NC, Bordoloi M, Yadav A, Bora TC, et al.

- Bio-transformation of artemisinin using soil microbe: Direct C-acetoxylation of artemisinin at C-9 by *Penicillium simplicissimum*. Bioorganic & Medicinal Chemistry Letters. 2010;**20**:359-361. DOI: 10.1016/j.bmcl.2009.10.097
- [16] Gaur R, Darokar MP, Ajayakumar PV, Shukla RS, Bhakuni RS. In vitro antimalarial studies of novel artemisinin biotransformed products and its derivatives. Phytochemistry. 2014;**107**:135-140. DOI: 10.1016/j.phytochem.2014.08.004
- [17] Liu J-H, Chen Y-G, Yu B-Y, Chen Y-J. A novel ketone derivative of artemisinin biotransformed by *Streptomyces griseus* ATCC 13273. Bioorganic & Medicinal Chemistry Letters. 2006;**16**:1909-1912. DOI: 10.1016/j.bmcl.2005.12.076
- [18] Saraswati S, Agrawal S-S, Alhaider A-A. Ursolic acid inhibits tumor angiogenesis and induces apoptosis through mitochondrial-dependent pathway in Ehrlich ascites carcinoma tumor. Chemico-Biological Interactions. 2013;**206**:153-165. DOI: 10.1016/j.phytol.2019.04.019
- [19] Zhang C, Xu S-H, Ma B-L, Wang W, Yu B-Y, Zhang J. New derivatives of ursolic acid through the biotransformation by *Bacillus megaterium* CGMCC 1.1741 as inhibitors on nitric oxide production. Bioorganic & Medicinal Chemistry Letters. 2017;**27**:2575-2578. DOI: 10.1016/j.bmcl.2017.03.076
- [20] Zhang C-X, Ma W-J, Liu D-L, Jia X-J, Zhao Y-M. Biotransformation of ursolic acid by *Alternaria longipes* AS3.2875. Natural Product Research. 2018;**32**:536-543. DOI: 10.1080/14786419.2017.1327860
- [21] Wang Y, Xiang L, Chen M, Zhang Z-X, He X. Substrate specify for the 2 β -hydroxylation of ursolic acid by *Alternaria alternate* and the antitumor activities of those metabolites. Journal of Molecular Catalysis B: Enzymatic. 2012;**83**:51-56. DOI: 10.1016/j.molcatb.2012.07.005
- [22] S-b F, Yang J-S, J-l C, Sun D-A. Biotransformation of ursolic acid by *Syncephalastrum racemosum* CGMCC 3.2500 and anti-HCV activity. Fitoterapia. 2013;**86**:123-128. DOI: 10.1016/j.fitote.2013.02.007
- [23] Huang F-X, Yang W-Z, Ye F, Tian J-Y, Hu H-B, Feng L-M, et al. Microbial transformation of ursolic acid by *Syncephalastrum racemosum* (Cohn). Phytochemistry. 2012;**82**:56-60. DOI: 10.1016/j.phytochem.2012.06.020
- [24] Fu S-B, Yang J-S, Cui J-L, Meng Q-F, Feng X, Sun D-A. Multihydroxylation of ursolic acid by *Pestalotiopsis microspora* isolated from the medicinal plant *Huperzia serrate*. Fitoterapia. 2011;**82**:1057-1061. DOI: 10.1016/j.fitote.2011.06.09
- [25] Zhanga S-S, Hea L-S, Zhaoa Y-M, Fub S, Liua D-L, Yua Z-H, et al. Three new triterpenoids transformed from ursolic acid by *Mucor spinosus* (AS 3.3450) and their cytotoxicity. Phytochemistry Letters. 2019;**32**:33-37. DOI: 10.1016/j.phytol.2019.04.019
- [26] Wang Y, Sun Y, Wang C, Huo X, Liu P, Wang C, et al. Biotransformation of 11-keto-b-boswellic acid by *Cunninghamella blakesleeana*. Phytochemistry. 2013;**96**:330-336. DOI: 10.1016/j.phytochem.2013.07.018
- [27] Yu-Juan Q, Bing F, Xin-Bo S, Wen-Bin Z, He-Shui Y, Li-Li Z, et al. Biotransformation of glycyrrhetic acid by *Cunninghamella blakesleeana*. Chinese Journal of Natural Medicines. 2010;**8**:373-381. DOI: 10.3724/SP.J.1009.2010.00373
- [28] Maatooq GT, Marzouk AM, Gray AI, Rosazza JP. Bioactive microbial metabolites from glycyrrhetic acid.

- Phytochemistry. 2010;**71**:262-270. DOI: 10.1016/j.phytochem.2009.09.014
- [29] Mufler K, Leipold D, Scheller M-C, Haas C, Steingroewer J, Bley T, et al. Biotransformation of triterpenes. *Process Biochemistry*. 2011;**46**:1-15. DOI: 10.1016/j.procbio.2010.07.015
- [30] Chi H, Ji G. Transformation of Ginsenosides Rb1 and Re from *Panaxginseng* by food microorganisms. *Biotechnology Letters*. 2005;**27**:765-771. DOI: 10.1007/s10529-005-5632-y
- [31] Han Y, Sun B, Jiang B, Hu X, Spranger MI, Zhang Y, et al. Microbial transformation of ginsenosides Rb1, Rb3 and Rc by *Fusarium sacchari*. *Journal of Applied Microbiology*. 2010;**109**:792-798. DOI: 10.1111/j.1365-2672.2010.04707.x
- [32] Gao J, Xu W, Fang Q, Liang F, Jin R, Wu D, et al. Efficient biotransformation for preparation of pharmaceutically active ginsenoside compound K by *Penicillium oxalicum* sp. *Annales de Microbiologie*. 2013;**63**:139-149. DOI: 10.1007/s13213-012-0454-3
- [33] Dong A, Ye M, Guo H, Zhen J, Guo D. Microbial transformation of ginsenoside Rb1 by *Rhizopus stolonifer* and *Curvularia lunata*. *Biotechnology Letters*. 2003;**25**:339-344
- [34] Shirane N, Hashimoto Y, Ueda K, Takbnaka H, Katoh K. Ring a cleavage 3-oxo-olean-12-en-28-oic acid by the fungus *Chaetomium longirostre*. *Phytochemistry*. 1996;**43**:99-104
- [35] Martinez A, Perojil A, Rivas F, Parra A, Garcia-Granados A, Fernandez-Vivas A. Biotransformation of oleanolic and maslinic methyl esters by *Rhizomucor miehei* CECT 2749. *Phytochemistry*. 2015;**117**:500-508. DOI: 10.1016/j.phytochem.2015.07.020
- [36] Kouzi SA, Chatterjee P, Pezzuto JM, Hamann MT. Microbial transformations of the antimelanoma agent betulinic acid. *Journal of Natural Products*. 2000;**63**:1653-1657
- [37] Chatterjee P, Kouzi SA, Pezzuto JM, Hamann MT. Biotransformation of the antimelanoma agent betulinic acid by *Bacillus megaterium* ATCC 13368. *Applied and Environmental Microbiology*. 2000;**66**:3850-3855
- [38] Chatterjee P, Pezzuto JM, Kouzi SA. Glucosidation of betulinic acid by *Cunninghamella* species. *Journal of Natural Products*. 1999;**62**:761-763
- [39] Akihisa T, Takamine Y, Yoshizumi K, Tokuda H, Kimura Y, Ukiya M, et al. Microbial transformations of two lupane-type triterpenes and anti-tumor promoting effects of the transformation products. *Journal of Natural Products*. 2002;**65**:278-282
- [40] Goswami A, Guo Z, Tullya TP, Rinaldia FA, Huangb XS, Swidorskib JJ, et al. Microbial transformations of betulinic and betulonic acids. *Journal of Molecular Catalysis B: Enzymatic*. 2015;**117**:45-53. DOI: 10.1016/j.molcatb.2015.04.012
- [41] Wang W-W, Xu S-H, Zhao Y-Z, Zhang C, Zhang Y-Y, Yu B-Y, et al. Microbial hydroxylation and glycosylation of pentacyclic triterpenes as inhibitors on tissue factor procoagulant activity. *Bioorganic & Medicinal Chemistry Letters*. 2017;**27**:1026-1030. DOI: 10.1016/j.bmcl.2016.12.066
- [42] Yana S, Lina H, Huang H, Yanga M, Xua B, Chena G. Microbial hydroxylation and glycosidation of oleanolic acid by *Circinella muscae* and their anti-inflammatory activities. *Natural Product Research*. 2019;**33**:1849-1855. DOI: 10.1080/14786419.2018.1477150

- [43] Akihisa T, Watanabe K, Yoneima R, Suzuki T, Kimura Y. Biotransformation of cycloartane-type triterpenes by the fungus *Glomerella fusarioides*. Journal of Natural Products. 2006;**69**:604-607
- [44] Wang F-Q, Li B, Wang W, Zhang C-G, Wei D-Z. Biotransformation of diosgenin to nuatigenin-type steroid by a newly isolated strain, *Streptomyces virginiae* IBL-14. Applied Microbiology and Biotechnology. 2007;**77**:771-777. DOI: 10.1007/s00253-007-1216-1
- [45] Adham NZ, Zaki RA, NAim N. Microbial transformation of diosgenin and its precursor furestanol glycosides. World Journal of Microbiology and Biotechnology. 2009;**25**:481-487. DOI: 10.1007/s11274-008-9913-1
- [46] Wu G-W, Gao J-M, Shi X-W, Hang Q, Wei S-P, Ding K. Microbial transformations of diosgenin by the white-rot Basidiomycete *Coriolus versicolor*. Journal of Natural Products. 2011;**14**:2095-2101. DOI: 10.1021/np2003484
- [47] Xu M, Huo X-K, Tian X-G, Dong P-P, Wang C, Huang S-S, et al. Microbial transformation of diosgenin by *Cunninghamella blakesleana* AS 3.970 and potential inhibitory effects on P-glycoprotein of its metabolites. RSC Advances. 2015;**78**(78):081-78089. DOI: 10.1039/c5ra122553h
- [48] Zhao Y, Sun LM, Wang XN, Shen T, Ji M, Li X, et al. Microbial transformation of diosgenin by *Syncephalastrum racemosum* (Cohn) Schroeter. Chinese Chemical Letters. 2010;**21**:76-80. DOI: 10.1016/j.ccllet.2009.08.015
- [49] Dong T, Wua G-W, Wang X-N, Gao J-M, Chen J-G, Lee S-S. Microbiological transformation of diosgenin by resting cells of filamentous fungus, *Cunninghamella echinulata* CGMCC3.2716. Journal of Molecular Catalysis B: Enzymatic. 2010;**67**: 251-266. DOI: 10.1016/j.molcatb.2010.09.001
- [50] Peterson DH, Murray HC. Microbiological oxygenation of steroids at carbon-11. Journal of the American Chemical Society. 1952;**74**:1871-1872
- [51] Habibi Z, Yousefi M, Ghanian S, Mohammadi M, Ghasemi S. Biotransformation of progesterone by *Absidia griseolla* var. *igachii* and *Rhizomucor pusillus*. Steroids. 2012;**77**: 1446-1449. DOI: 10.1016/j.steroids.2012.08.010
- [52] Shan L, Jiao K, Yin M, Huang J, Chen Y, Qin S, et al. Biotransformation of 5-en-3 β -ol steroids by *Mucor circinelloides lusitanicus*. Biocatalysis and Biotransformation. 2016;**34**:83-88. DOI: 10.3109/10242422.2015.1089865
- [53] Choudhary MI, Erum S, Atif M, Malik R, Tameen K, Atta-ur-Rhaman. Biotransformation of (20S)-20-hydroxymethylpregna-1,4-dien-3-one by four filamentous fungi. Steroids. 2011;**76**:1288-1296. DOI: 10.1016/j.steroids.2011.06.007
- [54] Cano-Flores A, Gómez X, Ramos R. Biotransformation of steroids using different microorganisms. In: JAR S, editor. Chemistry and Biological Activity of Steroids. Rijeka: IntechOpen; 2018. DOI: 10.5772/intechopen.85849
- [55] Peart PC, McCook KP, Rusell FA, Reynolds WF, Reese PB. Hydroxylation of steroids by *Fusarium oxysporum*, *Exophiala jeanselmei* and *Ceratocystis paradoxa*. Steroids. 2011;**76**: 1317-1330. DOI: 10.1016/j.steroids.2011.06.010
- [56] Kristan K, Lanisnik Rizner T. Steroid-transforming in fungi. The Journal of Steroid Biochemistry and Molecular Biology. 2012;**129**:79-91. DOI: 10.1016/j.jsbmb.2011.08.012

- [57] Fernandes P, Cruz A, Angelova B, Pinheiro HM, Cabral JMS. Microbial conversion of steroid compounds: Recent developments. *Enzyme and Microbial Technology*. 2003;**32**:688-705. DOI: 10.1016/S0141-0229(03)00029-2
- [58] Suzuki K, Sanga K, Chikaoka Y, Itagaki E. Purification and properties of cytochrome P-450 (P-450_{1un}) catalyzing steroid 11 β -hydroxylation in *Curvularia lunata*. *Biochimica et Biophysica Acta*. 1993;**1203**:215-223. DOI: 10.1016/0167-4838(93)90086-7
- [59] Lu W, Du L, Wang M, Jia X, Wen J, Huang Y, et al. Optimisation of hydrocortisone production by *Curvularia lunata*. *Applied Biochemistry and Biotechnology*. 2007; **142**:17-28. DOI: 10.1007/s12010-007-0005-8
- [60] Bhosale S, Saratale G, Govindwar S. Biotransformation enzymes in *Cunninghamella blakesleeana* (NCIM-687). *Journal of Basic Microbiology*. 2006;**46**:444-448. DOI: 10.1002/jobm.200510117
- [61] Petrica S, Hakkib T, Bernhardt R, Zigonc D, Cresna B. Discovery of a steroid 11-hydroxylase from *Rhizopus oryzae* and its biotechnological application. *Journal of Biotechnology*. 2010;**150**:428-437. DOI: 10.1016/j.jbiotec.2010.09.928
- [62] Schmitz D, Janocha S, Kiss FM, Berhardt R. CYP106A2—A versatile biocatalyst with high potential for biotechnological production of selectively hydroxylated steroid and terpenoid compounds. *BBA Proteins and Proteomics*. 2018;**1866**:11-22. DOI: 10.1016/j.bbapap.2017.07.011
- [63] Berrie JR, Williams RAD, Smith KE. Microbial transformation of steroids—XI. Progesterone transformation by *Streptomyces roseochromogenes*—Purification and characterisation of the cytochrome P450 progesterone 16 α -hydroxylase system. *The Journal of Steroid Biochemistry and Molecular Biology*. 1999;**71**:153-165
- [64] Berrie JR, Williams RAD, Smith KE. Microbial transformation of steroids—XII. Progesterone hydroxylation profiles are modulated by post-translational modification of an electron transfer protein in *Streptomyces roseochromogenes*. *The Journal of Steroid Biochemistry and Molecular Biology*. 2001;**77**:87-96
- [65] Tang R, Shen Y, Wang M, Zhou H, Zhao Y. Highly efficient synthesis of boldenone from androst-4-ene-3,17-dione by *Arthrobacter simplex* and *Pichia pastoris*. *Process Biochemistry*. 2019;**42**: 933-940. DOI: 10.1007/s00449-019-02092-y
- [66] Hunter AC, Khuenl-B RH, Barret P, Dodd HT, Dedi C. Transformation of some 3-substituted steroids by *Aspergillus tamarii* KITA reveals stereochemical restriction of steroid binding orientation in the minor hydroxylation pathway A. *Journal of Steroid Biochemistry and Molecular Biology*. 2010;**118**:171-176. DOI: 10.1016/j.jsbmb.2009.12.003
- [67] Donova MV, Ergova OV. Microbial steroid transformations: Current state and prospect. *Applied Microbiology and Biotechnology*. 2012;**94**:1423-1447. DOI: 10.1007/978-1-4939-7183-1
- [68] Sultana N. Microbial biotransformation of bioactive and clinically useful steroids and some salient features of steroids and biotransformation. *Steroids*. 2018;**136**: 76-92. DOI: 10.1016/j.steroids.2018.01.007
- [69] Tong W-Y, Dong X. Microbial biotransformation: Recent developments on steroid drugs. *Recent Patents on Biotechnology*. 2009;**3**: 141-153

- [70] Lin Y, Song X, Fu J, Lin J, Qu Y. Microbial transformation of androst-4-ene-3,17-dione by *Bordetella* sp B4 CGMC2229. *Journal of Chemical Technology & Biotechnology*. 2009;**84**: 789-793. DOI: 10.1002/jctb.2113
- [71] Lobastova TG, Gulevskaya SA, Sukhodolskaya GV, Donova MV. Dihydroxylation of dehydroepiandrosterone in positions 7 α and 15 α by mycelial fungi. *Applied Biochemistry and Microbiology*. 2009; **45**:617-622
- [72] Li H, Liu H-M, Ge W, Huang L, Shan L. Synthesis of 7 α -hydroxy-dehydroepiandrosterone and 7 α -hydroxy-dehydroepiandrosterone. *Steroids*. 2005;**70**:970-973
- [73] Lobastova TG, Khomutov SM, Vasiljeva LL, Lapitskaya MA, Pivnitsky KK, Donova MV. Synthesis of 3 β -hydroxy-androsta-5,7-dien-17-one from 3 β -hydroxyandrost-5-en-17-one via microbial 7 α -hydroxylation. *Steroid*. 2009;**74**:233-237. DOI: 10.1016/j.steroids.2008.10.019
- [74] Al-Awadi S, Afzal M, Oommen S. Studies of *Bacillus stearothermophilus*. Part III. Transformation of testosterone. *Applied Microbiology and Biotechnology*. 2003;**62**:48-52. DOI: 10.1007/s00253-003-1269-8
- [75] Al-Awadi S, Afzal M, Oommen S. Studies on *Bacillus stearothermophilus*. Part IV. Influence of enhancers on biotransformation of testosterone. *Steroids*. 2005;**70**:323-327
- [76] Janeczko T, Dmochowska-Gladysz J, Kostrezewa-Suslowa E, Bialonska A, Ciunik Z. Biotransformations of steroid compounds by *Chaetomium* sp. KCH 6651. *Steroids*. 2009;**74**:657-660. DOI: 10.1016/j.steroids.2009.02.006
- [77] Romano A, Romano D, Ragg E, Costantinoc F, Lennac R, Gandolfid R, et al. Steroid hydroxylations with *Botryodiplodia malorum* and *Colletotrichum lini*. *Steroid*. 2006;**71**: 429-434. DOI: 10.1016/j.steroids.2006.01.014
- [78] Lisowska K, Dlugonski J. Concurrent corticosteroid and phenanthrene transformation by filamentous fungus *Cunninghamella elegans*. *The Journal of Steroid Biochemistry and Molecular Biology*. 2003;**85**:63-69
- [79] Manosroi J, Chisti Y, Manosroi A. Biotransformation of cortexolone to hydrocortisone by molds using a rapid color-development assay. *Applied Biochemistry and Microbiology*. 2006; **42**:479-483
- [80] Wu Y, Li H, Zhang X-M, Gong JS, Rao Z-M, Shi J-S, et al. Efficient hydroxylation of functionalized steroids by *Colletotrichum lini* ST-1. *Journal of Molecular Catalysis B: Enzymatic*. 2015; **120**:111-118. DOI: 10.1016/j.molcatb.2015.07.003
- [81] Agnieszka Bartmanska A, Dmochowska-Gladysz J. Transformation of steroids by *Trichoderma hamatum*. *Enzyme and Microbial Technology*. 2007;**40**: 1615-1621. DOI: 10.1016/j.enzmictec.2006.11.011
- [82] Choudhary MI, Siddiqui ZA, Musharraf SG, Nawaz SA, Atta-ur-Rahman. Microbial transformation of prednisone. *Natural Product Research*. 2005;**19**: 311-317. DOI: 10.1080/14786410410001729168
- [83] Cano A, Ramírez-Apan MT, Delgado G. Biotransformation of sclareolide by filamentous fungi: Cytotoxic evaluations of the derivatives. *Journal of the Brazilian Chemical Society*. 2011;**22**:1177-1182. DOI: 10.1590/S0103-50532011000600025

- [84] Atta-ur-Rahman, Farooq A, Choudhary MI. Microbial transformation of sclareolide. *Journal of Natural Products*. 1997;**60**:1038-1040. DOI: 10.1021/np970076h
- [85] Choudhary MI, Musharraf SG, Sami A, Atta-ur-Rahman. Microbial transformation of sesquiterpenes, (–)-ambrox and (+)-sclareolide. *Helvetica Chimica Acta*. 2004;**87**:2685-2694
- [86] Ata A, Conci LJ, Betteridge J, Orhan I, Sener B. Novel microbial transformations of sclareolide. *Chemical and Pharmaceutical Bulletin*. 2007;**55**: 118-123. DOI: 10.1248/cpb.55.118
- [87] Cao H, Chen X, Jassbi AR, Xiao J. Microbial biotransformation of bioactive flavonoids. *Biotechnology Advances*. 2015;**33**:214-233. DOI: 10.1016/j.biotechadv.2014.10.012
- [88] Andrae-Marobela K, Ghislain FW, Okatch H, Majinda R. Polyphenols: A diverse class of multi-target anti-HIV-1 agents. *Current Drug Metabolism*. 2013; **7**:392-413. DOI: 10.2174/1389200211314990095
- [89] Liao XL, Luo JG, Kong LY. Flavonoids from *Millettia nitida* var. *Hirsutissima* with their anticoagulative activities and inhibitory effects on NO production. *Journal of Natural Medicines*. 2013;**67**:856-861. DOI: 10.1007/s11418-013-0745-4
- [90] Ogungbe IV, Erwin WR, Setzer WN. Antileishmanial phytochemical phenolics: Molecular docking to potential protein targets. *Journal of Molecular Graphics & Modelling*. 2014; **48**:105-117. DOI: 10.1016/j.jmgm.2013.12.010
- [91] Kim H, Bartley GE, Arvik T, Lipson R, Nah SY, Seo K, et al. Dietary supplementation of chardonnay grape seed flour reduces plasma cholesterol concentration, hepatic steatosis, and abdominal fat content in high-fat diet-induced obese hamsters. *Journal of Agricultural and Food Chemistry*. 2014; **62**:1919-1925. DOI: 10.1021/jf404832s
- [92] Asha S, Vidyavathi M. *Cunninghamella*—A microbial model for drug metabolism studies—A review. *Biotechnology Advances*. 2009;**27**:16-29. DOI: 10.1016/j.biotechadv.2008.07.005
- [93] Alarcón J, Alderete J, Escobar C, Araya R, Cespedes CL. *Aspergillus niger* catalyzes the synthesis of flavonoids from chalcones. *Biocatalysis and Biotransformation*. 2013;**31**:160-167. DOI: 10.3109/10242422.2013.813489
- [94] Sordon S, Madej A, Poplonski J, Bartmanska A, Tronina T, Brzezowska E, et al. Regioselective ortho-hydroxylations of flavonoids by yeast. *Journal of Agricultural and Food Chemistry*. 2016;**64**:5525-5530. DOI: 10.1021/acs.jafc.6b02210
- [95] Bartmanska A, Huszcza E, Tronina T. Transformation of isoxanthohumol by fungus. *Journal of Molecular Catalysis B: Enzymatic*. 2009; **61**:221-224. DOI: 10.1016/j.molcatb.2009.07.008
- [96] Chen X, Mukwaya E, Wong MS, Zhang Y. A systematic review on biological activities of prenylated flavonoids. *Pharmaceutical Biology*. 2014;**52**:655-660. DOI: 10.3109/13880209.2013.853809
- [97] Cao H, Chen X, Jassbi AR, Xiao J. Microbial transformation of bioactive flavonoids. *Biotechnology Advances*. 2015;**33**:214-223. DOI: 10.1016/j.biotechadv.2014.10.012
- [98] Bartmanska A, Tronina T, Huszcza E. Transformation of 8-prenylnaringenin by *Absidia coerulea* and *Beauveria bassiana*. *Bioorganic & Medicinal Chemistry Letters*. 2012;**22**: 6451-6453. DOI: 10.1016/j.bmcl.2012.08.060

- [99] Tronina T, Strugala P, Poplonski J, Wloch SS, Bartmanska A, Huszcza E. The influence of glycosylation of natural and synthetic prenylated flavonoids on binding to human serum albumin and inhibition of cyclooxygenases COX-1 and COX-2. *Molecules*. 2017;**22**: 1230-1250. DOI: 10.3390/molecules22071230
- [100] Haraguchi H, Inoue J, Tamura Y, Mizutani K. Antioxidative components of *Psoralea corylifolia* (Leguminosae). *Phytotherapy Research*. 2002;**16**:539e544. DOI: 10.1002/ptr.972
- [101] Lim SH, Ha TY, Ahn J, Kim S. Estrogenic activities of *Psoralea corylifolia* L, seed extracts and main constituents. *Phytomedicine*. 2011;**18**: 425-430. DOI: 10.1016/j.phymed.2011.02.002
- [102] Luo J, Liang Q, Shen Y, Chen X, Yin Z, Wang M. Biotransformation of bavachinin by three fungal cell cultures. *Journal of Bioscience and Bioengineering*. 2014;**117**:191-196. DOI: 10.1016/j.jbiosc.2013.08.001
- [103] Tronina T, Bartmanska A, Milczarek M, Wietrzyk J, Poplonski J, Roj E, et al. Antioxidant and antiproliferative activity of glycosides obtained by biotransformation of xanthohumol. *Bioorganic & Medicinal Chemistry Letters*. 2013;**23**: 1957-1960. DOI: 10.1016/j.bmcl.2013.02.031
- [104] Kim HJ, Lee I-S. Microbial metabolism of the prenylated chalcone xanthohumol. *Journal of Natural Products*. 2006;**69**:1522-1524. DOI: 10.1021/np060310g
- [105] Bartmanska A, Huszcza E, Tronina T. Transformation of isoxanthohumol by fungi. *Journal of Molecular Catalysis B: Enzymatic*. 2009; **61**:221-224. DOI: 10.1016/j.molcatb.2009.07.008
- [106] Tronina T, Bartmanska A, Poplonski J, Huszcza E. Transformation of xanthohumol by *Aspergillus ochraceus*. *Journal of Basic Microbiology*. 2014;**54**: 66-71. DOI: 10.1002/jobm201200320
- [107] Tronina T, Bartmanska A, Filip-Psurska B, Wietrzyk J, Poplonski J, Huszcza EA. Fungal metabolites of xanthohumol with potent antiproliferative activity on human cancer cell lines in vitro. *Bioorganic & Medicinal Chemistry*. 2013;**21**: 2001-2006. DOI: 10.1016/j.bmc.2013.01.026
- [108] Stompor M, Potaniec B, Szumny A, Zieliński P, Żołnierczyk AK, Anioł M. Microbial synthesis of dihydrochalcones using *Rhodococcus* and *Gordonia* species. *Journal of Molecular Catalysis B: Enzymatic*. 2013a;**97**:283-288
- [109] Abourashed EA, Mikell JR, Khan IA. Bioconversion of silybin to phase I and II microbial metabolites with retained antioxidant activity. *Bioorganic & Medicinal Chemistry*. 2012;**20**:2784-2788. DOI: 10.1016/j.bmc.2012.03.046
- [110] Choi J-M, Han S-S, Kim H-S. Industrial applications of enzyme biocatalysis: Current status and future aspects. *Biotechnology Advances*. 2015; **33**:1443-1454. DOI: 10.1016/j.biotechadv.2015.02.014
- [111] De Carvalho CCCR. Enzymatic and whole cell catalysis: Finding new strategies for old processes. *Biotechnology Advances*. 2011;**29**:75-83
- [112] Muñoz SD, Hoyos P, Hernáiz MJ, Alcántara AR, Sánchez-Moreno JM. Industrial biotransformations in the synthesis of building blocks leading to enantiopure drugs. *Bioresource Technology*. 2012;**115**:196-207. DOI: 10.1016/j.biortech.2011.11.131
- [113] Patel RN. Biocatalysis for synthesis of pharmaceuticals. *Bioorganic &*

- Medicinal Chemistry. 2018;**26**: 1252-1274. DOI: 10.1016/j.bmc.2017.05.023
- [114] Van Den Burg B. Extremophiles as a source for novel enzymes. *Current Opinion in Microbiology*. 2003;**6**: 213-218. DOI: 10.1016/S1369-5274(03)00060-2
- [115] Antranikian G, Vorgias CE, Bertoldo C. Extreme environments as a resource for microorganisms and novel biocatalysts. *Advances in Biochemical Engineering/Biotechnology*. 2005;**96**: 219-262. DOI: 10.1007/b135786
- [116] Elleuche S, Schroder C, Sahm K, Antranikian G. Extremozymes-biocatalysts with unique properties from extremophilic microorganisms. *Current Opinion in Biotechnology*. 2014;**29**:116-123. DOI: 10.1016/j.copbio.2014.04.003
- [117] Falcicchio P, Wolterink-Van Loo S, Franssen MC, van der Oost J. DHAP-dependent aldolases from (hyper) thermophiles: Biochemistry and applications. *Extremophiles*. 2014;**18**: 1-13. DOI: 10.1007/s00792-013-0593-x
- [118] Urbietta MS, Donati ER, Chan KG, Shahar S, Li Sin L, Goh KM. Thermophiles in the genomic era: Biodiversity, science, and applications. *Biotechnology Advances*. 2015;**33**: 633-647. DOI: 10.1016/j.biotechadv.2015.04.007
- [119] Woodward J, Orr M, Cordray K, Greenbaum E. Biotechnology: Enzymatic production of biohydrogen. *Nature*. 2000;**405**:1014-1015
- [120] Soliman N, Abdel-Fattah Y, Mostafa HE, Gaballa A. Heterologous expression of thermostable esterase gene from *Geobacillus thermoleovorans* YN under different expression promoters. *International journal of Environmental Science and Technology*. 2014;**11**:119-126. DOI: 10.1007/s13762-013-0360-7
- [121] Saranya P, Kumari HS, Rao BP, Sekaran G. Lipase production from a novel thermo-tolerant and extreme acidophile *Bacillus pumilus* using palm oil as the substrate and treatment of palm oil-containing wastewater. *Environmental Science and Pollution Research International*. 2014;**21**: 3907-3919. DOI: 10.1007/s11356-013-2354-x
- [122] Gudiukaitė R, Gegeckas A, Kazlauskas D, Citavicius D. Influence of N-and/or C-terminal regions on activity, expression, characteristics and structure of lipase from *Geobacillus* sp. 95. *Extremophiles*. 2014;**18**:131-145. DOI: 10.1007/s00792-013-0605-x
- [123] Abraham J, Gea T, Sánchez A. Substitution of chemical dehairing by proteases from solid state fermentation of hair wastes. *Journal of Cleaner Production*. 2014;**74**:191-198
- [124] Staufenberger T, Imhoff JF, Labes A. First crenarchaeal chitinase found in *Sulfolobus tokodaii*. *Microbiological Research*. 2012;**167**: 262-269
- [125] Barabote RD, Parales JV, Guo Y-Y, Labavitch JM, Parales RE, Berry AM. Xyn10A, a thermostable endoxylanase from *Acidothermus cellulolyticus* 11B. *Applied and Environmental Microbiology*. 2010;**76**:7363-7366
- [126] Kang J, Park K-M, Choi K-H, Park C-S, Kim G-E, Kim D, et al. Molecular cloning and biochemical characterization of a heat-stable type I pullulanase from *Thermotoga neapolitana*. *Enzyme and Microbial Technology*. 2011;**48**:260-266