We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Fungal Endophyte-Host Plant Interactions: Role in Sustainable Agriculture

Tamanreet Kaur

Abstract

Fungal endophytes that live inside plant tissues without causing any apparent symptoms in the host plant are important components of plant micro-ecosystems. Endophytic fungi confer profound impacts on their host plants by enhancing their growth, increasing their fitness, strengthening their tolerances to pests and diseases. Moreover, fungal endophytes symbiotic with host plant produce a plethora of bioactive secondary metabolites that are expressed as defensive weapons to protect the host plant against various abiotic stresses. Currently, main focus in endophytic fungi research is associated with the ability of these microorganisms to produce and accumulate biologically active metabolites as these are potent source of novel natural products useful in agriculture sector.

Keywords: fungal endophyte, symbiosis, secondary metabolites, stress, sustainable agriculture

1. Introduction

Over reliance of synthetic pesticides in crop fields from late 1940 to mid-1960s resulted in a number of adverse environmental impacts such as secondary pest outbreak, insect resurgence, effects on non-target organisms, residual problem, environmental pollution, prompted an urgent need for alternative tactics to help make crop protection more sustainable. Biological control using micro-organism has gained much interest, being specific, low relative cost and low risk to ecosystem [1]. Among the various micro-organisms, endophytic fungi can make the chemical intensive crop production system more sustainable as it has ability to enhance plant growth, yield and increase plant fitness by providing biotic and abiotic stress tolerance [2, 3]. Endophytes (“endo” = within, “phyte” = plant) are the microorganisms that inhabit interior of plants especially leaves, stems, roots without causing any apparent harm to the host [4]. These are ubiquitous having rich biodiversity and found in every plant species as nearly 3,00,000 plant species exist on earth with each individual plant host having one or more than one endophytes [5]. Endophytic fungi are considered as plant mutualists as they receive nutrition and protection from host plant while the host plant may benefit from enhanced competitive abilities and increased resistance to herbivores, pathogens and various abiotic stresses [6]. It spends whole or part of their life cycle colonizing inters- and/or intra-cellularly within the healthy tissues of the host plant without causing visible signs

of infection [7, 8]. Moreover, fungal endophytes have gained significant interest in sustainable agriculture due to their great potential to contribute to secondary compounds with unique structure, including alkaloids, benzopyranones, chinones, flavonoids, phenolic acids, quinones, steroids, terpenoids, tetralones, xanthenes, etc. [9–11] produced by the fungi or by the plant due to interaction with the fungi. Among the microorganisms, fungal endophytes are the largest group producing secondary metabolites. Fungal toxins produced by these biotic metabolites contribute to plants tolerance towards various biotic and abiotic stresses. Fungal endophytes are known to produce bioactive compounds toxic to insects, nematodes, produces extracellular enzymes (cellulases, proteinase, lipases, esterases) for degradation of dead soil biomass, solubilize insoluble phosphates and produce plant growth-promoting hormones (auxins, cytokinins, gibberellins). Endophyte infected plants manage plant growth under adverse conditions of drought, salinity, temperature and heavy metal stress through different mechanisms. This chapter outlines various approaches for the use of endophytic fungal inoculants to combat various stresses in agricultural fields, thus increasing global crop productivity.

2. Fungal endophyte-host plant association

The association between fungal endophytes and their host plant is due to their unique adaptations which enable the endophytes to harmonize their growth with their host plant [12]. The origin of endophytes is not clear due to complex association between the endophyte and its host plant and the multiplicity of the host's living environment. Exogenous and endogenous are the two hypotheses explaining the origin of endophytes. According to endogenous hypothesis, endophytes are gaged from the mitochondria and chloroplast of the plant, and so it has comparable genetic backgrounds to the host [13], whereas exogenous hypothesis believes that endophytes arrive from outside of the plant and got inserted into the host from root wound, induced channels, or surface [14]. During the long period of co-existence and evolutionary processes, different relationships have been established between endophytic fungi and their host plants ranging from (i) a continuum of mutualism, (ii) antagonism, and (iii) neutralism. As once inside the tissues of a host plant, the endophytic fungi assumed a quiescent (latent) state, either for the whole lifetime of the host plant (neutralism) or for an extended period of time (mutualism or antagonism) until environmental conditions are favorable for endophytic fungi [15]. Endophytes due to its cryptic existence also have its role of decomposers in ecosystem, as they are among the primary colonizers of dead plant tissues [16, 17].

2.1 Fungal endophytes

2.1.1 Transmission

The life history of endophytes in symbiotum with host plant has three modes of reproduction (**Figure 1**). They can either be transmitted (i) vertically from infected plant to offspring via seeds (*Neotyphodium* spp.), (ii) horizontally by sexual spore s from infected individuals (e.g. *Epichloe* spp.) or (iii) mixture of two life cycles [19]. The pure vertical transmission is asexual reproduction of intercellular hyphae of above ground tissues with no symptoms and transmitted vertically via seeds from infected plants to offspring (e.g. *Neotyphodium* spp.). In contrast, the pure horizontal transmission evolves sexual life cycle, relies on the production of contagious

Figure 1.
 Asexual and sexual life cycles of *Epichloe festucae* symbiotic with *Festuca* spp. [18].

sexual spores. These spores can only be produced on a fungal structure (stroma) surrounding the grass flag leaf sheath (e.g. some *Epichloe* spp.). Leaves accumulate numerous infections shortly after emergence by means of epiphytic germination of fungal propagules, followed by cuticular penetration or entry through stomata's [20–22] and grow intercellularly within healthy tissues [20, 23]. However, many *Epichloe* spp. use a third mode of reproduction. In this fungi choke some flowering tillers and produce sexual spores leaving majority of tillers uninfected and transmitted asexually via seeds [18]. Endophytes are transmitted vertically (systemic) and horizontally (non-systemic). Vertically transmitted endophytes are mutualistic, whereas those transmitted horizontally depict antagonism to the host [6, 24].

3. Fungal endophytes for sustainable agriculture

In view of escalating pollution and cost due to indiscriminate use of chemical pesticides, diverted researchers interest towards alternative eco-friendly and safe approaches to meet increasing demand of agriculture productivity. Sustainable agriculture requires the use of various strategies to increase or maintain the current rate of food production while minimizing damage to the environment and human health. Symbiotic endophytic fungal associations with crops offer wide range of benefits ranging from the promotion of plant growth to improvements in the tolerance of various biotic and abiotic stresses. Moreover, loss of useful endophytic microbes from crop plants during their domestication and long term cultivation also requires transfer of endophytes from wild relatives of crops to crop species.

4. Fungal endophytes: Biotic stress management

Endophytic fungi have gained importance in the area of agriculture because of their ability to confer resistance to various biotic stress conditions like insect herbivory, nematocidal attack and by aiding plant growth processes.

4.1 Fungal endophytes

4.1.1 Biocontrol agents

Fungal endophytes act as biocontrol agents as they can protect their host plants from pathogens and pests [25, 26]. The mechanism whereby endophytes deter herbivory is through production of antiherbivory/bioactive compounds [27–29] or complex interacting factors of metabolic processes in both the fungus and the plant after infection [26, 30]. These defensive compounds may deter feeding (antixenosis) or reduce insect performance (antibiosis) [31, 32]. Endophytic fungi release the specialized biologically active compounds without any observable damage to their host tissues [33]. Defensive compounds may be categorized into various functional groups: alkaloids, terpenoids, isocoumarin derivatives, quinones, flavonoids, chlorinated metabolites, phenol and phenolic acids and many others [7, 34].

1. **Alkaloids:** Alkaloids are the first reported fungal metabolites to have insecticidal activity. Alkaloids produced by the fungus or by plant in response to fungal infection increase host resistance to herbivores [4, 35]. Endophyte infected grasses contain a variety of alkaloids such as peramines, ergot alkaloids, lolitrems, loline alkaloids and which are absent in non-infected conspecifics [36, 37]. Alkaloids are the first reported fungal metabolites to have insecticidal activity. Most of the alkaloids have been detected in the cultures of grass associated endophytic fungi, such as sexual *Epichloe* spp. and asexual *Neotyphodium* spp. Fungal isolate determines the types of alkaloids produced and plant/fungal genotype interaction can modify the quantities of these alkaloids [38]. The alkaloids from fungal endophytes are categorized into three groups, amines and amides, indole derivatives and pyrrolizidines. Among amines and amides, peramine is toxic to insects without being harmful to mammals [39, 40]. It is a strong feeding deterrent for argentine stem weevil and several other insects [41, 42]. The levels of alkaloids and other toxins may be altered qualitatively depending on the plants physiological state. Ball et al. [43] verified that with plant aging, the amount of peramine decreases in leaves and reaches lower levels during inflorescence phase. The second group of amine and amide alkaloids is ergot alkaloids that also provide significant resistance against insect pests [44]. Feeding experiments with a variety of mammals indicate that ergot alkaloids have significant detrimental effects on mammalian health and reproduction [45, 46]. Among indole derivatives, the lolitrem C and F have been shown to confer resistance against a number of insect species [47]. Other indole derivatives namely chanoclavine, agroclavine and elymoclavine isolated from culture of *Neotyphodium* endophyte [34] were reported to be toxic to some insects and mammals [48]. Among Pyrrolizidines, the saturated aminopyrrolizidine alkaloids as norloline, N-formylloline, N-acetylnorloline, N-acetylloline were exclusively found in endophyte infected grasses of *F. arundinacea* (infected with *Neotyphodium coenophialum*) and *Festuca pratensis* (with *Neotyphodium uncinatum*) [49]. A number of feeding experiments have demonstrated the insecticidal and insect feeding deterrent activities of these lolines [50–52]. Lolines in addition to the well documented effect on insects are also nematicidal [53].
2. **Terpenoids:** Second group of endophytic toxins include terpenoids isolated from some endophytic cultures originating from a variety of host plants. Sesquiterpenes and diterpenes are among the identified terpenoids. Sesquiterpenes as of heptelidic acid and hydroheptelidic acid isolated from *Phyllosticta* sp., an endophytic fungus of balsam fir (*Abies balsamea*) exhibited toxicity to

spruce budworm, *Choristoneura fumiferana* (Clemens) larvae [54]. Two insect toxins, pimarane and diterpene were isolated from an unidentified endophytic fungus symbiotic with needle of *A. balsamea* [54]. Two benzofuran carrying normonoterpene derivatives, toxic to spruce budworm larvae were characterized from an endophytic culture obtained from wintergreen (*Gaultheria procumbens*) [55].

3. **Isocoumarin derivatives:** Toxicity of isocoumarin related metabolites from the conifer endophyte cultures showed toxicity against cells and/ or larvae of spruce budworm [56].
4. **Quinones:** Rugulosin, a metabolite of endophytic fungus *Hormonema dematioides* from balsam fir has been reported to have insecticidal activity [54]. An unidentified endophytic culture isolated from eastern larch (*Larix laricina*) produced a quinone derivative, which was toxic to spruce budworm larvae [55].
5. **Flavonoids:** Among the flavonoids, triclin and related flavone glycosides isolated from endophyte infected blue grass (*Poa ampla*) exhibited toxicity against mosquito larvae [56].
6. **Chlorinated metabolites:** Insecticidal chlorinated metabolite, heptelidic acid chlorohydrins were isolated from cultures of balsam fir needle endophyte *Phyllosticta* spp. [57].
7. **Phenol and phenolic acids:** Phenol and phenolic acids are frequently detected in cultures of endophytes and have pronounced biological activities. Singh et al. [58] purified phenolic compound from ethyl acetate extract of endophytic *Cladosporium* sp. isolated from guduchi (*Tinospora cordifolia*), which induced significant mortality and adversely affected development and survival of tobacco cutworm, *Spodoptera litura* (Fabricius).

Since the 1980s, there is accumulating evidence about factors that influence the outcome of grass–endophyte–insect interactions. Webber [59] was probably the first worker to report plant protection given by fungal endophyte *Phomopsis oblonga* in elm trees (*Ulmus* spp.) against the elm bark beetle, *Physocnemum brevilineum* (Say). Majority of studies for herbivore performance on native grass species symbiotic with endophytic fungi are more consistent showing negative effects including increased mortality [60], reduced mass [61, 62] and decelerated development time [63]. Afkhami et al. [62] reported that bird cherry oat aphid, *Rhopalosiphum padi* (Linnaeus) damaged more endophyte free nodding fescue (*Festuca subverticillata*) than endophyte symbiotic *F. subverticillata*, while positive effect of endophyte infection was reported on eastern lubber grasshopper, *Romalea guttata* (Houttuyn) that preferentially consumed endophyte symbiotic *F. subverticillata* over endophyte free. Similarly increase in growth rate was recorded in third to fifth instars of fall armyworm, *Spodoptera frugiperda* (J.E. Smith) feeding on *N. coenophialum* infected tall fescue [63].

4.2 Fungal endophytes

4.2.1 Nematicidal agents

Fungal endophytes act as nematicidal agents as they are known to produce some compounds which are toxic to nematodes. Diedhiou et al. [64] demonstrated reduced

nematicidal activity by an endophytic fungus, *Fusarium oxysporum*, against the plant parasitic nematode *Meloidogyne incognita* in tomato plant. Schwarz et al. [65] reported that several endophytic fungi isolated from above-ground plant organs produced bioactive compound, 3-hydroxypropionic acid (HPA) extracted by bioactivity-guided fractionation of fungal extracts that showed selective nematicidal activity against *M. incognita* with LD50 values of 12.5–15 µg/ml. Similarly, Felde et al. [66] found that combined inoculations of endophytic fungal isolates *Trichoderma atroviride* and *F. oxysporum* is considered an alternative to improve and increase banana yield that reduces the population of burrowing nematode, *Radopholus similis* (Cobb), an important parasitic nematode on banana.

4.3 Fungal endophytes

4.3.1 Phytohormone production

Endophytes can actively or passively regulate the plant growth by solubilization of phosphate, enhance uptake of phosphorus (P), and/ or plant hormones such as auxin, abscisins, ethylene, gibberellic acid (GA), and indole acetic acid (IAA) [67, 68], among these Gibberellic acid is an important phytohormone. The phytohormone GA, a diterpenoid complex, controls the growth of plants, and promotes flowering, stem elongation, seed germination, and ripening [69, 70]. Fungal endophytes *Sebacina vermifera*, *Piriformospora indica*, *Colletotrichum* and *Penicillium* are distinguished to have better plant growth promoting effects under unfavorable conditions due to their ability to synthesize enzymes and bioactive metabolites [71–73]. Hamayun et al. [69] reported that fungal endophyte, *Cladosporium sphaerospermum* isolated from soybean plant (*Glycine max*) produced gibberellic acid that induced plant growth in rice and soybean. Metabolite pestalotin analogue, isolated from the endophytic *Pestalotiopsis microspora* exhibited significant gibberellin activity in winter-hazel seeds (*Distylium chinense*) and increased their germination rate [74]. Endophytes, *Fusarium tricinctum* and *Alternaria alternata* produced derivatives of plant hormone indole acetic acid that enhanced the plant growth [75]. A study conducted by Johnson et al. [76] on root colonizing endophyte *P. indica* found that association of fungal endophytes with roots modulated phytohormones involved with growth and development of host plant and enhanced nutrient uptake and translocation especially of phosphorus and nitrogen from the soil.

4.4 Fungal endophytes

4.4.1 Agriculturally important enzyme production

Degradation of the dead soil biomass by fungal endophytic is a major step in bringing the utilized nutrients back to the ecosystem that improves soil quality. Endophytic fungi is reported to produce various extracellular hydrolases including cellulase, laccase, pectinase, phosphatase, lipase, xylanase, and proteinase as a resistance mechanism against pathogenic invasion [77] and to obtain nutrition from host as these enzymes break macromolecules such as lignin, sugar-based polymers, proteins, organic phosphate, and carbohydrates to micromolecules that are transported throughout the cells for metabolism and help in host symbiosis process [78]. Sunitha et al. [79] isolated and identified approximately 50 endophytic fungal strains having amylase, laccase, cellulase, pectinase, lipase and protease enzymes. Study conducted by He et al. [80] explained that endophytic fungal species have ability to decompose organic components, including lignin,

cellulose, and hemicelluloses that facilitates nutrient cycling. Chathurdevi and Gowrie [81] reported that the endophytic fungi species isolated from medicinal plant *Cardiospermum halicacabum* can support plant growth to overcome the adverse conditions through producing different extracellular enzymes. Fungal chitinases enzyme have vital role in degradation and cycling of carbon and nitrogen from chitin molecule. Chitin molecule is a linear homopolymer of β -1,4N-acetylglucosamine can be obtained from insect's exoskeleton, crustacean's shells, and fungal cell wall. Many fungal endophytes isolated from leaves of trees of Southern India have shown the production of chitinases [82]. An endophyte, *Acremonium zeae*, isolated from maize is reported to produce the extracellular enzyme hemicellulase, which may be used in the bioconversion of lignocellulosic biomass into fermentable sugars [83].

5. Fungal endophytes: Abiotic stress management

Agricultural productivity is significantly threatened by various abiotic stresses. Environmental stresses such as drought, salinity, temperature can collectively cause more than 50% yield losses worldwide [84]. Plants can tolerate abiotic stress by two mechanisms: (i) via activation of response systems directly after exposure to stress [67] (ii) biochemical compounds that are synthesized by fungal endophytes, acts as anti-stress agents [85]. Experimental studies also confirmed that endophytic fungi can help the host plants from environmental stress conditions such as drought, salts, high temperatures and heavy metals and can thus increase the plant growth.

5.1 Drought stress

Among the abiotic stresses, water stress commonly, known as 'drought', is considered as one of the major challenges to crop production worldwide [86]. Drought has a negative impact on the plant growth rate, germination rates, membrane loss of its integrity, repression of photosynthesis, and increase in the productivity of reactive oxygen species [87, 88]. Fungal endophyte infected plants enhance drought tolerance by increased accumulation of solutes in tissues, or by reduced leaf conductance and a slowdown of the transpiration stream, or due to formation of thicker cuticle as compared to non-infected plants [67]. Chippa et al. [89] reported that endophytic, *Neotyphodium* spp. is reported to enhance drought tolerance in grass plant by stomatal and osmoregulations and protect plants in drought and nitrogen starvation. Experimental studies on lavender plants inoculated with *Glomus* spp. showed that these plants accumulated solutes in tissues thereby exhibiting better drought tolerance by improving water contents, N and P contents and root biomass [90, 91]. Moreover, plants harboring endophytes consumes significantly less water and had enhanced biomass than non-symbiotic plants. For instance, endophytes *Chaetomium globosum* and *P. resedanum* isolated from sweet pepper (*Capsicum annuum*) plants enhanced shoot length and biomass of the host plants challenged by drought stress [92, 93]. Similarly, Redman et al. [72] found that inoculation of endophytes *Fusarium culmorum* and *Curvularia protuberata* in drought-affected rice plants resulted in increased biomass than of non-inoculated plants. Fungal endophyte colonization also results in higher chlorophyll content and leaf area in plants under drought stress than non-colonized plant. Higher chlorophyll concentration is related with higher photosynthetic rate [94]. For instance, enhanced photosynthesis rate was recorded in drought stressed *C. annuum* plants colonized by endophytes *C. globosum* [95] and *P. resedanum* [96].

5.2 Salinity stress

High salinity due to extreme climatic conditions and misuse of agricultural land over the past few decades has led to high salinity, which is a limiting factor to global agricultural productivity [97]. Soil salinity is the accumulation of water soluble salts in soil that affects its physical and chemical properties thereby reducing soil's agricultural output [98]. Reactive oxygen species (SOD, CAT, APX) are formed in plants on onset of salt and osmotic stress. Endophytic *Piriformospora indica* induces salt stress tolerance by elevation of antioxidant enzymes [99]. These are involved in the removal of reactive oxygen species either directly or indirectly via regeneration of ascorbate and glutathione in the cell. Experimental studies by Rodriguez et al. [100] reported that constant exposure of non-symbiotic plants dunegrass (*Leymus mollis*) to 500 mmol/l NaCl solution, became severely wilted and desiccated within 7 days and were dead after 14 days. In contrast, symbiotic plants infected with *F. culmorum* showed wilting symptoms only after they were exposed to 500 mmol/l NaCl solution for 14 days.

5.3 Temperature stress (low and high)

High temperature is a major obstacle in crop production that results in major cellular damage such as protein degradation and aggregation [101]. Whereas, low temperature can cause impaired metabolism due to inhibition of enzyme reactions, interactions among macromolecules, changes in protein structure, and modulating cell membrane properties [102]. Endophytic, *Curvularia* spp. is proven to confer thermal tolerance ability plants like tomato, watermelon, and wheat [103]. Herbal plant woolly rosette grass (*Dichanthelium lanuginosum*) that lives in the areas where soil temperatures can reach up to 57°C, the presence of endophytic fungi *Curvularia* sp. protects the plant from temperature stress better than endophyte free plants [104]. Experimental demonstration by Redman et al. [103] showed that grass *D. lanuginosum* survival in soil temperatures ranging between 38 and 65°C is directly linked to its association with the fungus *C. protuberata* and its mycovirus, *Curvularia* thermal tolerance virus (CThTV). Moreover, cold stress tolerance was conferred in germinated seeds of rice under laboratory conditions by *C. protuberata* isolated from *D. lanuginosum* thriving in geothermal soils [72].

5.4 Heavy metal stress

Heavy metal contamination due to increased industrialization has recently received attention because heavy metals cannot be itself degraded [105]. Toxicity by heavy metals can cause the loss of about 25–80% of various cultivated crops. Heavy metals being very toxic to roots of cultivated crop plants can cause poor development of the root system [106]. Endophytic fungi possess metal sequestration or chelation systems that increases tolerances of their host plants to heavy metals via enhancements of antioxidative system thereby changing heavy metal distribution in plant cells and detoxification of heavy metal, thus assisting their hosts to survive in contaminated soil [107, 108]. For instance, dark septate root endophytes (DSEs), *Phialocephala fortinii* can produce the black biopolymer melanin, which can be synthesized from phenolics and binds heavy metals [109] that keep heavy metal ions away from living, plant cells [110]. Siderophores being metal-chelating compounds [111, 112] released from roots into the rhizosphere can be helpful in inhibiting absorption of heavy metals into plant cells as siderophores can form complexes with heavy metals which are not easily absorbed by plant

roots. Yamaji et al. [113] recorded that endophytes *P. fortinii* and *Rhizodermea veluensis* showed an ability to produce siderophores that probably affects heavy metal exclusion in the rhizosphere.

6. Conclusion

Fungal endophytes can be a significant component of sustainable agriculture, being safe, cost-effective, have ability to produce various compounds like phytohormones, defensive compounds, solubilize phosphates, extracellular enzymes, siderophore production, inhibiting plant pathogens, and promoting plant growth. Over the last decade, sharp rise in study of fungal endophytes is seen as they hold huge potential in agricultural sector. However, most of the research on endophytes is still at an experimental level in lab or greenhouse. For permitting the practical use of these endophytes in agriculture it is extremely necessary to encourage field experiments to determine the effectiveness of the endophytes under real world conditions. Simultaneously, it is also necessary to build awareness of this new research field among farmers to improve interactions and collaboration with scientists working in different fields, thereby encouraging the adoption of endophytes in agriculture and maximizing their benefits. If endophytes become feasible in agricultural sector, their practical aspects will also have to be researched so that farmers can learn how to integrate these endophyte species within pre-existing eco-friendly agricultural methods so as to ensure continuity in the approach to sustainability. Moreover, scientific research has to be also focused on use of genetically modified endophytes made by combining endophytes having two or more different ecological roles, such as the suppression of diseases and insect pests to simultaneously improve plant yields and its defensive properties. Thus, optimization of microbial functions to enhance crop production and protection is also required.

Conflict of interest

No conflict of interest is indulged.

Abbreviations

HPA	3-hydroxypropionic acid
P	phosphorus
GA	Gibberellic acid
IAA	indole acetic acid
SOD	superoxide dismutase
CAT	catalase
APx	ascorbate peroxidase
CThTV	<i>Curvularia</i> thermal tolerance virus
DSEs	dark septate root endophytes

IntechOpen

IntechOpen

Author details

Tamanreet Kaur

Department of Zoology, Kanya Maha Vidyalaya, Jalandhar, India

*Address all correspondence to: tamanreetkaur@gmail.com

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Castillo MA, Moya P, Hernandez E, Primo-Yufera E. Susceptibility of *Ceratitidis capitata* Wiedemann (Diptera: Tephritidae) to entomopathogenic fungi and their extracts. *BioControl*. 2000;**19**:274-282. DOI: 10.1006/bcon.2000.0867
- [2] Tanaka A, Tapper BA, Popay A, Parker EJ, Scott B. A symbiosis expressed non-ribosomal peptide synthetase from a mutualistic fungal endophyte of perennial ryegrass confers protection to the symbiotum from insect herbivory. *Molecular Microbiology*. 2005;**57**:1036-1050. DOI: 10.1111/j.1365-2958.2005.04747.x
- [3] Vega FE, Posada F, Aime MC, Pava-Ripoll M, Infante F, Rehner SA. Entomopathogenic fungal endophytes. *Biological Control*. 2008;**46**:72-82. DOI: 10.1016/j.biocontrol.2008.01.008
- [4] Azevedo JL, Maccheroni W Jr, Periera JO, Araujo WL. Endophytic microorganisms: A review on insect control and recent advances on tropical plants. *Electronic Journal of Biotechnology*. 2000;**3**:40-65. DOI: 10.2225/vol3-issue1-fulltext-4
- [5] Smith SA, Tank DC, Boulanger LA, Bascom-Slack CA, Eisenman K, Kingery D, et al. Bioactive endophytes warrant intensified exploration and conservation. *PLoS One*. 2008;**3**:e3052. DOI: 10.1371/journal.pone.0003052
- [6] Saikkonen K, Faeth SH, Helander M, Sullivan TJ. Fungal endophytes: A continuum of interactions with host plants. *Annual Review of Ecology and Systematics*. 1998;**29**:319-343. DOI: 10.1146/annurev.ecolsys.29.1.319
- [7] Tan RX, Zou WX. Endophytes: A rich source of functional metabolites. *Natural Product Reports*. 2001;**18**:448-459. DOI: 10.1039/B100918O
- [8] Hartley SE, Gange AC. Impacts of plant symbiotic fungi on insect herbivores: Mutualism in a multitrophic context. *Annual Review of Entomology*. 2009;**54**:323-342. DOI: 10.1146/annurev.ento.54.110807.090614
- [9] Jalgaonwala RE, Mohite BV, Mahajan RT. Natural products from plant associated endophytic fungi. *Journal of Microbiology and Biotechnology Research*. 2011;**1**:21-32
- [10] Joseph B, Priya RM. Bioactive compounds from endophytes and their potential in pharmaceutical effect: A review. *American Journal of Biochemistry and Molecular Biology*. 2011;**1**:291-309. DOI: 10.3923/ajbmb.2011.291.309
- [11] Pimentel MR, Molina G, Dionisio AP, Marostica MR, Pastore GM. Use of endophytes to obtain bioactive compounds and their application in biotransformation process. *Biotechnology Research International*. 2011:1-11. Article ID: 566286. DOI: 10.4061/2011/576286
- [12] Verma VC, Singh SK, Kharwar RN. Histological investigation of fungal Endophytes in healthy tissues of *Azadirachta indica* A. Juss. *Kasetsart Journal (Natural Science)*. 2012;**46**:229-237
- [13] Wen CY. Recent advances and issues on the endophyte. *Chinese Journal of Ecology*. 2004;**2**:86-91
- [14] Li WK. Endophytes and natural medicines. *Chinese Journal of Natural Medicines*. 2005;**4**:193-199
- [15] Sieber TN. Endophytic fungi in forest trees: Are they mutualists? *Fungal Biology Reviews*. 2007;**21**:75-89. DOI: 10.1016/j.fbr.2007.05.004
- [16] Kumaresan V, Suryanarayanan TS. Endophyte assemblage in young,

mature and senescent leaves of *Rhizophora apiculata*: Evidence for the role of endophytes in mangrove litter degradation. *Fungal Diversity*. 2002;**9**:81-91

[17] Oses R, Valenzuela S, Freer J, Sanfuentes E, Rodriguez J. Fungal endophytes in xylem of healthy chilean trees and their possible role in early wood decay. *Fungal Diversity*. 2008;**33**:77-86

[18] Bush LP, Wilkinson HH, Schardl CL. Bioprotective alkaloids of grass-fungal endophyte symbioses. *Plant Physiology*. 1997;**114**:1-7. DOI: 10.1104/pp.114.1.1

[19] Schardl CL, Leuchtman A, Chung KR, Penny D, Siegel MR. Coevolution by common descent of fungal symbionts (*Epichloe* spp.) and grass hosts. *Molecular Biology and Evolution*. 1997;**14**:133-143. DOI: 10.1093/oxfordjournals.molbev.a025746

[20] Lebron L, Lodge DJ, Laureano SL, Bayman P. Where is the gate to party? *Phytopathology*. 2001;**91**:116

[21] Arnold AE. Fungal endophytes in Neotropical trees: Abundance, diversity and ecological interactions [Thesis]. Tucson: University of Arizona; 2002

[22] Arnold AE, Herre EA. Canopy cover and leaf age affect colonization by tropical fungal endophytes: Ecological pattern and process in *Theobroma cacao* (Malvaceae). *Mycologia*. 2003;**95**:388-398. DOI: 10.1080/15572536.2004.11833083

[23] Arnold AE, Mejia L, Rojas E, Maynard Z, Robbins N, Herre EA. Organismos endofiticos: Microorganismos en plantas. In: Barrantes RM, editor. *El uso de Microorganismos Beneficos en la Agricultura Moderna*. Costa Rica: Guacimo; 2001. pp. 1-20

[24] Schardl CL, Liu JS, White JF, Finkel RA, An Z, Siegel MR. Molecular

phylogenetic relationships of nonpathogenic grass mycosymbionts and clavicipitaceous plant pathogens. *Plant Systematics and Evolution*. 1991;**178**:27-41. DOI: 10.1007/BF00937980

[25] Bae H, Sicher RC, Kim MS, Kim S-H, Strem MD, Melnick RL, et al. The beneficial endophyte *Trichoderma hamatum* isolate DIS 219b promotes growth and delays the onset of the drought response in *Theobroma cacao*. *Journal of Experimental Botany*. 2009;**60**:3279-3295. DOI: 10.1093/jxb/erp165

[26] Akello J, Dubois T, Gold CS, Coyne D, Nakavuma J, Paparu P. *Beauveria bassiana* (Balsamo) Vuillemin as an endophyte in tissue culture banana (*Musa* spp.). *Journal of Invertebrate Pathology*. 2007;**96**:34-42. DOI: 10.1016/j.jip.2007.02.004

[27] Clay K, Schardl C. Evolutionary origins and ecological consequences of endophyte symbiosis with grasses. *American Naturalist*. 2002;**160**:99-127. DOI: 10.1086/342161

[28] Faeth SH, Hayes CJ, Gardner DR. Asexual endophytes in a native grass: Tradeoffs in mortality, growth, reproduction, and alkaloid production. *Microbial Ecology*. 2010;**60**:496-504. DOI: 10.1007/s00248-010-9643-4

[29] Torres MS, White JF Jr. Grass endophyte-mediated plant stress tolerance: Alkaloids and their functions. In: Seckbach J, Grube M, editors. *Symbiosis and Stress: Joint Ventures in Biology, Cellular Origin, Life in Extreme Habitats and Astrobiology*. New York: Springer; 2010. pp. 477-493

[30] Barbosa P, Krischik VA, Jones CG. *Microbial Mediation of Plant-Herbivore Interactions*. New York: Wiley; 1991. 530p

[31] Braman SK, Duncan RR, Engelke MC, Hanna WW, Hignight K,

Rush D. Grass species and endophyte effects on survival and development of fall armyworm (Lepidoptera: Noctuidae). *Journal of Economic Entomology*. 2002;**95**:487-492. DOI: 10.1603/0022-0493-95.2.487

[32] Crawford KM, Land JM, Rudgers JA. Fungal endophytes of native grasses decrease insect herbivore preference and performance. *Oecologia*. 2010;**164**:431-444. DOI: 10.1007/s00442-010-1685-2

[33] Liarzi O, Bucki P, Braun Miyara S, Ezra D. Bioactive volatiles from an endophytic *Daldinia* cf. *concentrica* isolate affect the viability of the plant parasitic nematode *Meloidogyne javanica*. *PLoS One*. 2016;**11**:e0168437. DOI: 10.1371/journal.pone.0168437

[34] Strobel G, Daisy B, Castillo U, Harper J. Natural products from endophytic microorganisms. *Journal of Natural Products*. 2004;**67**:257-268. DOI: 10.1021/np030397v

[35] Powell RG, Petroski RJ. Alkaloid toxins in endophyte infected grasses. *Natural Toxins*. 1992;**1**:163-170. DOI: 10.1002/nt.2620010304

[36] Muller CB, Krauss J. Symbiosis between grasses and asexual fungal endophytes. *Current Opinion in Plant Biology*. 2005;**8**:450-456. DOI: 10.1016/j.pbi.2005.05.007

[37] Zhang Y, Han T, Ming Q, Wu L, Rahman K, Qin L. Alkaloids produced by endophytic fungi: A review. *Natural Product Communications*. 2012;**7**:963-968

[38] Lane GA, Christensen MJ, Miles CO. Coevolution of fungal endophytes with grasses: The significance of secondary metabolites. In: Bacon CW, White JF Jr, editors. *Microbial Endophytes*. New York: Marcel Dekker; 2000. pp. 341-388

[39] Dew RK, Boissonneault GA, Gay N, Boling JA, Cross RJ, Cohen DA. The

effect of the endophyte (*Acremonium coenophialum*) and associated toxin of tall fescue on serum titer response to immunization and spleen cell flow cytometry analysis and response to mitogens. *Veterinary Immunology and Immunopathology*. 1990;**26**:285-295. DOI: 10.1016/0165-2427(90)90097-c

[40] Rowan DD, Latch GCM. Utilization of endophyte-infected perennial ryegrasses for increased insect resistance. In: Bacon CW, White JF Jr, editors. *Biotechnology of Endophytic Fungi of Grasses*. Boca Raton: CRC Press; 1994. pp. 169-183

[41] Rowan DD, Dymock JJ, Brimble MA. Effect of fungal metabolite peramine and analogs on feeding and development of argentine stem weevil (*Listronotus bonariensis*). *Journal of Chemical Ecology*. 1990;**16**:1683-1695. DOI: 10.1007/BF01014100

[42] Rowan DD. Lolitrems, peramine and paxilline: Mycotoxins of the ryegrass/endophyte interaction. *Agriculture, Ecosystems and Environment*. 1993;**44**:103-122. DOI: 10.1016/0167-8809(93)90041-M

[43] Ball OJP, Barker GM, Prestidge RA, Lauren DR. Distribution and accumulation of the alkaloid peramine in *Neotyphodium lolii*-infected perennial ryegrass. *Journal of Chemical Ecology*. 1997;**23**:1419-1434. DOI: 10.1023/B:J OEC.0000006473.26175.19

[44] Potter AD, Stokes TJ, Redmond TC, Schardl LC, Panaccione GD. Contribution of ergot alkaloids to suppression of a grass-feeding caterpillar assessed with gene knockout endophytes in perennial ryegrass. *Entomologia Experimentalis et Applicata*. 2008;**126**:138-147. DOI: 10.1111/j.1570-7458.2007.00650.x

[45] Gadberry MS, Denard TM, Spiers DE, Piper EL. Effects of feeding ergovaline on lamb performance in a heat stress environment. *Journal of*

Animal Science. 2003;**81**:1538-1545.
DOI: 10.2527/2003.8161538x

[46] Parish JA, McCann MA, Watson RH, Hoveland CS, Hawkins LL, Hill NS, et al. Use of nonergot alkaloid-producing endophytes for alleviating tall fescue toxicosis in sheep. Journal of Animal Science. 2003;**81**:1316-1322. DOI: 10.2527/2003.8151316x

[47] Parish JA, McCann MA, Watson RH, Piava NN, Hoveland CS, Parks AH, et al. Use of nonergot alkaloid-producing endophytes for alleviating tall fescue toxicosis in stocker cattle. Journal of Animal Science. 2003;**81**:2856-2868. DOI: 10.2527/2003.81112856x

[48] Ball OJP, Barker GM, Prestidge RA, Sprosen JM. Distribution and accumulation of the Mycotoxin Lolitrem B in *Neotyphodium lolii*-infected perennial ryegrass. Journal of Chemical Ecology. 1997;**23**:1435-1449. DOI: 10.1023/B:JOEC.0000006474.44100.17

[49] Schardl CL, Phillips TD. Protective grass endophytes: Where are they from and where are they going? Plant Disease. 1997;**81**:430-438. DOI: 10.1094/PDIS.1997.81.5.430

[50] Siegel MR, Latch GCM, Bush LP, Fannin FF, Rowan DD, Tapper BA, et al. Fungal endophyte-infected grasses: Alkaloid accumulation and aphid response. Journal of Chemical Ecology. 1990;**16**:3301-3315. DOI: 10.1007/BF00982100

[51] Riedell WE, Kieckhefer RE, Petroski RJ, Powell RG. Naturally occurring and synthetic loline alkaloid derivatives: Insect feeding behavior modification and toxicity. Journal of Entomological Science. 1991;**26**:122-129. DOI: 10.18474/0749-8004-26.1.122

[52] Jensen JG, Popay AJ, Tapper BA. Argentine stem weevil adults are affected by meadow fescue endophyte and its loline alkaloids. New Zealand

Plant Protection. 2009;**62**:12-18. DOI: 10.30843/nzpp.2009.62.4800

[53] Bacetty AA, Snook ME, Glenn AE, Noe JP, Nagabhyru P, Bacon CW. Chemotaxis disruption in *Pratylenchus scribneri* by tall fescue root extracts and alkaloids. Journal of Chemical Ecology. 2009;**35**:844-850. DOI: 10.1007/s10886-009-9657-x

[54] Calhoun LA, Findlay JA, Miller JD, Whitney NJ. Metabolites toxic to spruce budworm from balsam fir needle endophytes. Mycological Research. 1992;**96**:281-282. DOI: 10.1016/S0953-7562(09)80939-8

[55] Findlay JA, Li G, Penner PE, Miller JD. Bioactive isocoumarins and related metabolites from conifer endophytes. Journal of Natural Products. 1995;**58**:1759-1768. DOI: 10.1021/np50125a021

[56] Findlay JA, Buthelezi S, Li G, Seveck M, Miller JD. Insect toxins from an endophytic fungus from wintergreen. Journal of Natural Products. 1997;**60**:1214-1215. DOI: 10.1021/np970222j

[57] Ju Y, Sacalis JN, Still CC. Bioactive flavonoids from endophyte-infected blue grass (*Poa ampla*). Journal of Agricultural and Food Chemistry. 1998;**46**:3785-3788. DOI: 10.1021/jf980189m

[58] Singh B, Kaur T, Kaur S, Manhas RK, Kaur A. An alpha-glucosidase inhibitor from an endophytic *Cladosporium* sp. with potential as a biocontrol agent. Applied Biochemistry and Biotechnology. 2015;**175**:2020-2034. DOI: 10.1007/s12010-014-1325-0

[59] Webber J. A natural control of Dutch elm disease. Nature. 1981;**292**:449-451. DOI: 10.1038/292449a0

[60] Brem D, Leuchtman A. *Epichloe* grass endophytes increase herbivore

resistance in the woodland grass *Brachypodium sylvaticum*. *Oecologia*. 2001;**126**:522-530. DOI: 10.1007/s004420000551

[61] Bazely DR, Vicari M, Emmerich S, Filip L, Lin D, Inman A. Interactions between herbivores and endophyte-infected *Festuca rubra* from the Scottish islands of St Kilda, Benbecula and Rum. *Journal of Applied Ecology*. 1997;**34**:847-860. DOI: 10.2307/2405276

[62] Afkhami ME, Rudgers JA. Endophyte-mediated resistance to herbivores depends on herbivore identity in the wild grass *Festuca subverticillata*. *Environmental Entomology*. 2009;**38**:1086-1095. DOI: 10.1603/022.038.0416

[63] Bultman TL, Bell GD. Interaction between fungal endophytes and environmental stressors influences plant resistance to insects. *Oikos*. 2003;**103**:182-190. DOI: 10.1034/j.1600-0706.2003.11574.x

[64] Diedhiou PM, Hallmann J, Oerke EC, Dehne HW. Effects of arbuscular mycorrhizal fungi and a non-pathogenic *Fusarium oxysporum* on *Meloidogyne incognita* infestation of tomato. *Mycorrhiza*. 2003;**13**:199-204. DOI: 10.1007/s00572-002-0215-4

[65] Schwarz M, Kopcke B, Weber RWS, Sterner O, Anke H. 3-Hydroxypropionic acid as a nematicidal principle in endophytic fungi. *Phytochemistry*. 2004;**65**:2239-2245. DOI: 10.1016/j.phytochem.2004.06.035

[66] Felde ZA, Pocasangre LE, Carnizares Monteros CA, Sikora RA, Rosales FE, Riveros AS. Effect of combined inoculations of endophytic fungi on the biocontrol of *Radopholus similis*. *InfoMusa*. 2006;**15**:12-18

[67] Malinowski DP, Belesky DP. Adaptations of Endophyte-infected cool-season grasses to environmental

stresses: Mechanisms of drought and mineral stress tolerance. *Crop Science*. 2000;**40**:923-940. DOI: 10.2135/cropsci2000.404923x

[68] Firakova S, Sturdikova M, Muckova M. Bioactive secondary metabolites produced by microorganisms associated with plants. *Biologia*. 2007;**62**:251-257. DOI: 10.2478/s11756-007-0044-1

[69] Hamayun M, Khan SA, Iqbal I, Na CI, Khan AL, Hwang YH, et al. *Chrysosporium pseudomercurarium* produces gibberellins and promotes plant growth. *Journal of Microbiology*. 2009;**47**:425-430. DOI: 10.1007/s12275-009-0268-6

[70] Yamaguchi S. Gibberellin metabolism and its regulation. *Annual Review of Plant Biology*. 2008;**59**:225-251. DOI: 10.1146/annurev.arplant.59.032607.092804

[71] Waller F, Achatz B, Baltruschat H, Fodor J, Becker K, Fischer M, et al. The endophytic fungus *Piriformospora indica* reprograms barley to salt stress tolerance, disease resistance, and higher yield. *Proceedings of the National Academy of Sciences of the United States of America*. 2005;**102**:13386-13391

[72] Redman RS, Kim YO, Woodward CJ, Greer C, Espino L, Doty SL, et al. Increased fitness of rice plants to abiotic stress via habitat adapted symbiosis: A strategy for mitigating impacts of climate change. *PLoS One*. 2011;**6**:e14823. DOI: 10.1371/journal.pone.0014823

[73] Hamilton CE, Bauerle TL. A new currency for mutualism? Fungal endophytes alter antioxidant activity in hosts responding to drought. *Fungal Diversity*. 2012;**54**:39-49. DOI: 10.1007/s13225-012-0156-y

[74] Li X, Guo Z, Deng Z, Yang J, Zou K. A new α -Pyrone derivative from

- endophytic fungus *Pestalotiopsis microspora*. Records of Natural Products. 2015;**9**:503-508
- [75] Khan AL, Hussain J, Al-Harrasi A, Al-Rawahi A, Lee IJ. Endophytic fungi: Resource for gibberellins and crop abiotic stress resistance. Critical Reviews in Biotechnology. 2015;**35**:62-74. DOI: 10.3109/07388551.2013.800018
- [76] Johnson JM, Alex T, Oelmuller R. *Piriformospora indica*: The versatile and multifunctional root endophytic fungus for enhanced yield and tolerance to biotic and abiotic stress in crop plants. Journal of Tropical Agriculture. 2014;**52**:103-122
- [77] Leo VV, Passari AK, Joshi JB, Mishra VK, Uthandi S, Ramesh N, et al. A novel triculture system (CC3) for simultaneous enzyme production and hydrolysis of common grasses through submerged fermentation. Frontiers in Microbiology. 2016;**7**:447. DOI: 10.3389/fmicb.2016.00447
- [78] Strong PJ, Claus H. Laccase: A review of its past and its future in bioremediation. Critical Reviews in Environmental Science and Technology. 2011;**41**:373. DOI: 10.1080/10643380902945706
- [79] Sunitha VH, Devi DN, Srinivas C. Extracellular enzymatic activity of endophytic fungal strains isolated from medicinal plants. World Journal of Agricultural Sciences. 2013;**9**:1-9. DOI: 10.5829/idosi.wjas.2013.9.1.72148
- [80] He X, Han G, Lin Y, Tian X, Xiang C, Tian Q, et al. Diversity and decomposition potential of endophytes in leaves of a *Cinnamomum camphora* plantation in China. Ecological Research. 2012;**27**:273-284. DOI: 10.1007/s11284-011-0898-0
- [81] Chathurdevi G, Gowrie SU. Endophytic fungi isolated from medicinal plant—A source of potential bioactive metabolites. International Journal of Current Pharmaceutical Research. 2016;**8**:50-56
- [82] Rajulu MBG, Thirunavukkarasu N, Suryanarayanan TS, Ravishankar JP, Gueddari NEE, Moerschbacher BM. Chitinolytic enzymes from endophytic fungi. Fungal Diversity. 2011;**47**:43-53. DOI: 10.1007/s13225-010-0071-z
- [83] Bischoff KM, Wicklow DT, Jordan DB, de Rezende ST, Liu S, Hughes SR, et al. Extracellular hemicellulolytic enzymes from the maize endophyte *Acremonium zeae*. Current Microbiology. 2009;**58**:499-503. DOI: 10.1007/s00284-008-9353-z
- [84] Wang Y, Frei M. Stressed food—The impact of abiotic environmental stresses on crop quality. Agriculture, Ecosystems and Environment. 2011;**141**:271-286. DOI: 10.1016/j.agee.2011.03.017
- [85] Schulz B, Boyle C, Draeger S, Rommert A-K, Krohn K. Endophytic fungi: A source of novel biologically active secondary metabolites. Mycological Research. 2002;**106**:996-1004. DOI: 10.1017/S0953756202006342
- [86] IPCC. Climate change 2007: Synthesis report. In: Pachauri RK, Reisinger A, editors. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Geneva: IPCC; 2007. p. 104
- [87] Granier C, Tardieu F. Reductions in area, cell number and cell area in sunflower leaves subjected to short water deficits with different timings. Variability in responses can be simulated using a simple model of leaf development. Plant Physiology. 1999;**119**:609-619. DOI: 10.1104/pp.116.3.991
- [88] Greenberg BM, Huang XD, Gerwing P, Yu XM, Chang P, Wu SS, et al. Phytoremediation of salt impacted

soils: Greenhouse and the field trials of plant growth promoting rhizobacteria (PGPR) to improve plant growth and salt phyto-accumulation. In: Proceeding of the 33rd AMOP Technical Seminar on Environmental Contamination and Response. Ottawa: Environment Canada; 2008. pp. 627-637

[89] Chhipa H, Deshmukh SK. Fungal Endophytes: Rising tools in sustainable agriculture production. In: Jha S, editor. *Endophytes and Secondary Metabolites*. Switzerland: Springer International Publishing; 2019. pp. 1-24. DOI: 10.1007/978-3-319-76900-4_26-1

[90] Porcel R, Aroca R, Cano C, Bago A, Ruiz-Lozano JM. Identification of a gene from the arbuscular mycorrhizal fungus *Glomus intraradices* encoding for a 14-3-3 protein that is upregulated by drought stress during the AM symbiosis. *Microbial Ecology*. 2006;**52**:575-582. DOI: 10.1007/s00248-006-9015-2

[91] Marulanda A, Barea JM, Azcon R. Stimulation of plant growth and drought tolerance by native microorganisms (AM fungi and bacteria) from dry environments: Mechanisms related to bacterial effectiveness. *Journal of Plant Growth Regulation*. 2009;**28**:115-124. DOI: 10.1007/s00344-009-9079-6

[92] Khan AL, Hamayun M, Hussain J, Kang SM, Lee IJ. The newly isolated endophytic fungus *Paraconiothyrium* sp. LK1 produces ascotoxin. *Molecules*. 2012;**17**:1103-1112. DOI: 10.3390/molecules17011103

[93] Khan AL, Waqas M, Hussain J, Al-Harrasi A, Lee IJ. Fungal endophyte *Penicillium janthinellum* LK5 can reduce cadmium toxicity in *Solanum lycopersicum* (Sitens and Rhe). *Biology and Fertility of Soils*. 2014;**50**:75-85. DOI: 10.1007/s11274-013-1378-1

[94] Davies FT, Potter JR, Linderman RG. Drought resistance of

mycorrhizal pepper plants independent of leaf P concentration response in gas exchange and water relations. *Plant Physiology*. 1993;**87**:45-53. DOI: 10.1111/j.1399-3054.1993.tb08789.x

[95] Khan AL, Shinwari ZK, Kim Y, Waqas M, Hamayun M, Kamran M, et al. Role of endophyte *Chaetomium globosum* lk4 in growth of *Capsicum annuum* by production of gibberellins and indole acetic acid. *Pakistan Journal of Botany*. 2012;**44**:1601-1607

[96] Khan AL, Waqas M, Lee IJ. Resilience of *Penicillium resedanum* LK6 and exogenous gibberellin in improving *Capsicum annuum* growth under abiotic stresses. *Journal of Plant Research*. 2014;**128**:259-268. DOI: 10.1007/s10265-014-0688-1

[97] Wicke B, Smeets E, Dornburg V, Vashev B, Gaiser T, Turkenburg W, et al. The global technical and economic potential of bioenergy from salt-affected soils. *Energy & Environmental Science*. 2011;**4**:2669-2681. DOI: 10.1039/C1EE01029H

[98] Hu Y, Schmidhalter U. Limitation of salt stress to plant growth. In: Hock B, Elstner CF, editors. *Plant Toxicology*. New York: Marcel Dekker Inc; 2002. pp. 91-224. DOI: 10.1201/9780203023884.ch5

[99] Baltruschat H, Fodor J, Harrach BD, Niemczyk E, Barna B, Gullner G, et al. Salt tolerance of barley induced by the root endophyte *Piriformospora indica* is associated with a strong increase in antioxidants. *New Phytologist*. 2008;**180**:501-510. DOI: 10.1111/j.1469-8137.2008.02583

[100] Rodriguez RJ, Henson J, Van Volkenburgh E, Hoy M, Wright L, Beckwith F, et al. Stress tolerance in plants via habitat-adapted symbiosis. *The ISME Journal*. 2008;**2**:404-416. DOI: 10.1038/ismej.2007.10

- [101] Hussain SS, Mehnaz S, KHM S. Harnessing the plant microbiome for improved abiotic stress tolerance. In: Egamberdieva D, Ahmad P, editors. *Plant Microbiome: Stress Response*. Singapore: Springer Nature; 2018. pp. 21-43. DOI: 10.1007/978-981-10-5514-0_2
- [102] Andreas T, Christophe C, Essaid AB. Physiological and molecular changes in plants at low temperatures. *Planta*. 2012;235:1091-1105. DOI: 10.1007/s00425-012-1641-y
- [103] Redman RS, Sheehan KB, Stout RG, Rodriguez RJ, Henson JM. Thermotolerance eneredated by plant/fungal symbiosis. *Science*. 2002;298:1581-1581
- [104] Stierle A, Strobel G, Stierle D. Taxol and taxane production by *Taxomyces andreanae*, an endophytic fungus of pacific yew. *Science*. 1993;260:214-216. DOI: 10.1126/science.8097061
- [105] Kidd P, Barcelo J, Bernal MP, Navari-Izzo F, Poschenrieder C, Shilev S, et al. Trace element behavior at the root-soil interface: Implications in phytoremediation. *Environmental and Experimental Botany*. 2009;67:243-259. DOI: 10.1016/j.envexpbot.2009.06.013
- [106] Singh LP, Gill SS, Tuteja N. Unraveling the role of fungal symbionts in plant abiotic stress tolerance. *Plant Signalling and Behaviour*. 2011;6:175-191. DOI: 10.4161/psb.6.2.14146
- [107] Likar M. Dark septate endophytes and mycorrhizal fungi of trees affected by pollution. In: Pirttila AM, Frank AC, editors. *Endophytes of Forest Trees*. Dordrecht: Springer Science+Business Media; 2011. pp. 189-201. DOI: 10.1007/978-94-007-1599-8
- [108] Wang JL, Li T, Liu GY, Smith JM, Zhao ZW. Unraveling the role of dark septate endophyte (DSE) colonizing maize (*Zea mays*) under cadmium stress: Physiological, cytological and genic aspects. *Scientific Reports*. 2016;6:22028. DOI: 10.1038/srep22028
- [109] Senesi N, Sposito G, Martin JP. Copper (II) and iron (III) complexation by humic acid-like polymers (melanins) from soil fungi. *Science of the Total Environment*. 1987;62:241-252
- [110] Fogarty RV, Tobin JM. Fungal melanins and their interaction with metals. *Enzyme and Microbial Technology*. 1996;19:311-317. DOI: 10.1016/0141-0229(96)00002-6
- [111] Bultreys A. Siderotyping, a tool to characterize, classify and identify fluorescent pseudomonads. In: Varma A, Chincholkar S, editors. *Microbial Siderophores*. New York: Springer; 2007. pp. 67-90. DOI: 10.1007/978-3-540-71160-5_3
- [112] Miethke M, Marahiel MA. Siderophore-based iron acquisition and pathogen control. *Microbiology and Molecular Biology Reviews*. 2007;71:413-451. DOI: 10.1128/MMBR.00012-07
- [113] Yamaji K, Watanabe Y, Masuya H, Shigeto A, Yui H, Haruma T. Root fungal endophytes enhance heavy-metal stress tolerance of *Clethra barbinervis* growing naturally at mining sites via growth enhancement, promotion of nutrient uptake and decrease of heavy-metal concentration. *PLoS One*. 2016;11:e0169089. DOI: 10.1371/journal.pone.0169089