We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Capsicum Seeds as a Source of Bioactive Compounds: Biological Properties, Extraction Systems, and Industrial Application

Javier Echave, Antía G. Pereira, María Carpena, Miguel Ángel Prieto and Jesús Simal-Gandara

Abstract

Recent research has substantially focused on residual subproducts containing chemical compounds with bioactive properties. Even though there are some culinary or medicinal uses of *Capsicum* seeds, there is still a seed mass waste from pepper processing. Many pepper leading producer countries generally lack the facilities and infrastructure required for such processing technologies and so, pepper seeds are usually either destroyed or employed as landfilling or as animal feed. This involves an inadvertent economic loss for producers as well as a detrimental environmental impact. However, there is a hidden potential within the pepper processing industry related to valorization of pepper seeds to obtain added value by-products and thus reduce generated waste. Pepper seeds are a good source of antioxidants, carotenoids, phenolic acids, flavonoids, and vitamins C, E, and A and are also rich in volatile compounds, among others. The unique alkaloids of this genus are capsaicinoids and capsinoids, which have been linked to many beneficial biochemical and pharmacological effects including anti-oxidative or anti-inflammatory activities. Other prominent bioactive compounds of peppers seeds include saponins, lectins, and polyunsaturated fatty acids. In this context, an overview of the biological properties, extraction systems, and possible industrial application of bioactive compounds of pepper whole fruit and seeds is presented.

Keywords: *Capsicum* spp., seeds, extraction, bioactive, activity

1. Introduction

1.1 Characteristics of the genre

The genus *Capsicum* belongs to the Solanaceae family, which consist of different variants of peppers that can be easily recognized by their size, shape, color and degree of pungency. This latter characteristic allows the classification of chili peppers depending on their Scoville heat units (SHU), a measurement of their pungency, into spicy or hot foods (generally those with small fruits) and sweet or

non-pungent (generally those with large fruits). This aspect is very interesting for marketing them and makes them a demanded product [1]. Considering this, we can consider for example the pungency of paprika peppers (10–30 parts per million capsaicinoids), chili peppers (30–600 parts per million) and red peppers (600–13,000 parts per million) [2]. Ripe fruits display a range of colors varying from white to deep red. Likewise, the intensity of red color and the degree of pungency are valued as major quality parameters. That is why the fruit is harvested when its red color has completely developed in order to ensure the highest quality of these features [3].

Capsicum is a very homogeneous genus with 33 species and 10 varieties (Table 1), with only five of them being domesticated: *Capsicum baccatum* L., *Capsicum pubescens* Ruiz & Pav., *Capsicum frutescens* L., *Capsicum chinense* Jacq. and *Capsicum. annuum* L. (Figure 1) native to Central and South America [5]. Each one of them was farmed independently in Pre-Columbians times in diverse regions of the American tropics [4]. They have been genetically modified to obtain varieties with agronomic interests, as in the case of *C. annuum* L. in which varieties more resistant to adverse factors were created, with a higher content and/or yield of the compounds of interest and fruit quality [6].

Amid the species *C. annuum*, there are different well know varieties like paprika, cayenne, jalapeños or chiltepin. Inside the specie *C. frutescens* are tabasco chilis; among *C. chinense*, the hottest chills (naga, habanero and Scotch bonnet) and among *C. pubescens* and *C. baccatum*, peppers emblematic from South America like rocoto or aji [7].

Although the characteristics of each species differ from each other (Table 2), it can be said that the genus *Capsicum* is characterized by being an annual or perennial herb or undershrub with entire or repand leaves that alternate. Its flowers are pedicelled, enclosing five petals; it is axillary and can appear solitary or in groups of two or three. Its sepals connate in a subentire or minutely five-toothed calyx that is much shorter than the fruit. It also has 5 stamens, which are adnate nearly to base of corolla-tube and are characterized for being short filaments. Its carpels connate in a two-celled (in some cases three) ovary, and its anthers are dehiscence longitudinal

Kingdom	Plantae
Subkingdom	Tracheobionta
Division	Magnoliophyta
Class	Magnoliopsida
Subclass	Asteridae
Order	Solanales
Family	Solanaceae
Genus	<i>Capsicum</i>
Species	<i>C. baccatum</i> , <i>C. pubescens</i> , <i>C. frutescens</i> , <i>C. chinense</i> , <i>C. annuum</i>
Varieties	<i>C. annuum</i> var. <i>annuum</i> , <i>C. annuum</i> var. <i>glabriusculum</i> , <i>C. baccatum</i> var. <i>pendulum</i>

Table 1.
Taxonomic classification of the genus *Capsicum* [1, 4].

Figure 1.
Some of the major cultivated pepper species. (A) *Capsicum frutescens*; (B) *Capsicum annuum*; (C) *Capsicum chinense*, and (D) *Capsicum baccatum*.

Species	Flower color	N° flower	Seed color	Calyx constriction	Distribution
<i>C. annuum</i>	White	1	Tan	Absent	Colombia, USA
<i>C. frutescens</i>	Green	2–5	Tan	Absent	No studies
<i>C. chinese</i>	White-green	2–5	Tan	Present	South America
<i>C. baccatum</i>	White with yellow spot	1–2	Tan	Absent	South America
<i>C. pubescens</i>	Purple	1–2	Black	Absent	South America

Table 2.
Characteristics that allow to distinguish between the most common species of the *Capsicum* genus [8].

without any exceeding filaments. As for fruit and seeds, the first is irregularly shaped (globose or elongate) with many seeded berries. Seeds are discoid, smooth or subscabrous [8].

Thus, its physical appearance defers from one species to another. For example, *Capsicum annuum* is an annual cultivate that reaches a height of 1 m and has glabrous or pubescent lanceolate leaves, white flowers, and fruit of varying length, color, and pungency depending upon the cultivar and growth conditions. It also the most widely cultivated pepper species around the world. Considering another species, *Capsicum frutescens* is a short-lived perennial with woody stems that reach

a height of 2 m, with glabrous or pubescent leaves, with two or more greenish-white flowers per node, and extremely pungent fruit [2].

Being a cold sensitive plant, the best conditions for production are between 7 and 29°C and an annual precipitation of 0.3–4.6 mm. It grows best in well-drained, sandy or silt-loam soil and a soil pH of 4.3–8.7. Hot and dry weather is also desirable for fruit ripening [2]. To carry out its cultivation it is necessary to seed or transplant the peppers, harvesting 3 months after planting [2].

1.2 *Capsicum* seeds sources and production

Chili (a variety of *C. annuum*) is one of the first plants cultivated in Mesoamerica, existing evidence of its use for the last 9000 years. It can be used in multiple ways: fresh, dry, as a spice powder, natural dye, antioxidant, bactericide and fungicide, as a drug in the pharmaceutical industry, in the cosmetology industry or in food industry (sausage, canned meats) [9]. These characteristics made them an essential part of daily cooking in many Latin American and Asian countries, for example, curry blends in India or in many meat sausages both fresh and dehydrated in the Mediterranean region [3].

On the basis of extracts obtained from pre-ceramic in the Coxcatlan caves, it is believed that the domestication of *C. annuum* probably occurred in the northeast or central-east of Mexico, being older than the remains of corn, beans and pumpkin [1]. In fact, peppers presently represent one of the vegetables of greater economic importance just behind the tomato for several American tropical countries [10].

Of the five domesticated species of *Capsicum* spp., the largest cultivated and with higher production per hectare are *C. annuum* and *C. frutescens* [11]. As for its world production, it is estimated that it is nearby 24.9 million tons (*C. annuum*), making it the ninth most produced vegetable in the world. Likewise, it has had an annual average growth of 6.26% in the last 10 years, with Mexico being its main consumer with an annual average of 8 kg/person. Its main producers are China (60.6%), Turkey (8.4%), Mexico (7.8%), Spain (5.0%), USA (4.3%), Indonesia (3.4%), Nigeria (2.8%), Egypt (2.2%), Korea (2.0%) and Italy (1.7%) [7]. In Mexico, one of the countries with the highest production levels, there are about a hundred cultivars of hot *Capsicum* spp. with varying degrees of spiciness, size, shape and colors. Nonetheless, there are also varieties that are cultivated throughout the world. This is the case of *C. annuum* or *C. chinense* [12]. Hence, it is a product that is traded internationally. This means that a series of elements such as aflatoxins, pesticides, residuals, microbial contaminations or infections, capsaicin levels and color values are subjected to inspection. All these aspects are controlled by law in several countries (EU, USA) [8].

Pepper quality depends on their composition, which is determined by factors such as environmental cultivation conditions, variety, ripeness, and pre-harvest and post-harvest handling and preservation [13]. The degree of ripening required may be one of the most important factors in quality, but it will also depend on the destined market, since not the same degree of maturity is desired for all the possible uses. Notably, the moment of harvesting is also important for the maintenance of the quality as metabolic activity persists after harvesting [14].

1.3 Chemical composition

Regardless of the enormous consumption and production of this kind of vegetables, there is little data about the chemical composition of the different varieties (Table 3). However, the demand and cultivation of peppers, especially “hot” cultivars, has increased due to its flavoring and medicinal properties. Some of the latter have been described as anticancer, antioxidant and antimicrobial. Its edible and

Common name	State	Carbohydrates	Protein	Fat	Capsaicinoids	Fiber	Ash
Guajillo	D	58.00	12.89	12.43	5.97	Nd	7.52
Ancho	D	60.21	12.05	9.82	8.50	Nd	7.81
Pasado	D	66.18	12.61	5.41	9.74	Nd	7.18
Pasilla	D	60.53	12.28	13.76	11.80	Nd	5.85
Puya	D	63.76	13.25	8.11	12.13	Nd	7.82
M. Tres venas	D	61.05	13.28	9.61	14.40	Nd	7.02
Chiplote Meco	D	57.68	15.22	9.08	29.01	Nd	9.54
Jalapeno	D	63.97	14.36	4.23	58.40	Nd	7.32
Mirasol	D	58.96	14.05	7.49	58.55	Nd	9.61
Morita	D	58.91	14.12	7.60	67.32	Nd	8.59
Serrano	D	67.93	12.78	2.26	102.73	Nd	5.81
Chiplote	D	62.92	12.72	8.66	143.57	Nd	6.92
De Arbol	D	59.41	12.75	13.38	193.51	Nd	8.82
Piquin	D	62.25	13.72	11.02	368.83	Nd	7.28
Habanero	D	61.13	13.52	4.63	1312.10	Nd	7.51
Marako fana	D	35.3	11.8	11.2	Nd	27.3	5.3
Bako local	D	39.5	8.7	9.5	Nd	26.0	7.3
Oda haro	D	37.1	9.2	9.2	Nd	28.6	7.3
Arnoia red	F	6.23	0.15	0.54	Nd	1.62	0.62
Arnoia green	F	3.84	0.14	0.22	Nd	1.63	0.40
Arnoia green B	F	3.51	0.12	0.16	Nd	1.31	0.33
Hot pepper	D	Nd	21.29	23.65	Nd	38.76	4.94
Chunhamuchuk	D	Nd	15.05	29.27	Nd	48.72	3.59
Amhanegosa	D	Nd	14.66	26.70	Nd	52.10	3.28
Hanbando	D	Nd	14.08	27.84	Nd	38.43	3.49
Dachon I	D	Nd	15.99	19.53	Nd	50.61	3.76
Samgang	D	Nd	13.90	23.50	Nd	50.71	3.71
Chunhajeil	D	Nd	14.67	21.87	Nd	52.54	3.46
Daejangbu	D	Nd	14.88	26.50	Nd	53.78	3.47
Hongjangkun	D	Nd	15.17	21.61	Nd	54.66	3.33
Kumbit	D	Nd	15.09	23.28	Nd	46.17	3.18
Dokyachungjung	D	Nd	15.75	25.13	Nd	53.36	3.86
Dangchan	D	Nd	15.36	19.99	Nd	55.63	3.46
Chohyang	D	Nd	13.25	18.05	Nd	59.13	3.77
Taesan	D	Nd	14.71	23.45	Nd	48.80	3.11
Ganggun	D	Nd	15.55	20.63	Nd	52.71	3.43
Chungsan	D	Nd	15.06	20.45	Nd	50.75	3.05
Dachon II	D	Nd	15.89	18.83	Nd	45.73	3.61
Wangdaebak	D	Nd	16.53	23.65	Nd	54.39	3.28
Chunhailpum	D	Nd	15.70	19.79	Nd	53.34	3.38

Common name	State	Carbohydrates	Protein	Fat	Capsaicinoids	Fiber	Ash
Daechan	D	Nd	13.88	20.64	Nd	54.68	3.72
Mixed	D	Nd	14.01	24.09	Nd	50.26	3.26
Sandia	D	Nd	14.95	23.07	Nd	58.34	3.22
R-Naky	D	Nd	14.36	23.57	Nd	60.19	3.57
New Mexico 6	D	Nd	14.79	21.95	Nd	60.61	3.48
LB-25	D	Nd	14.87	25.06	Nd	52.98	3.29

F, fresh; D, dried; Nd, not described.

Table 3.
Different Capsicum spp. and their proximate chemical composition (g/100 g) [14–18].

nutritional value is acknowledge as well, since it is rich in vitamins (A, C, B₆, E), carotenoids (β-carotene), flavonoids, oils, oleoresins and alkaloids [19]. Therefore, the compounds that can be found in this genre are carbohydrates (accounting for approximately 85% of dry weight), polyphenols (0.5% of dry weight) and important molecules such as capsaicinoids, carotenoids and vitamins [20]. Given these facts, peppers are considered a good source of most essential nutrients [14].

The capsaicinoids content, which depends on the variety and maturation stage, will determinate the pungency. *C. chinense* and *Capsicum annuum* var. *aviculare* contain larger amounts of capsaicin and dihydrocapsaicin (ratio of 2:1), while some varieties of *Capsicum annuum* var. *annuum* showed an average proportion of 1:1. These variations could be attributed to environment, genetics and extraction methodologies [15]. Even though in some cases peppers are sought to be spicy, this can also be a limitation. This is the case, for example, in obtaining dyes from this raw material in the food industry. In this case, extraction methods need to be improved to prepare non-pungent oleoresins from pungent *Capsicum* fruits. This is achieved by the selective removal of capsaicinoids which allows the exploration of a large number of pungent varieties with good oleoresin yielding [20].

Capsaicinoids are the characteristic pungent compounds of the *Capsicum* genus. They include capsaicin, dihydrocapsaicin, nordihydrocapsaicin, homocapsaicin, and homodihydrocapsaicin. Capsaicin is the most abundant. As an example, capsaicin constitutes together with dihydrocapsaicin approximately 90% of total capsaicinoids content of chili peppers [15].

The characteristic red color of many peppers is determined by the presence of different carotenoids. Actually, more than 50 different carotenoids can be found in this kind of material. Some of them are capsanthin, capsorubin, and cryptocapsin which give brilliant red color (ripe fruits) or β-carotene, violaxanthin, zeaxanthin and β-cryptoxanthin which give yellow–orange color [3]. Nevertheless, the color will depend on the state of maturity. For example, jalapeño (*C. annuum*) has a green color, and when it is mature it presents an intense red color [7]. Therefore, depending on the degree of maturity, the physicochemical parameters, flavor and mineral composition differs. For example, fat, protein, ascorbic acid (vitamin C), soluble solid content and titratable acidity will increase during ripening [14].

Using jalapeño as an example to study composition, its main component is water, then carbohydrates (5.3%), fiber (2.3%), protein (1.2%), fat (0.1%) and minerals, being the most important potassium (340 mg per 100 g of fresh product). It also has calcium (25 mg per 100 g of fresh product), magnesium (25 mg per 100 g of fresh product), sodium (7 mg per 100 g of fresh product), iron (2 mg per 100 g of fresh product) and zinc (0.3 mg per 100 g of fresh product). As for vitamins, the most important are ascorbic acid, retinol and folic acid (72, 20 and 23 mg per 100 g

of fresh product, respectively). Other vitamins that can be found in peppers are thiamine, riboflavin, niacin and pyridoxine. Moreover, jalapeños also contain important amino acids such as lysine, methionine and valine (252, 40 and 23 mg per 100 g of protein, respectively) [7].

2. *Capsicum* seeds as a source of bioactive compounds

Like many fruits and plants, peppers are an excellent nutritional source. As aforementioned, it has an abundance of minerals, vitamins, aminoacids, carotenoids as also phytochemicals like phenolic compounds or polyunsaturated fatty acids (PUFAs). Likewise, capsaicins are unique to the genus *Capsicum*, being responsible of the pungency of many pepper species. Capsaicinoids have demonstrated to induce a vast range of bioactivities, such as anti-inflammatory, anticancer analgesic, antimicrobial, hypotensive and induce lower adipogenesis or lower body temperature by isolated action or in synergy with other compounds [21–25].

Different bioactive compounds have been isolated and extracted from *Capsicum* spp. fruits and seeds and its concentrations vary among species. More information on reported compounds studied may be found in (Table 4).

2.1 Phenolic compounds

Phenolic compounds, also referred as phenolics, are secondary metabolites that may be found in a wide spectrum of plant species. They are synthesized as a result of adaptation to biotic and abiotic stress through the phenylpropanoid pathway playing an important role in plant development, because they act as a defensive mechanism that eases plant growth against harsh conditions [33, 34]. Based on recorded knowledge, phenolic compounds exhibit numerous potential health benefits that are already well described in scientific literature and are currently a major current focus of nutritional and pharmacological research [35, 36]. Phenolic acids and flavonoids are the main phenolic phytochemicals found in peppers. Likewise, capsaicinoids are synthesized in the same biochemical pathway and exhibit some similar properties such as antioxidative activity [34].

The yield of phenolic compounds recovered from an extraction can be very different, depending largely on the extraction method, the conservation of vegetal material and maturity state [37]. Generally, fresh raw material preserves the highest quantities of phenolic compounds [16]. Several factors contribute to yielding disparity, such as heterogeneous genotypes, growing and harvesting conditions of the samples [38]. Some of the most prominent non-capsaicinoid phenolic compounds because of their valuable health benefits are phenolic acids like gallic acid, caffeic acid, chlorogenic acid or ellagic acid (Figure 2). Flavones are another important group of phenolic compounds, being some of the most prominent kaempferol, quercetin, luteolin or rutin (Figure 3) [26]. These phenolics have demonstrated great health benefits and many of them are commercially available in purified products extracted from other plant species [39, 40]. Even so, their concentration differs among species but not much among varieties [26]. Other phenolic compounds are reviewed in the next paragraphs as in the case of some carotenoids or vitamins.

Regarding capsaicinoids, they are synthesized naturally in the placenta of pepper fruits by enzymatic transformation of vanillylamine, the phenolic portion of the molecule, which confers this alkaloid its antioxidant capacity [41]. The seeds are not the primary source of capsaicinoids but they may absorb them because they are in close proximity to the placenta, which is the richest capsaicin fraction [42]. Their presence in the seed and the high concentrations they achieve, has been observed to

Species and varieties	Bioactive compounds						
	Major phenolic compounds	Vitamins	Pigments	Volatiles	PUFAs	Peptides	Ref.
<i>C. annuum</i>	4-Hydroxybenzoic acid, vanillic acid, caffeic acid, <i>p</i> -coumaric acid, sinapic acid, ferulic acid, ellagic acid, naringein	Nd	Nd	Nd	Nd	Nd	[26]
<i>C. annuum</i>	Elagic acid, gallic acid, chlorogenic acid, caffeic acid, salicylic acid, rutin	Nd	Nd	Nd	Nd	Nd	[21]
<i>C. annuum</i>	Caffeic acid, gallic acid, ferulic acid, rutin, capsaicin, dihydrocapsaicin	Tocopherols	Capsanthin, zeaxanthin, capsorubin, lutein, β -carotene, cryptocapsin	Nd	Linoleic, palmitic, oleic acids	Nd	[27]
<i>C. annuum</i>	Nd	Nd	Nd	Nd	Nd	Mannose/ glucose Specific lectin	[28]
<i>C. annuum</i>	Nd	Ascorbic acid	Nd	Betulin, β -sitosterol	Oleic, palmitic, linoleic, elaidic acids	Nd	[29]
<i>C. annuum</i>	Gallic acid, caffeic acid, ferulic acid, chlorogenic acid, catechin, epicatechin, capsaicin, dihydrocapsaicin	Ascorbic acid	Nd	Nd	Nd	Nd	[30]
<i>C. annuum</i> var. <i>annuum</i>	Caffeic acid, cinnamic acid, coumaric acid, capsaicin, vanillin	Nd	Nd	Nd	Nd	Phenyl-alanine	[31]
<i>C. baccatum</i>	Tannins, flavonoids, coumarins, polyphenols	Nd	Nd	Saponins	Nd	Amino acids	[32]

Nd, not described.

Table 4.
Capsicum seeds bioactive compounds.

rise the riper the pepper is [43]. It is also confirmed that fresh seeds yield more capsaicin than dry seeds, which suggests that surface capsaicinoids are sensitive to heat and/or oxidation [29]. “Hot” pepper cultivars attribute their pungency to high levels of capsaicinoids whereas non-pungent or “sweet” peppers (e.g., bell pepper) have very low capsaicinoids quantities [44].

Capsinoids are other non-pungent capsaicin analogues noticeably found in *C. baccatum* var. *praetermissum* and the sweet cultivar CH-19 of *C. annuum* [45, 46].

Figure 2.
Chemical structure of some main phenolic acids found in peppers: (A) gallic acid; (B) ellagic acid; (C) caffeic acid and (D) chlorogenic acid.

Figure 3.
Chemical structure of the main flavonoids found in pepper: (A) luteolin; (B) quercetin; (C) kaempferol; and (D) rutin.

Capsinoids exhibit biological activities similar to capsaicin due to being capsaicinoids analogs, except that they display much lower pungency in comparison [45]. Capsaicinoids and capsinoids manifest many promising therapeutic properties such as apoptosis induction, antioxidation, anticancer (cytostatic), or immunomodulation, making them attractive targets for pharmaceutical research [24, 47].

Besides, it is worth mentioning that the well-known flavone chrysoeriol (**Figure 4**), present in a multitude of vegetables and at least in *C. annuum* and *C. frutescens* fruit and seeds, has been determined to inhibit the growth of several microorganisms at very low quantities [31].

Figure 4.
Chemical structure of chrysoeriol.

2.2 Fatty acids

Peppers are fruits rich in polyunsaturated fatty acids (PUFAs), and their seeds show even greater concentration per gram [42]. The main fatty acids are indeed PUFAs [48]. These PUFAs, which are in the whole fruit and seeds, are mainly linoleic acid, palmitic acid α -linolenic acid and stearic acid [16, 48]. Moreover, peppers appear to have very low levels of saturated fatty acids. Linoleic acid has the highest concentration ($\approx 70\%$) and the other fatty acids show much lower levels [48, 49]. PUFAs and specifically linoleic acid and α -linolenic acid, are recognized as essential fatty acids and are precursors of other important fatty acids in metabolism like arachidonic acid and eicosapentanoic acid or prostaglandins that in whole contribute to normal physiological performance [50].

2.3 Pigments

Chlorophylls and carotenoids constitute another group of valuable pepper nutrients that affect its color, but they also have important antioxidative, anti-inflammatory effects and promote immune response [22, 51, 52]. Zeaxanthin, β -carotene, violaxanthin, lutein and β -cryptoxanthin are the pigments with highest concentration among *Capsicum* spp. [37, 43]. Capsanthin is another red colored pigment mostly present in red bell pepper (*C. annuum*) [53]. They all show strong antioxidative properties and increase their levels the more mature the pepper is [44].

β -carotene and β -cryptoxanthin for instance, possess the added value of being able to be converted to vitamin A [54]. Vitamin A plays a vital role in disease prevention and development [54].

2.4 Vitamins

As it has been mentioned, peppers contain several vitamins like ascorbic acid and tocopherols like α -tocopherol, γ -tocopherol and δ -tocopherol, are all isomers of vitamin E [55, 56]. These essential vitamins contribute to the normal metabolism, promote immune response and also have antioxidative bioactivity [52, 57]. Because of this, ascorbic acid is an essential vitamin that is also used as a natural food preserver [52]. Indeed, peppers contain levels of vitamin C corresponding with those found in many citrus fruits and other vegetables considered good sources of this vitamin [14]. Tocopherols are well-known lipophilic antioxidants and appear to exert a vital part in the normal T lymphocyte maturation, lower age-related increase in tissue inflammation and lower interleukin production [55, 58]. This makes vitamin E an important modulator to an orderly and better inflammatory reaction, among other health benefits [58].

2.5 Volatile compounds

Volatile compounds in *Capsicum* spp. have been studied because of the importance they have in many plants as pollinator attractors and their influence in odor and flavor, which are key features of any edible vegetable [59, 60]. Some of the most valuable are hexyl isopentanoate, hexyl isobutanoate, and β -ionone [61]. Extracts of *C. annuum* have also showed high levels of 4-methoxyphenol, ethylhexadecanoate, hexanal or isopulegol [62].

2.6 Other minor compounds

There are a few other minor and less studied compounds amid the different pepper species, because research has focused almost entirely on finding and studying phytochemicals of metabolic importance. In regard to these bioactive compounds, *C. annuum* or *C. frutescens* saponins have showed antimicrobial activity against several opportunistic and pathogenic fungi genres [63]. Saponins are triterpene glycosides with the ability to disrupt the cell membrane and wall of fungal cells during cell proliferation, which leads to cell lysis [27]. As such, saponins do not have effect on non-germinated fungi, but they inhibit its growth. Although there are several present in pepper tissue, CAY-1 has been well-studied (**Figure 5**), presenting a high antifungal activity [60].

Another of these minor chemicals are lectins, a group of proteins with glycoside agglutination properties. Some lectins have been described in *C. frutescens* and *C. annuum* seeds and have been studied as possible antimicrobials and have yielded positive results against some fungi [28, 64].

Figure 5.
Chemical structure of CAY-1. Adapted from [60].

3. Bioactivities of the raw material

In recent years, nutraceutical and therapeutic research has focused its view towards both exotic and domestic fruits and vegetables as a source of phytochemicals with the ability to induce beneficial bioactivities [65]. These natural chemicals have been and are already used as a main source of therapeutics in traditional and modern medicine, reaching one third of the total production of therapeutics [66]. Furthermore, processing waste by-products of fruits and vegetables are being researched as a viable source of phytochemicals that would be affordable and reduce economic and ecological impacts of wasted by-products or taking produce out of the food market [67, 68]. Thus, fractions that are not employed in nourishment such as the placenta, seeds or leaves of many species can prove to be a valuable resource instead of end as waste or fertilizer, which is the most common use for vegetable non-edible parts [61].

Taking into account the aforementioned compounds present in the many different species of the genus *Capsicum* and different regional cultivars, as in the case of *C. annuum*, many biological activities are expected to be found in direct or indirect physiological response to the presence of these substances. It is widely known that capsaicinoids, capsinoids, saponins, many phenolic compounds and its valuable nutrients are behind the beneficial effects of this vegetable, which has been traditionally employed as a source of medicinal remedies all across South America, Asia and Africa [63]. The intensity and type of activity may vary depending on the concentration of these compounds and will differ in each species and cultivar of *Capsicum*, the bioavailability of each compound, the ripening state and from which fraction of the vegetable the extracts are obtained [43, 44].

These bioactivities have been described in scientific reports through tissue culture and both animal and human test research as antioxidant, antimicrobial, anti-inflammatory, anticancer, analgesic or even antidiabetic [21, 22, 34]. Considering this with the fact that *Capsicum* spp. is widely grown around the world, it makes it to be considered an excellent source of new therapeutics and dietary supplements that could lead to a healthier well-being. Indeed, many extracts of pepper fruits and seeds with a varying purification degree are marketed and can be purchased [69].

Some of the most recognized bioactivities found by chemicals in the whole pepper and its seeds will be reviewed and a brief compendium of the bioactivity research can be found in **Table 5**.

3.1 Antioxidant

Oxidative stress is caused as a result of the presence of reactive oxygen species (ROS) which may be produced in oxidative metabolism and exposure to the environment [33]. The term ROS englobes the molecules superoxide radical (O_2^{2-}), the hydroxyl radical ($OH^\cdot$) and hydrogen peroxide (H_2O_2). They are produced by the sequential reduction of molecular oxygen in various metabolic reactions [76]. O_2^{2-} is the most unstable form but it may dismutate to H_2O_2 by the action of endogenous superoxide dismutase (SOD) enzyme reaction or non-enzymatically [76]. The effect of oxidative stress mainly translates into changes in the rate of metabolic reduction reactions and an increased rate of DNA mutations and cell mitosis. They are also main signals indicating cell death, which in turn triggers inflammation via the pro-inflammatory factor nitric oxide (NO), which is released by macrophages [33, 76]. Since these are both cause and result of cancer development, it is considered that high levels of ROS are detrimental to health [40, 65, 76]. All living aerobic

Activity	Bioactives	Species	Type of study	Test results	Ref.
Antioxidant	Capsaicin, dihydrocapsaicin	<i>C. annuum</i> , <i>C. baccatum</i> var. <i>pendulum</i> , <i>C. chinense</i> , <i>C. frutescens</i>	In vitro	DPPH, ABTS	[68]
	Total phenolic compounds	<i>C. annuum</i>	In vitro	DPPH, ABTS, ORAC	[26]
	Phenolic extracts	<i>C. annuum</i>	In vitro	DPPH, ferrus chelating	[39]
	Seed oil	<i>C. annuum</i>	In vitro	DPPH, ABTS	[42]
Antimicrobial	Capsicosides A, G, D	<i>C. annuum</i> var. <i>acuminatum</i>	In vitro	Gia. various yeasts	[32]
	Capsaicin	<i>C. annuum</i> , <i>C. baccatum</i> , <i>C. chinense</i> , <i>C. frutescens</i>	In vitro	Gia. <i>Clostridium tetani</i> & <i>C. sporogenes</i>	[31]
	Phenolic extracts	<i>C. annuum</i>	In vitro	Gia. <i>S. aureus</i> , <i>L. monocytogenes</i> , <i>S. typhimurium</i> , <i>B. cereus</i>	[70]
	CAY-1	<i>C. frutescens</i>	In vitro	Gia. <i>A. flavus</i> , <i>A. fumigatus</i> , <i>A. parasiticus</i> , <i>A. niger</i> , <i>C. albicans</i> , <i>P. carinii</i>	[60]
	Lectins	<i>C. annum</i>	In vitro	Gia. <i>A. flavus</i> , <i>F. graminearum</i>	[28]
	Total polyphenol content, capsaicins	<i>C. annuum</i>	In vitro	Gia. <i>E. faecalis</i> , <i>P. aeruginosa</i> , <i>E. coli</i> , <i>S. aureus</i>	[21]
	Jalapeño extracts	<i>C. annuum</i> var. <i>annuum</i>	In vitro	Gia. <i>L. monocytogenes</i>	[71]
	Capsaicinoids, chrysoeriol	<i>C. frutescens</i>	In vitro	Gia <i>E. faecalis</i> , <i>B. subtilis</i> , <i>S. aureus</i> , <i>P. aeruginosa</i> , <i>E. coli</i> , <i>C. albicans</i>	[31]
	Residual aqueous extract	<i>C. baccatum</i>	In vitro	Gia <i>S. epidermidis</i> , <i>P. aeruginosa</i> biofilm	[72]
Anti-inflammatory	Capsaicin	Nd	In vitro	Inhibition of inflammatory transcription factor NF-κB and AP-1	[73]
	Capsaicin	Nd	In vitro	Inhibition of Iκα and Iκβ via NF-κB	[74]
	Capsaicin	Nd	In vitro	Inhibition of adipose tissue inflammation (interleukin 8, c-Jun)	[75]

Nd, not described; Gia., growth inhibition against; DPPH, 2,2-diphenyl-1picrylhydrazyl scavenging method; ABTS, 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt scavenging methods; ORAC, oxygen radical absorbance capacity.

Table 5.
Properties and mechanisms of bioactive compounds and Capsicum spp. extracts evaluation.

organisms have developed defense mechanisms against oxidative stress through the synthesis of reductive biochemicals or enzymes such as SOD. The most important antioxidative biochemicals are phenolic compounds, which are prominently found

in plants [52, 66]. These antioxidants present in peppers, mainly identified as the mentioned phenolic compounds, vitamins and pigments, have the potential to reduce biological oxidative stress and thus preventing the incidence of many related diseases, but also to further food preservation by inhibiting the oxidative metabolism of decomposing fungi and bacteria [33, 35, 38]. Phenolic compounds are the principal antioxidants in nature and generally show the greatest antioxidative capacities [65]. However, high concentrations of few phenolic compounds showed pro-oxidant effects due to a concentration imbalance between ROS and the phenols, which is why it is important to maintain intake of different phenolic compounds in order to benefit from their antioxidant properties [52]. Thus, antioxidants have a wide extent of applications, and many natural and synthetic antioxidants have been used by the food industry in order to better preserve raw or precooked products that would otherwise have a much shorter shelf-life [22].

The antioxidative effect of the aforementioned phenolic compounds essentially works by scavenging free superoxide and hydroxyl radicals and thus preventing high levels of ROS, NO and oxidation of sensitive biomolecules like proteins or lipids (**Figure 6**) [36, 40]. This results in a better physiological performance, immunomodulation and DNA mutation protection. Furthermore, the decrease of oxidative stress results in better cardiovascular health. This is due to the fact that low-density lipoprotein (LDL), as the main cholesterol carrier in the circulatory system, is susceptible to oxidation by ROS [58]. Oxidized LDL presence has been found to be cause of atherosclerosis, a vascular ailment caused by the formation of plaques inside the arteries that may weaken blood flow and lead to cardiovascular diseases [77]. On top of that, pepper and its seeds are described to be one of the richest vegetables in phenolic compounds [29]. Then, it should be taken into account the antioxidant properties of the many phenolic acids, capsaicinoids, vitamins and pigments present among the *Capsicum* genus and the fact that these exploitable compounds are available not only in the fruit, but also in the non-edible fractions [39, 57, 78].

3.2 Antimicrobial

Pepper seed extracts and selected pigments, phenols, capsaicins and capsiates have been tested against the most common microorganisms present in foods and/or potential pathogens. The major antimicrobial effect of capsaicinoids has been found to be against common opportunistic and pathogenic fungi like *Fusarium* spp., *Aspergillus* spp. and yeasts like *Saccharomyces cerevisiae* or *Candida albicans*, but also against certain bacteria like *Streptococcus* spp., *Helicobacter pylori*, *Pseudomonas aeruginosa* or

Figure 6. Scheme of radical scavenging performed by gallic acid as an example of phenolic compounds. Adapted from [36].

Listeria monocytogenes [21, 32, 38, 79]. It has been reported that it can also reduce the virulence of pathogenic infections by *Vibrio cholerae* or *L. monocytogenes*, by inhibiting pro-inflammatory factors or toxin production [79–81]. There are few *in vivo* studies of these effects, though there are some reports of poultry showing less hepatic damage from *Salmonella enteritidis* when given feed supplemented with capsaicinoids, which could be related to its antioxidative activity [24].

Capsaicinoids appear to inhibit, fungi, and both Gram positive and negative bacterial growth in several studies. In all of them, capsaicin and/or dihydrocapsaicin showed to stop or slow colony development in a significant degree [24, 35]. Some interesting results of antimicrobial tests with capsaicinoids reported that it inhibits the growth of *Escherichia coli*, *L. monocytogenes*, *Bacillus subtilis* and many strains of *Staphylococcus aureus* [21].

Even though capsaicinoids possess these interesting antimicrobial properties, the positive correlation between antimicrobial activities, concentration and its pungency and potential irritation effects make them less suitable when it comes to applying this compound as a food preserver [46]. Thus, it is interesting to study the effects on this area of alternative chemicals found in pepper fruit and seeds. CAY-1, chrysoeriol and lectins have been previously mentioned as such.

CAY-1 is found in *C. frutescens* and a study demonstrated to produce cell membrane lysis in fungi like *Aspergillus flavus*, *Aspergillus niger*, *Fusarium solani*, *Fusarium moniliforme*, *Pneumocystis carinii* or *Candida albicans* [60]. It is ineffective though against bacteria though, probably because of the different type of cell membrane and wall [60]. It has also been suggested to combine this saponin with certain metallic elements such as silver in order to produce antimicrobial surfactants [82]. Other saponins isolated from *C. annuum* seeds, like capsicosides, which are furostanol saponins, have showed similar effects on growth inhibition against yeasts [27].

Chrysoeriol is another potent antimicrobial compound present in a variety of plants as well as in peppers [83]. Several studies have presented that chrysoeriol is capable of greatly inhibit the growth of *Enterococcus faecalis*, *Klebsiella pneumoniae*, *S. aureus*, *L. monocytogenes*, *B. subtilis* or *C. albicans* [31, 84]. Given the very low concentrations at which it exerts its antimicrobial effect, it may be an optimal food preserver against many opportunistic and nosocomial microbes.

Certain lectins isolated from peppers have showed antifungal properties, inhibiting growth of some common opportunistic fungi like *Aspergillus flavus* and *Fusarium moniliforme* by limiting surface adherence. However, its inhibitory action is not as effective across all species of these genres [28, 85].

Furthermore, a recent study analyzed the antiadhesive capacity of natural peptides from *C. baccatum* var. *pendulum* on biofilms made by *Staphylococcus epidermidis* [64]. Considering that biofilms are a source of significant virulence, new studies on this field present extremely interesting. Altogether, these findings may contribute to develop new uses for natural antimicrobials, which would be an application of vital priority, given the global concern on the appearance of new-born antibiotic resistant microorganisms [31, 84].

3.3 Anti-inflammatory

Although an anti-inflammatory action is carried indirectly by the antioxidant capacity of other compounds to scavenge ROS and inhibit NO production, it is known that capsaicins and capsiates directly induce an anti-inflammatory effect through the activation of the transient receptor potential vanilloid 1 receptor (TRPV1) [24, 86]. TRPV1 is an ion channel receptor located in several glia that gives sensations of heat, but is also relevant in pain perception and induces this sensation,

as well as inflammation [87]. By activating this receptor, capsaicins cause the well-known pungency sensation, as well as irritation in high enough concentrations. However, after activation by capsaicinoids, the excited neurons become resistant to further stimuli [87]. Hence, after a brief burning sensation, capsaicin may act as a local analgesic in neuropathic pain [84].

Research on capsaicinoids has showed that these molecules can also decrease inflammation of adipose tissue linked to obesity [73, 75]. It seems that capsaicin is able to inhibit or at least partially decrease the production of pro-inflammatory signals like interleukin 8 (IL-8), nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) or active protein 1 (AP-1) [74, 80]. This anti-inflammatory effect has the potential to make inflamed tissue less prone to tumor development as well as reduce the infection caused inflammation [25].

Still, the pungency of capsaicins poses a setback for using them without potentially hazardous secondary effects. That is why capsinoids may be a good alternative to induce similar effects to capsaicin, even though they are not found in so many pepper species [41, 45].

Although the importance of researching capsaicinoids bioactivities in pepper is due to its unique presence in this genre, the also present phenolic compounds, vitamins and pigments also bear significant anti-inflammatory properties [25, 88, 89]. Indeed, the antioxidative properties of phenolic compounds could be a key factor in reducing age-related extended tissue inflammation [25, 58]. Nonetheless, further research assessing the suitability of using as anti-inflammatory drug these phytochemicals, and specifically capsaicin, is needed.

4. Extraction systems for bioactive compounds from *Capsicum* spp.

The bioactive compounds obtained from nature are secondary metabolites produced by the organisms. The concentrations of each type of compound are very variable, so in order to have enough quantities, the development of new and advanced technologies is needed.

Although there are several extraction methodologies, there is a demand for more appropriate and standardized extraction strategies. For choosing one method over another, it must be taken into account the quantitative and qualitative characteristics of the compound of interest. In addition, to improve the efficiency of a method it is important that the nature of the source, the different parameters of the method and the possible interaction are taken into account. Experiments carried out with peppers can be seen in **Table 6**.

4.1 Conventional extraction systems

4.1.1 Maceration

Maceration (MA) is a type of solid-liquid extraction, in which the solid would be peppers. This raw material is characterized by having several compounds soluble in the liquid phase. The type of molecules extracted will depend on two factors: the type of starting material and the type of solvent used.

In order to carry out this process, the sample is suspended in the solvent at the desired temperature during the chosen time while stirring. Once the process has been carried out, the sample is centrifuged and the supernatant, which is where the compounds of interest are found, is filtered. Therefore, the process of extraction consists of two stages. On the first (washing) there is a rapid transfer of the compounds from the surface of the solid to the solvent. On the second one (transfer), the matter passes from the interior to the exterior of the solid by diffusion

Compound	Species	Extraction		Medium	Yield	Ref.
		Type	Method			
Pigments	<i>Capsicum</i> spp.	Emergent	SFE	—	6%	[90]
	<i>C. annuum</i>	Green	MAE	Acetone-water	—	[91]
		Emergent	SFE	—	17.4%	[92]
				—	93%	[93]
				—	7.2%	[94]
Carotenoids	<i>C. annuum</i>	Green	EAE	Viscozyme L	87%	[95]
				Pectinase	80%	[95]
				Cellulase	Low	[95]
				Viscozyme L	78%	[96]
Phenolics	<i>C. annuum</i>	Emergent	SFE	—	84%	[93]
				—	100%	[94]
Capsaicinoids	<i>Capsicum</i> spp.	Green	MAE	Acetone (30%)	0.48 mg/g	[97]
	<i>C. annuum</i>	Green	MAE	None	230 ppm	[98]
			UAE	None	200 ppm	[97]
				n-hexane (100%)	—	[99]
				Methanol (100%)	100%	[100]
	<i>C. annuum</i>	Green	EAE	Viscozyme L	88.8%	[96]
				Viscozyme L	22%	[101]
				Celluclast	20%	[101]
		Emergent	SFE	—	8.6%	[92]
				—	—	[102]
				—	93%	[93]
				—	710 µg/g	[103]
	<i>C. frutescens</i>	Green	MAE	Acetone (pure)	5.3 mg/g	[104]
				Ethanol (99.5%)	—	[105]
		Emergent	SFE	—	710 µg/g	[103]
				—	5.2%	[108]
	<i>C. chinense</i>	Green	UAE	Acetone (100%)	0.31 mg/g	[109]
				Methanol (100%)	2.88 mg/g	[110]
		Emergent	SFE	—	0.5%	[111]
	<i>C. baccatum</i>	Green	UAE	Methanol (100%)	50%	[112]
Antioxidants	<i>C. baccatum</i>	Green	UAE	Methanol (100%)	27%	[112]
OLEORESIN	<i>C. baccatum</i>	Green	UAE	Methanol (100%)	26%	[112]
	<i>C. annuum</i>	Green	EAE	Viscozyme L	6%	[101]
		Emergent	SFE	—	8.2%	[102]
				—	7.4%	[94]
				—	—	[113]
	<i>C. frutescens</i>	Emergent	SFE	—	0.3%	[108]
Other phytochemicals	<i>Capsicum</i> spp.	Green	UAE	Methanol (80%)	—	[114]

Table 6.
Bioactive compounds obtained from Capsicum spp. extracted with different extraction methods.

being this step the limiting stage. This method is used, for example, for the isolation of phenolic compounds of vegetable origin for their subsequent use in the food, pharmaceutical and cosmetic industries [115].

Water can be used as an extraction solvent, but it comes with the drawback is that it principally extracts the hydrophilic compounds present in vegetable materials. Several studies showed that water is not a very suitable solvent because its high polarity does not allow the extraction of capsaicinoids that are non-polar. However, it has the advantages of being safe, cheap and the simplest form to obtain essential oils. This technique can also be done with other solvents, being the ratio of solvent another parameter to take into account. Polarity is the reason why the most common solvents used are methanol, ethanol, water or a mixture of them [3].

Another parameter to optimize will be temperature. This makes it possible to differentiate two types of MA: cold maceration and heat maceration. Maceration with or without stirring, mild heating or heating under reflux are also possible variations to the method.

It is a quite old simple method with the inconvenience of long extraction times and large amounts of sample and solvents. Additionally, high temperatures can destroy thermolabile compounds such as phenolic compounds. In a study in which the effect of MA and ultrasound was related, it was detected that they have similar extraction yield of oleoresin; nevertheless, MA needs longer extraction times with the consequent loss of quality due to their higher times of exposure to high temperatures. It also was observed that n-hexane was a better solvent than ethanol [116].

4.1.2 Heat assisted extraction

This procedure is also a type of solid-liquid extraction in which the extraction is usually carried out in thermostatic and sealed water baths so that the solvent does not evaporate. The determining parameters are extraction time, temperature and the proportion of solvent that is used as well as the solid-liquid ratio. Due to the application of high temperatures (as far as 100°C), it is not the best extraction process for this type of matrix because certain compounds present in peppers (vitamins, phenolic compounds, etc.) are degraded by heat.

4.1.3 Cold pressure extraction

Cold pressure extraction is one of the oldest techniques of extraction for obtaining oils. It consists of mechanical pressing with the absence of heating. By using this method, little to no heat is generated, however it gives low yields. This technique was applied to Capiá pepper seed, resulting in a lower extraction yield compared to traditional Soxhlet extraction with hexane. Moreover, the final oil was unpopular among consumers [117].

4.1.4 Organic solvent extraction

Organic solvent extraction (OSE) allows the extraction of many compounds (oils, fats and proteins). Normally, after the extraction process another step is done. It consists of concentrating the extract by removing the solvent under atmospheric or reduced pressure. OSE is the most extensively used technique to obtain oleoresins from peppers. These oleoresins are. Usually used as color additives [118]. In this method, the most determining limitation will be the solvent properties (polarity). When choosing the solvent, it must also be taken into account if the legislation allows its use. In the EU is regulated in Commission Regulation N° 231/2012.

In order to improve these conventional extraction techniques, other methods have been developed (ultrasound assisted extraction, pressurized hot water extraction, negative pressure cavitation-assisted extraction and pulsed electric) to partner with them and solve some of the inopportuneness of traditional techniques. In this way the extraction efficiency is substantially improved.

4.2 Green extraction techniques

This group includes several techniques with the aim of not only preventing pollution but also reducing sample preparation costs by, for example, lowering solvents consumption.

4.2.1 Microwave-assisted extraction

Microwave-assisted extraction (MAE) is based on the heating of solvents that are in contact with solid samples with the use of microwaves. This allows the partition of compounds of interest from the sample into the solvent. Microwave energy accelerates a great variety of chemical reactions as well as the extraction of organic compounds from different matrices [98]. During the process, electromagnetic energy is transformed into calorific energy by two mechanisms: ionic conduction and dipole rotation [119]. It is a method that could be an alternative for avoiding thermal degradation and oxidation, with no influence on cell integrity and shape. However, due to thermal stress and localized high pressures, cell rupture is more rapid than with another techniques, which is an inconvenience, for example, in the extraction of volatile oils [91].

Some of the advantages of this technique are simplicity, effectiveness, low processing time, low solvent consumption and energy, no generation of secondary waste and can be used for larger volumes [120]. Another advantage of this method is that it produces a uniform heating, so extraction is simultaneous regardless of the area where the compound is [121]. By using this method, lipids, pigments, carbohydrates, vitamins and proteins can be extracted [122]. Moreover, it is characterized by having a superior extraction rate of volatile compounds in *Capsicum* spp.

The selectivity and efficiency of MAE largely depends on the dielectric constant of the extraction solvent mixture, which defines its chemical polarity and thus what compounds will be extracted [91]. It can be expected that by using water as a single solvent, the quantity of capsaicinoids extracted decrease due to its negligible polarity [109]. Therefore, the parameters to take into account in this type of extraction are time, power, temperature and type of solvent and its ratios.

Several experiments can be seen in a summarized form in **Table 7**, which shows the variation between the values of the parameters among different experiments with solvents of different polarity. It is worth highlighting the values obtained for capsaicinoids using ethanol [105] as solvents since it is the one with the highest yield.

4.2.2 Ultrasound assisted extraction

In every extraction process there is a critical step which is the degradation of cell walls and membranes. Ultrasound assisted extraction (UAE) makes this possible by applying pressure waves that are transmitted across a medium as compression and expansion (rarefaction) cycles, finding an area with maximum pressure in the compression phase and one with minimum pressure in the rarefaction phase. This pressure difference makes the cavitation phenomenon possible, which breaks the

Target	Species	Extracting conditions				Yield	Ref.
		Temp.	Time	Power	Solvent ratios		
		°C	min	W			
Volatiles	<i>Capsicum annuum</i>	55	10	250	n-hexane (pure)	42 compounds	[59]
Pigments	<i>Capsicum annuum</i>	60	2	50	Acetone-water (50%)	—	[91]
Capsaicinoids	<i>Capsicum</i> spp.	21	15	300	Acetone (30%)	0.48 mg/g	[98]
	<i>Capsicum chinense</i>	50	5	800	Methanol (100%)	0.32 mg/g	[109]
	<i>Capsicum frutescens</i>	—	20	320	Ethanol (99.5%)	5.3 mg/g	[105]
	<i>Capsicum frutescens</i>	120	15	150	Acetone (pure)	0.673%	[104]
	<i>Capsicum annuum</i>	76	1	500	Methanol (60%)	230 ppm	[97]
—, not described.							

Table 7.
Different experimental conditions carried out in peppers by microwave-assisted extraction.

structure to make the compounds accessible and therefore available to extract them [123].

This technique has been applied to extract different bioactive compounds from herbs or algae due to its ease to disrupt their cell walls despite its resistance which differentiates this method from the previous ones, since strong disruption of the cell wall is achieved and as a result the extraction of intracellular materials is increased with the increment of energy input [106, 120, 124].

The ultrasonic technique, as well as the other green techniques, has been proven to have several advantages such as reduction of solvents consumption, temperature and time, very important parameters in the extraction of thermolabile and unstable compounds as it has no effect on the chemical structure and biological properties. Furthermore, UAE has low equipment investment and easy implementation, so it can be basically industrially employed in local companies [106]. Moreover, there are studies that showed that this method also achieves a greater supercritical extraction of pungent compounds from ginger owing to physical effects on the surface of particles [106].

There are several parameters to take into account to optimize the method which include ultrasound power intensities, frequency, wavelength and time. There are several studies about it (**Table 8**). One of them studied the influence of several of these parameters on the extraction of capsaicinoids, observing that their release was very fast in the first 5 min of the process and then decreased. Temperature influence was also observed since an increase of 15°C (up to 45°C) improved the extraction using 95% ethanol (v/v) as a solvent. Higher temperatures did not lead to significant improvements [106]. The effect of the solvent was also studied, being the most common solvents used for extracting capsaicinoids methanol, ethanol, acetone, acetonitrile and water. It was proved that the yield of extraction is worse with the addition of water. Methanol and ethanol have similar recoveries but ethanol was better than acetone, the best concentration of ethanol being 95% [100, 106].

Target	Species	Conditions					Yield	Ref.
		Freq.	Intensity	Time	Temp.	Solvent		
		kHz	W/cm ³	min	°C	(%)		
Capsaicinoids	<i>Capsicum chinense</i>	20	—	10	40	Acetone (100)	0.31 mg/g	[109]
	<i>Capsicum chinense</i>	40	—	10	24	Methanol (100)	2.88 mg/g	[110]
	<i>Capsicum annuum</i>	31.5	—	20	76	Methanol (60)	200 ppm	[97]
	<i>Capsicum annuum</i>	—	—	20	50	n-hexane (100)	—	[99]
	<i>Capsicum annuum</i>	20	360	10	50	Methanol (100)	100%	[100]
	<i>Capsicum frutescens</i>	35	600	180	45	Ethanol (95)	87.4%	[106]
	<i>Capsicum frutescens</i>	20	360	10	50	Methanol (100)	100%	[100]
	<i>Capsicum frutescens</i>	—	—	40	25	Acetone (100)	3.92 mg/g	[107]
	<i>Capsicum baccatum</i>	20	150	20	40	Methanol (100)	50%	[112]
Antioxidants	<i>Capsicum baccatum</i>	20	150	20	40	Methanol (100)	27%	[112]
Oleoresin	<i>Capsicum baccatum</i>	20	450	20	60	Methanol (100)	26%	[112]
Phytochemicals	<i>Capsicum</i>	35	—	20	50	Methanol (80)	—	[114]
—, not described.								

Table 8.
Different experimental conditions carried out in peppers by ultrasound assisted extraction.

4.2.3 Enzyme-assisted extraction

Enzyme-assisted extraction (EAE) is an extraction system that allows the avoidance of processing conditions like temperature or drastic pH changes and so, maintain the quality and yield of multiple biomolecules [125].

This type of extraction supports isolation for recovering bioingredients from different plant materials. An enzymatic pre-treatment before applying traditional methods will help to isolate high yields of bioingredients due to enzyme assisted extraction facilities. This is due to the fact that degradation of cell walls and membranes is the critical step of extraction which [101]. Among its advantages over traditional methods are high selectivity, overall efficacy, eco-friendly procedures, low-energy consumption, minimal usage of harsh chemicals, maximum yield, low to no wasteful protection or deprotection steps, easy recovery, and process recyclability [126]. However, it also presents some drawbacks such as the cost of enzymes, requirement of holding tanks that may require long term incubation, lack of knowledge about optimal or compatible enzyme formulations for cell disruption and inability to completely hydrolyse the bonds in plant cell wall [127].

The factors to take into account in this technique are fundamentally temperature, pH, and type of enzyme. A range of enzymes (lipases, carbohydrase, celluloses, proteases, pectinases) have been widely used as specific catalysts. Each enzyme has different substrates. For example, cellulase and hemi-cellulase have their greater hydrolysing activity on the cellulose found in plant cell walls, hence their name. This enzymatic processing increases the permeability of the cell wall, resulting in a better recovery of some compounds like volatile oil and resin, which are prone to degradation when extracted with more disruptive methods [101].

Several studies (**Table 9**) show that viscozyme L is an enzyme with a superior recovery rate of bioactive compounds fractions like total carotenoid content, total phenolic content, total flavonoids and total antioxidant activity with high total suspended solids (TSS). It also has at this moment the better extract yield [95, 96]. Among its applications is the extraction of pigments or capsaicinoids. In almost all cases the pH of the medium is 4.5 as it is the pH of the sample and it is in the range of optimal activity of the different enzymes [95].

4.3 Emerging technologies for extraction

Due to the disadvantages of traditional techniques, there is an interest in the development of new extraction techniques. Some of the most sought-after features include shortened extraction time, automation or reduced organic solvent consumption.

4.3.1 Hydrostatic high-pressure extraction

Hydrostatic high pressure (HHP) is considered an emerging technology that have been applied in the preservation of food since the end of eighties [128]. The first food products treated with this method began commercialization in Japan in 1990 [129]. This method is established on the application of high pressures (100–900 MPa) to the product of interest.

Among the advantages of this type of extraction there is the possibility to conduct it at room temperature, meaning no thermal degradation and derived bioactivity loss from extracted components. It also does not modify chemical structures of the different compounds of interest’s independent form of molecular weight [130]. Moreover, in comparison with conventional techniques it is faster,

Target	Enzyme	Conditions			Yield	Ref.
		Temp	pH	Time		
		°C		h	%	
Carotenoids	Viscozyme L	60	4.5	1	87	[95]
	Pectinase	60	4.5	1	80	[95]
	Cellulase	60	4.5	1	low	[95]
	Viscozyme L	50	4.5	5	78	[96]
Capsaicinoids	Viscozyme L	50	4.5	5	88.8	[96]
	Viscozyme L	45	4.5	1	22	[101]
	Celluclast	45	4.5	1	20	[101]
Oleoresin	Viscozyme L	45	4.5	1	6	[101]

Table 9.
Different experimental conditions carried out in C. annum by enzyme-assisted extraction.

gives higher extraction yields and fewer impurities [131] giving, unlike other preservation technologies as thermal treatment, uniform and nearly instantaneous effects throughout the foodstuff and thus independent of foodstuff geometry and equipment size which makes an easy scale-up from laboratory findings to full-scale production possible [132].

This method can be applied to multiple matrices for the extraction of natural compounds [133] like fruit and vegetables for different targets as for example carotenoids [134], antioxidants [131] and pigments [135].

The main parameters to be taken into account are pressure, time, temperature and type and quantity of solvent, studying each parameter for each variety in particular.

HHP can change physiological and biochemical properties of pepper based on a study carried out with *C. annuum* [136]. In this research, different pressures (50, 100, 200 and 300 MPa) were applied for 5 min at 25°C observing that it produces remarkable changes in seedlings, but no or very little physiological and biochemical variations, creating an antioxidant system that is positive for the plant itself in the process. Therefore, HHP (500 MPa) can be used as a preservation treatment similar to pasteurization which was the traditional thermal treatment for sweet pepper preservation [137].

4.3.2 Supercritical-fluid extraction

In supercritical-fluid extraction (SFE), the fluid must reach temperature and pressure above the critical point, so as the fluid behaves like liquid and gas simultaneously which makes extraction easier. This technology has been used in a wide diversity of fields (food, pharmaceutical, chemical and fuel industries) due to its advantages, there is an absence of toxic residue in the final product which allows not only the extraction of valuable active compounds (fatty acids, pigments, polyphenols and vitamins) free of solvents, but also to remove undesirable compounds (pollutants, toxins and pesticides) [138]. Additional advantages are great extraction selectivity, short processing times, requirement of minimal solvents, low degradability of the extracted product and the fact that the remaining biomass can be treated with other techniques in order to continue extraction.

The most important conditions are temperature, pressure and co-solvent. The selection of each parameter will depend on the specific compound searched [138]. The most used solvent is CO₂ due to its thermodynamics and heat transfer properties. Moreover, it has a low critical point (31°C, 73 bar). Furthermore, the polarity of CO₂ can be modified by the use of co-solvents such as ethanol, and in this way also extract polar components.

Studies demonstrate that the extracts obtained with this technique are better than natural spice for flavoring purposes as SFE could reduce aflatoxin in the final products [139]. In addition, numerous bibliographic references show that carotenoid extraction is better at higher pressures [113].

Furthermore, when analyzing the extraction of β -carotene and capsaicin at the same time, capsaicin shows lower solubility, yet the solubility of β -carotene did not change in the presence of capsaicin, a factor to bear in mind when designing the separation process of coloring and hot components from paprika [140]. Several pepper species have been used to obtain natural compounds of interest using supercritical fluid extraction including capsaicinoids, oleoresins, pigments, tocopherols and even aflatoxins (**Table 10**) observing large variations between the pressures applied in each of the methods referred to in the bibliography. The greatest amount of studies is related to the specie *C. annuum* since it is the best known, widespread and cultivated specie of the genus *Capsicum*.

Target	Species	Pressure	Temp.	Flow rate	Time	Pepper	Yield	Ref.
		MPa	°C	cm s ⁻¹	min	g	%	
Capsaicinoids	<i>C. frutescens</i>	20.5	40	0.064	10	50	5.2	[108]
	<i>C. frutescens</i>	7.84 × 10 ⁻³	55	1	—	70	7.1 × 10 ⁻⁴	[103]
	<i>C. annuum</i>	55	40	—	—	—	8.6	[92]
	<i>C. annuum</i>	12	40	—	390	—	—	[102]
	<i>C. annuum</i>	40	55	0.9–1.2 ^a	—	—	93	[93]
	<i>C. annuum</i>	7.84 × 10 ⁻³	55	1	—	250	7.1 × 10 ⁻⁴	[103]
	<i>C. chinense</i>	15	60	2.15 × 10 ⁻³	90	2.5	0.5	[111]
Oleoresins	<i>C. frutescens</i>	21.5	40	0.071	10	50	0.3	[108]
	<i>C. annuum</i>	30	40	—	360	—	8.2	[102]
	<i>C. annuum</i>	40	35	1–1.5	—	—	7.4	[94]
	<i>C. annuum</i>	43–54	40	1	—	25–30	—	[113]
Pigments	<i>Capsicum</i> spp.	47.5	80	sm	—	—	6	[90]
	<i>C. annuum</i>	36	45	—	—	—	17.4	[92]
	<i>C. annuum</i>	20	35	0.9–1.2	—	—	93	[93]
	<i>C. annuum</i>	40	35	1–1.5	—	—	7.2	[94]
Aflatoxins	<i>C. annuum</i>	30	50	—	2	7.8	16.2	[139]
Tocopherols	<i>C. annuum</i>	5	25	0.9–1.2 ^a	—	—	84	[93]
	<i>C. annuum</i>	40	35	1–1.5	—	—	100	[94]

sm, semi-continuous; —, not described.
^aPropane.

Table 10.
Different experimental conditions carried out in peppers by supercritical-fluid extraction.

4.3.3 Pulse electric field extraction

Pulsed electric field (PEF) consists of a non-thermal method that is extensively used in food processing applications due to, among other things, its ability to kill microorganisms in liquid foods. This method involves the application of short duration electric pulses, making several pores on the cell membrane in what is called electroporeabilization or electroporation. This makes it possible for the selective recovery of intracellular components with low energy consumption as the dielectric breakdown theory explains. According to this theory, the membrane of the cell has a low-dielectric constant that when exposed to a strong electric field provokes ion migration, forming free charges of the opposite sign which accumulate at both membrane sides, generating a potential difference across the membrane which depends on the size and shape of the cells and the concentration of cells in suspension. This difference in charges makes cell walls undergo a compression, reducing the membrane thickness that results in the formation of micropores and increasing permeability (electroporation) [119, 141]. It can be used directly or as a pre-treatment prior to solvent extraction [142]. The parameters to take into account with this method are number of pulses, length of the pulses, energy input and biomass concentration.

Among the advantages of PEF is the short extraction time (usually under 1 s), efficiency at low temperatures, decreased energy losses and a successful cell wall breakdown [120]. The disadvantages are that membrane changes can be reversible,

air bubbles make the process less effective and the efficiency of the method depends on electric field strength and electrode gap [142].

Further research is necessary since there is no data on the extraction of compounds with this technique. However, there are studies of the application of PEF for the production of dehydrated products or juices. As for drying, a study [143] done with red bell pepper at 320 J/kg, 2.0 kV/cm, 1 Hz and 30 pulses of 400 μ s reported good results. Another study [144] with the same aim used different extraction conditions, which were 2.5 kV/cm, 100 Hz and 30 μ s as the pulse time. As for juice production, *C. annuum* was studied with extracting conditions of 500 J/kg, 1.7 kV/cm, 1 Hz and 30 pulses of 300 μ s demonstrating that juice from PEF treated paprika compared well in quality with enzymatic treatment [145].

5. Seed valorisation and industrial application

As it is being already done by some factories and companies in pepper growing countries, seeds, stalks or peels are extracted through whole fruit processing with or without semi-automated equipment. This is mainly done by the food industry in order to take away the non-edible parts of the vegetable to elaborate commodities as pepper powders (e.g., paprika), sauces, jams or pickled peppers. This way, stalks, seeds and placenta are usually removed and discarded.

But in many cases, placenta and seeds are subjected to extraction techniques in order to obtain essential oils to treat chronic ailments via topical administration in traditional remedies [146]. Even so, it is clear that pepper seeds are a multivalent source of many important compounds that could be used in different applications.

The elevated phenolic concentrations found in seeds make them an excellent target for extraction and use of these phenolic compounds, whether it is as antioxidant supplements, as food coating in order to increase shelf-life or as a potential source of functional animal feed. Moreover, the implementation of an extraction process in the food industry is dependent on the type of compounds of interest, as the yields of phytochemicals obtained may differ greatly. It should be noted that highly efficient and sustainable extraction methods continue to be a focus of research both in industry and academia [65].

Also, antimicrobial potential of phytochemicals present in pepper should be considered. Even though this would involve highly specialized equipment and maintain strict aseptic conditions, there is promise in their use as food preservers from common food-borne pathogenic microorganisms besides their antioxidative capability. Furthermore, their use as alternative antibiotics has hardly been researched and the safety of their administration or clinical use has not yet been assessed.

Regarding their nutrition attributes, the high linoleic acid and palmitic acid present in the seeds could lead to use them as a primary source of cooking oil, or as a main component in foods such as margarine, or preserver in certain pickled products [48]. Also, as they are a rich natural source of vitamins E and C, extracts of these vitamins could also be incorporated as constituents of many juices and other foods. Pepper-processing industry could find very rewarding to investigate on the yet unknown beneficial properties of its by-products and obtain additional profit by integrating existing extraction and purification technologies into their process chain.

6. Conclusions

It is clear that the wide spectrum of historical applications of peppers is supported by the latest findings on the properties of peppers. The diversity among

Capsicum species properties have been studied over the years, although many of the possible bioactivities have only been characterized in the whole fruit and there are few *in vivo* experiments reported testing phytochemicals of this genus or assessing its safety. Moreover, research on produce by-products has acquired weight within the scientific community and industry as of lately. The exploitation of food processing by-products means not only an affordable raw material for the extraction of valuable phytochemicals, but also a reduction of ecological and economic impact of agricultural and industrial activities. Further studies are necessary to explore in depth more efficient bio-active compounds from the underutilized and non-edible biomass of *Capsicum* spp. Furthermore, the great genotypic diversity of the many pepper varieties grown in different regions, have a high variety of bioactive compounds, some still to be discovered. Currently available knowledge about the medicinal phytochemicals found in the genus *Capsicum* should be considered to develop novel drugs to treat many diseases and ailments.

Acknowledgements

Many thanks to MICINN for the financial support for the Ramón & Cajal researcher of M.A. Prieto, to the Regional Government of Galicia to the Regional Government of Galicia for “Programa de axudas á etapa predoutoral da Xunta de Galicia” for the pre-doctoral researchers of A.G. Pereira and the authors are also grateful to Interreg España-Portugal for financial support through the 0377_Iberphenol_project.

Author details

Javier Echave¹, Antía G. Pereira^{1,2}, María Carpena¹, Miguel Ángel Prieto^{1*} and Jesús Simal-Gandara^{1*}

¹ Nutrition and Bromatology Group, Analytical and Food Chemistry Department, Faculty of Food Science and Technology, University of Vigo, Ourense, Spain

² Centro de Investigação de Montanha (CIMO), Instituto Politécnico de Bragança, Bragança, Portugal

*Address all correspondence to: mprieto@uvigo.es and jsimal@uvigo.es

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Pérez-castañeda M, Castañón-nájera G, Ramírez-meraz M. Avances y perspectivas sobre el estudio del origen y la diversidad genética de *Capsicum* spp. Ecosistemas y Recursos Agropecuarios. 2015;2:117-128
- [2] Simon JE, Chadwick AF, Craker LE. Herbs: An Indexed Bibliography 1971–1980: The Scientific Literature on Selected Herbs, and Aromatic and Medicinal Plants of the Temperate Zone. Hamden, CT: Archon Books; 1984
- [3] Melgar-Lalanne G, Hernández-Álvarez AJ, Jiménez-Fernández M, Azuara E. Oleoresins from *Capsicum* spp.: Extraction methods and bioactivity. Food and Bioprocess Technology. 2017;10:51-76
- [4] Pickersgill B. Migrations of chili peppers, *Capsicum* spp., in the Americas. In: Pre-Columbian Plant Migration. Papers of the Peabody Museum of Archaeology and Ethnology. 1984. pp. 105-123
- [5] Kraft KH, Brown CH, Nabhan GP, Luedeling E, De Jesús Luna Ruiz J, D'Eeckenbrugge GC, et al. Multiple lines of evidence for the origin of domesticated chili pepper, *Capsicum annuum*, in Mexico. Proceedings of the National Academy of Sciences of the United States of America. 2014;111:6165-6170
- [6] Castañón-Nájera G, Mayek-Pérez N, Ruiz-Salazar R, García AC. Molecular study of a diallel chilli with amplified fragment length polymorphism (AFLP) markers. African Journal of Agricultural Research. 2011;6:6126-6131
- [7] Hui YH, Chen F, Nollet LML, Guiné RPF, Le Quéré JL, Martín-Belloso O, et al. Handbook of Fruit and Vegetable Flavors. Hoboken, New Jersey: John Wiley and Sons Inc.; 2010
- [8] Basu SK, De AK. Capsicum: Historial and Botanical Perspectives in *Capsicum*, The genus *Capsicum*. London, UK: CRC Press; 2003. pp. 1-15
- [9] Martín NC, González WG. Caracterización de accesiones de Chile (*Capsicum* spp.). Agronomía Mesoamericana. 2016;2:31
- [10] Barrios O, Fuentes V, Abreu S. Especies cultivadas de ajíes y pimientos (*Capsicum* spp. div.) en Cuba. In: Proceedings of the Convención Trópico 2004. II Congreso de Agricultura Tropical. 2004. pp. 1-15
- [11] Castañón-Nájera G, Latournerie-Moreno L, Leshner-Gordillo JM, De La Cruz-Lázaro E, Mendoza-Elos M. Identification of variables for the morphological characterisation of hot peppers (*Capsicum* spp) in Tabasco, Mexico. Universidad y Ciencia. 2010;26(3):225-234
- [12] Katz E. Chili Pepper, from Mexico to Europe: Food, imaginary and cultural identity. In: Estudios del Hombre, Serie Antropología de la Alimentación. Food, Imaginaries and Cultural Frontiers. Essays in Honour of Helen Macbeth. Guadalajara, Mexico: Universidad de Guadalajara; 2009. pp. 213-232
- [13] Rajput J, Parulekar Y. El pimiento. In: Tratado de Ciencia y Tecnología de las hortalizas. Zaragoza: Acriba; 2004. pp. 203-225
- [14] Martínez S, Curros A, Bermúdez J, Carballo J, Franco I. The composition of Arnoia peppers (*Capsicum annuum* L.) at different stages of maturity. International Journal of Food Sciences and Nutrition. 2007;58:150-161
- [15] Orellana-Escobedo L, García-Amezquita LE, Olivas GI, Ornelas-Paz JJ, Sepulveda DR. Capsaicinoids content and proximate composition of Mexican chili peppers (*Capsicum* spp.) cultivated in the State of Chihuahua. CyTA—Journal of Food. 2013;11:179-184

- [16] Zou Y, Ma K, Tian M. Chemical composition and nutritive value of hot pepper seed (*Capsicum annuum*) grown in northeast region of China. *Journal of Food Science and Technology*. 2015; **35**:659-663
- [17] Esayas K, Shimelis A, Ashebir F, Negussie R, Tilahun B, Gulelat D. Proximate composition, mineral content and antinutritional factors of some capsicum (*Capsicum annum*) varieties grown in Ethiopia. *Bulletin of the Chemical Society of Ethiopia*. 2011; **25**: 451-454
- [18] Ku K-H, Choi E-J, Park JB. Chemical component analysis of red pepper (*Capsicum annuum* L.) seeds with various cultivars. *Journal of the Korean Society of Food Science and Nutrition*. 2008; **37**:1084-1089
- [19] Liu S, Li W, Wu Y, Chen C, De Lei J. Novo transcriptome assembly in chili pepper (*Capsicum frutescens*) to identify genes involved in the biosynthesis of Capsaicinoids. *PLoS One*. 2013; **8**:1-8
- [20] Arimboor R, Natarajan RB, Menon KR, Chandrasekhar LP, Moorkoth V. Red pepper (*Capsicum annuum*) carotenoids as a source of natural food colors: Analysis and stability—A review. *Journal of Food Science and Technology*. 2015; **52**: 1258-1271
- [21] Ciulu-Costinescu F, Chifiriuc MC, Popa M, Bleotu C, Neamtu J, Marina L, et al. Screening of polyphenol content and in vitro studies of antioxidant, antibacterial and cytotoxic activities of *Capsicum annuum* extracts. *Revista de Chimie—Bucharest*. 2015; **66**:1261-1266
- [22] Hernández-Ortega M, Ortiz-Moreno A, Hernández-Navarro MD, Chamorro-Cevallos G, Dorantes-Alvarez L, Necoechea-Mondragón H. Antioxidant, antinociceptive, and anti-inflammatory effects of carotenoids extracted from dried pepper (*Capsicum annuum* L.). *Journal of Biomedicine and Biotechnology*. 2012; **2012**:10. Article ID 524019
- [23] Srinivasan K. Biological activities of red pepper (*Capsicum annuum*) and its pungent principle capsaicin: A review. *Critical Reviews in Food Science and Nutrition*. 2016; **56**:1488-1500
- [24] Adaszek Ł, Gadomska D, Mazurek Ł, Łyp P, Madany J, Winiarczyk S. Properties of capsaicin and its utility in veterinary and human medicine. *Research in Veterinary Science*. 2019; **123**:14-19
- [25] Pandey MK, Gupta SC, Nabavizadeh A, Aggarwal BB. Regulation of cell signaling pathways by dietary agents for cancer prevention and treatment. *Seminars in Cancer Biology*. 2017; **46**:158-181
- [26] Rodrigues CA, Nicácio AE, Jardim ICSF, Visentainer JV, Maldaner L. Determination of phenolic compounds in red sweet pepper (*Capsicum annuum* L.) using a modified QuEChERS method and UHPLC-MS/MS Analysis and its relation to antioxidant activity. *Journal of the Brazilian Chemical Society*. 2019; **30**: 1229-1240
- [27] Iorizzi M, Lanzotti V, Giancarlo R, De Marino S, Zollo F. Antimicrobial Furostanol Saponins from the seeds of *Capsicum annuum* L. Var. *acuminatum*. *Journal of Agricultural and Food Chemistry*. 2002; **50**:4310-4316
- [28] Kuku A, Odekanyin O, Adeniran K, Adewusi M, Olonade T, et al. Purification of a mannose/glucose-specific lectin with antifungal activity from pepper seeds (*Capsicum annum*). *African Journal of Biochemistry Research*. 2009; **3**:272-278
- [29] Silva LR, Azevedo J, Pereira MJ, Valentão P, Andrade PB. Chemical assessment and antioxidant capacity of

- papper (
- Capsicum annuum*
- L.) seeds. Food and Chemical Toxicology. 2013;53: 240-248
- [30] Sandoval-Castro CJ, Valdez-Morales M, Oomah BD, Gutiérrez-Dorado R, Medina-Godoy S, Espinosa-Alonso LG. Bioactive compounds and antioxidant activity in scalded Jalapeño pepper industrial byproduct (*Capsicum annuum*). Journal of Food Science and Technology. 2017;54:1999-2010
- [31] Nascimento PLA, Nascimento TCES, Ramos NSM, Silva GR, Gomes JEG, Falcão REA, et al. Quantification, antioxidant and antimicrobial activity of phenolics isolated from different extracts of *Capsicum frutescens* (Pimenta Malagueta). Molecules. 2014;19: 5434-5447
- [32] Diz MSS, Carvalho AO, Rodrigues R, Neves-Ferreira AGC, da Cunha M, Alves EW, et al. Antimicrobial peptides from chilli pepper seeds causes yeast plasma membrane permeabilization and inhibits the acidification of the medium by yeast cells. Biochimica et Biophysica Acta (BBA)—General Subjects. 1760; 2006:1323-1332
- [33] Aguirre J, Ríos-Momberg M, Hewitt D, Hansberg W. Reactive oxygen species and development in microbial eukaryotes. Trends in Microbiology. 2005;13:111-118
- [34] Chamikara MDM, Dissanayake DRRP, Ishan M, Sooriyapathirana SDSS. Dietary, anticancer and medicinal properties of the phytochemicals in chili pepper (*Capsicum* spp.). Ceylon Journal of Science. 2016;3:5-20
- [35] Mokhtar M, Ginestra G, Youcefi F, Filocamo A, Bisignano C, Riazi A. Antimicrobial activity of selected polyphenols and capsaicinoids identified in pepper (*Capsicum annuum* L.) and their possible mode of interaction. Current Microbiology. 2017;74: 1253-1260
- [36] Badhani B, Sharma N, Kakkar R. Gallic acid: A versatile antioxidant with promising therapeutic and industrial applications. RSC Advances. 2015;5: 27540-27557
- [37] Kevers C, Falkowski M, Tabart J, Defraigne JO, Dommes J, Pincemail J. Evolution of antioxidant capacity during storage of selected fruits and vegetables. Journal of Agricultural and Food Chemistry. 2007;55:8596-8603
- [38] Bertão MR, Moraes MC, Palmieri DA, Silva LP, da Silva RMG. Cytotoxicity, genotoxicity and antioxidant activity of extracts from *Capsicum* spp. Research Journal of Medicinal Plant. 2016;10: 265-275
- [39] Sim KH, Sil HY. Antioxidant activities of red pepper (*Capsicum annuum*) pericarp and seed extracts. International Journal of Food Science and Technology. 2008;43:1813-1823
- [40] Yahfoufi N, Alsadi N, Jambi M, Matar C. The immunomodulatory and anti-inflammatory role of polyphenols. Nutrients. 2018;10:1-23
- [41] Aza-González C, Núñez-Palenius HG, Ochoa-Alejo N. Molecular biology of capsaicinoid biosynthesis in chili pepper (*Capsicum* spp.). Plant Cell Reports. 2011;30:695-706
- [42] Chouaibi M, Rezig L, Hamdi S, Ferrari G. Chemical characteristics and compositions of red pepper seed oils extracted by different methods. Industrial Crops and Products. 2019;128: 363-370
- [43] Howard LR, Talcott ST, Brenes CH, Villalon B. Changes in phytochemical and antioxidant activity of selected pepper cultivars (*Capsicum* species) as

influenced by maturity. *Journal of Agricultural and Food Chemistry*. 2000; **48**:1713-1720

[44] Deepa N, Kaur C, George B, Singh B, Kapoor HC. Antioxidant constituents in some sweet pepper (*Capsicum annuum* L.) genotypes during maturity. *LWT—Food Science and Technology*. 2007; **40**: 121-129

[45] Kobata K, Todo T, Yazawa S, Iwai K, Watanabe T. Novel capsaicinoid-like substances, capsiate and dihydrocapsiate, from the fruits of a nonpungent cultivar, CH-19 sweet, of pepper (*Capsicum annuum* L.). *Journal of Agricultural and Food Chemistry*. 1998; **46**:1695-1697

[46] Luo XJ, Peng J, Li YJ. Recent advances in the study on capsaicinoids and capsinoids. *European Journal of Pharmacology*. 2011; **650**:1-7

[47] Ludy MJ, Moore GE, Mattes RD. The effects of capsaicin and capsiate on energy balance: Critical review and meta-analyses of studies in humans. *Chemical Senses*. 2012; **37**:103-121

[48] Jarret RL, Levy IJ, Potter TL, Cermak SC. Seed oil and fatty acid composition in *Capsicum* spp. *Journal of Food Composition and Analysis*. 2013; **30**:102-108

[49] Matthäus B, Özcan MM. Chemical evaluation of some paprika (*Capsicum annuum* L.) seed oils. *European Journal of Lipid Science and Technology*. 2009; **111**:1249-1254

[50] Ristić-Medić D, Vučić V, Takić M, Karadžić I, Glibetić M. Polyunsaturated fatty acids in health and disease. *Journal of the Serbian Chemical Society*. 2013; **78**:1269-1289

[51] Materska M, Perucka I. Antioxidant activity of the main phenolic compounds isolated from hot pepper fruit (*Capsicum annuum* L.).

Journal of Agricultural and Food Chemistry. 2005; **53**:1750-1756

[52] Chu YF, Sun J, Wu X, Liu RH. Antioxidant and antiproliferative activities of common vegetables. *Journal of Agricultural and Food Chemistry*. 2002; **50**:6910-6916

[53] Hernández-Ortega M, Ortiz-Moreno A, Hernández-Navarro MD, Chamorro-Cevallos G, Dorantes-Alvarez L, Necoechea-Mondragón H, et al. Antioxidant, antinociceptive, and anti-inflammatory effects of carotenoids extracted from dried pepper (*Capsicum annuum* L.). *Journal of Biomedicine & Biotechnology*. 2012; **2012**:1-10

[54] Rao AV, Rao LG. Carotenoids and human health. *Pharmacological Research*. 2007; **55**:207-216

[55] Mocchegiani E, Costarelli L, Giacconi R, Malavolta M, Basso A, Piacenza F, et al. Vitamin E-gene interactions in aging and inflammatory age-related diseases: Implications for treatment. A systematic review. *Ageing Research Reviews*. 2014; **14**:81-101

[56] Saini RK, Keum YS. GC-MS and HPLC-DAD analysis of fatty acids and tocopherols in sweet peppers (*Capsicum annuum* L.). *Journal of Food Measurement and Characterization*. 2016; **10**:685-689

[57] Asnin L, Park SW. Isolation and analysis of bioactive compounds in capsicum peppers. *Critical Reviews in Food Science and Nutrition*. 2015; **55**: 254-289

[58] Shahidi F, de Camargo AC. Tocopherols and tocotrienols in common and emerging dietary sources: Occurrence, applications, and health benefits. *International Journal of Molecular Sciences*. 2016; **17**:1745

[59] Gogus F, Ozel MZ, Keskin H, Yanik DK, Lewis AC. Volatiles of fresh and commercial sweet red pepper

pastes: Processing methods and microwave assisted extraction. International Journal of Food Properties. 2015;18:1625-1634

[60] De Lucca AJ, Bland JM, Vigo CB, Cushion M, Selitrennikoff CP, Peter J, et al. CAY-1, a fungicidal saponin from *Capsicum* sp. fruit. Medical Mycology. 2002;40:131-137

[61] Simonovska J, Škerget M, Knez Ž, Srbinoska M, Kavrakovski Z, Grozdanov A, et al. Physicochemical characterization and bioactive compounds of stalk from hot fruits of *Capsicum annuum* L. Macedonian Journal of Chemistry and Chemical Engineering. 2016;35:199-208

[62] Jang H-W, Ka M-H, Lee K-G. Antioxidant activity and characterization of volatile extracts of *Capsicum annuum* L. and *Allium* spp. Flavour and Fragrance Journal. 2008;23:178-184

[63] Cichewicz RH, Thorpe PA. The antimicrobial properties of Chile peppers (*Capsicum* species) and their uses in Mayan medicine. Journal of Ethnopharmacology. 1996;52:61-70

[64] Von Borowski RG, Barros MP, da Silva DB, Lopes NP, Zimmer KR, Staats CC, et al. Red pepper peptide coatings control *Staphylococcus epidermidis* adhesion and biofilm formation. International Journal of Pharmaceutics. 2020;574:118872

[65] de Camargo AC, Schwember AR, Parada R, Garcia S, Maróstica Júnior MR, Franchin M, et al. Opinion on the hurdles and potential health benefits in value-added use of plant food processing by-products as sources of phenolic compounds. International Journal of Molecular Sciences. 2018:19

[66] Shahidi F, Ambigaipalan P. Phenolics and polyphenolics in foods, beverages and spices: Antioxidant

activity and health effects—A review. Journal of Functional Foods. 2015;18:820-897

[67] Azabou S, Ben Taheur F, Jridi M, Bouaziz M, Nasri M. Discarded seeds from red pepper (*Capsicum annuum*) processing industry as a sustainable source of high added-value compounds and edible oil. Environmental Science and Pollution Research. 2017;24:22196-22203

[68] Sora GTS, Haminiuk CWI, da Silva MV, Zielinski AAF, Gonçalves GA, Bracht A, et al. A comparative study of the capsaicinoid and phenolic contents and in vitro antioxidant activities of the peppers of the genus *Capsicum*: An application of chemometrics. Journal of Food Science and Technology. 2015;52:8086-8094

[69] Hallmann E, Rembial Kowska E. Characterisation of antioxidant compounds in sweet bell pepper (*Capsicum annuum* L.) under organic and conventional growing systems. Journal of the Science of Food and Agriculture. 2012;92:2409-2415

[70] Dorantes L, Colmenero R, Hernandez H, Mota L, Jaramillo ME, Fernandez E, et al. Inhibition of growth of some foodborne pathogenic bacteria by *Capsicum annuum* extracts. International Journal of Food Microbiology. 2000;57:125-128

[71] Bacon K, Boyer R, Denbow C, O'Keefe S, Neilson A, Williams R. Antibacterial activity of jalapeño pepper (*Capsicum annuum* var. *annuum*) extract fractions against select foodborne pathogens. Food Science & Nutrition. 2017;5:730-738

[72] Von Borowski RG, Zimmer KR, Leonardi BF, Trentin DS, Silva RC, de Barros MP, et al. Red pepper *Capsicum baccatum*: Source of antiadhesive and antibiofilm compounds against nosocomial bacteria. Industrial Crops and Products. 2019;127:148-157

- [73] Kang JH, Kim CS, Han IS, Kawada T, Yu R. Capsaicin, a spicy component of hot peppers, modulates adipokine gene expression and protein release from obese-mouse adipose tissues and isolated adipocytes, and suppresses the inflammatory responses of adipose tissue macrophages. *FEBS Letters*. 2007;**581**:4389-4396
- [74] Sancho R, Lucena C, Macho A, Calzado MA, Blanco-Molina M, Minassi A, et al. Immunosuppressive activity of capsaicinoids: Capsiate derived from sweet peppers inhibits NF- κ B activation and is a potent antiinflammatory compound in vivo. *European Journal of Immunology*. 2002;**32**:1753-1763
- [75] Choi S-E, Kim TH, Yi S-A, Hwang YC, Hwang WS, Choe SJ, et al. Capsaicin attenuates palmitate-induced expression of macrophage inflammatory protein 1 and interleukin 8 by increasing palmitate oxidation and reducing c-Jun activation in THP-1 (human acute monocytic leukemia cell) cells. *Nutrition Research*. 2011;**31**:468-478
- [76] Metcalfe NB, Alonso-Alvarez C. Oxidative stress as a life-history constraint: The role of reactive oxygen species in shaping phenotypes from conception to death. *Functional Ecology*. 2010;**24**:984-996
- [77] Rao V, Al-Weshahy A. Plant-based diets and control of lipids and coronary heart disease risk. *Current Atherosclerosis Reports*. 2008;**10**:478-485
- [78] Tundis R, Loizzo MR, Menichini F, Bonesi M, Conforti F, Statti G, et al. Comparative study on the chemical composition, antioxidant properties and hypoglycaemic activities of two *Capsicum annum* L. cultivars (*Acuminatum* small and *Cerasiferum*). *Plant Foods for Human Nutrition*. 2011;**66**:261-269
- [79] Marini E, Magi G, Mingoia M, Pugnali A, Facinelli B. Antimicrobial and anti-virulence activity of capsaicin against erythromycin-resistant, cell-invasive group A streptococci. *Frontiers in Microbiology*. 2015;**6**:1-7
- [80] Lee IO, Lee KH, Pyo JH, Kim JH, Choi YJ, Lee YC. Anti-inflammatory effect of capsaicin in *Helicobacter pylori*-infected gastric epithelial cells. *Helicobacter*. 2007;**12**:510-517
- [81] Yamasaki S, Asakura M, Neogi SB, Hinenoya A, Iwaoka E, Aoki S. Inhibition of virulence potential of *Vibrio cholerae* by natural compounds. *The Indian Journal of Medical Research*. 2011;**133**:232
- [82] De Lucca AJ, Boue S, Sien T, Cleveland TE, Walsh TJ. Silver enhances the in vitro antifungal activity of the saponin, CAY-1. *Mycoses*. 2011;**54**:e1-e9
- [83] Bashyal P, Parajuli P, Pandey RP, Sohng JK. Microbial biosynthesis of antibacterial chrysoeriol in recombinant *Escherichia coli* and bioactivity assessment. *Catalysts*. 2019;**9**(2):1-15
- [84] Salehi B, Hernández-Álvarez AJ, Contreras MDM, Martorell M, Ramírez-Alarcón K, Melgar-Lalanne G, et al. Potential phytopharmacy and food applications of *Capsicum* spp.: A comprehensive review. *Natural Product Communications*. 2018;**13**:1543-1556
- [85] Ng TB, Ngai PHK, Cheung RCF, Wong JH, Lam SK, Wang HX, et al. Antifungal and mitogenic activities of cluster pepper (*Capsicum frutescens*) seeds. *Nuts and Seeds in Health and Disease Prevention*. 2011:345-349
- [86] Malek N, Mrugala M, Makuch W, Kolosowska N, Przewlocka B, Binkowski M, et al. A multi-target approach for pain treatment: Dual inhibition of fatty acid amide hydrolase

and TRPV1 in a rat model of osteoarthritis. *Pain*. 2015;**156**:890-903

[87] Du Q, Liao Q, Chen C, Yang X, Xie R, Xu J. The role of transient receptor potential Vanilloid 1 in common diseases of the digestive tract and the cardiovascular and respiratory system. *Frontiers in Physiology*. 2019;**10** (1064):1-17

[88] Deng Y, Huang X, Wu H, Zhao M, Lu Q, Israeli E, et al. Some like it hot: The emerging role of spicy food (capsaicin) in autoimmune diseases. *Autoimmunity Reviews*. 2016;**15**: 451-456

[89] Sharma SK, Vij AS, Sharma M. Mechanisms and clinical uses of capsaicin. *European Journal of Pharmacology*. 2013;**720**:55-62

[90] Skerget M, Knez Z. Modeling high pressure extraction process. *Computers & Chemical Engineering*. 2001;**25**: 879-886

[91] Csiktusnádi Kiss GA, Forgács E, Cserhádi T, Mota T, Morais H, Ramos A. Optimisation of the microwave-assisted extraction of pigments from paprika (*Capsicum annuum* L.) powders. *Journal of Chromatography A*. 2000;**889**:41-49

[92] Fernández-Trujillo JP. Supercritical CO₂ extraction of sweet and hot paprika oleoresin and other fractions. *Grasas y Aceites*. 2008;**59**(1):7-15

[93] Gnayfeed MH, Daood HG, Illés V, Biacs PA. Supercritical CO₂ and subcritical propane extraction of pungent paprika and quantification of carotenoids, tocopherols, and capsaicinoids. *Journal of Agricultural and Food Chemistry*. 2001;**49**:2761-2766

[94] Illés V, Daood HG, Biacs PA, Gnayfeed MH, Mészáros B. Supercritical CO₂ and subcritical propane extraction of spice red pepper oil with special regard to carotenoid and tocopherol

content. *Journal of Chromatographic Science*. 1999;**37**:345-352

[95] Nath P, Kaur C, Rudra SG, Varghese E. Enzyme-assisted extraction of carotenoid-rich extract from red capsicum (*Capsicum annuum*). *Agricultural Research*. 2016;**5**:193-204

[96] Santamaría RI, Reyes-Duarte MD, Bárzana E, Fernando D, Gama FM, Mota M, et al. Selective enzyme-mediated extraction of capsaicinoids and carotenoids from chili guajillo puya (*Capsicum annuum* L.) using ethanol as solvent. *Journal of Agricultural and Food Chemistry*. 2000;**48**:3063-3067

[97] Paduano A, Caporaso N, Santini A, Sacchi R. Microwave and ultrasound-assisted extraction of capsaicinoids from chili peppers (*Capsicum annuum* L.) in flavored olive oil. *Journal of Food Research*. 2014;**3**:51

[98] Williams OJ, Raghavan GSV, Orsat V, Dai J. Microwave-assisted extraction of capsaicinoids from capsicum fruit. *Journal of Food Biochemistry*. 2004;**28**:113-122

[99] Rafajlovska V, Klopcevska J, Klopcevska J, Srbinoska M, Srbinoska M, Raicki SR. Ultrasound-assisted extraction of capsaicinoids and carotenoids from hot red pepper. In: *Proceedings of the 5th Black Sea Basin Conference of Analytical Chemistry*. 2009

[100] Barbero GF, Liazid A, Palma M, Barroso CG. Ultrasound-assisted extraction of capsaicinoids from peppers. *Talanta*. 2008;**75**:1332-1337

[101] Baby KC, Ranganathan TV. Effect of enzyme pretreatment on yield and quality of fresh green chilli (*Capsicum annuum* L.) oleoresin and its major capsaicinoids. *Biocatalysis and Agricultural Biotechnology*. 2016;**7**: 95-101

- [102] Govindarajan VS. Capsicum—Production, technology, chemistry, and quality—Part II. Processed products, standards, world production and trade. *Critical Reviews in Food Science and Nutrition*. 1986;**23**: 207-288
- [103] Peusch M, Müller-Seitz E, Petz M, Müller A, Anklam E. Extraction of capsaicinoids from chillies (*Capsicum frutescens* L.) and paprika (*Capsicum annuum* L.) using supercritical fluids and organic solvents. *European Food Research and Technology*. 1997;**204**:351-355
- [104] Nazari F, Ebrahimi SN, Talebi M, Rassouli A, Bijanzadeh HR. Multivariate optimisation of microwave-assisted extraction of capsaicin from *Capsicum frutescens* L. and quantitative analysis by ¹H-NMR. *Phytochemical analysis*. 2007;**18**:333-340
- [105] Chuichulcherm S, Prommakort S, Srinophakun P, Thanapimmetha A. Optimization of capsaicin purification from *Capsicum frutescens* Linn. with column chromatography using Taguchi design. *Industrial Crops and Products*. 2013;**44**:473-479
- [106] Boonkird S, Phisalaphong C, Phisalaphong M. Ultrasound-assisted extraction of capsaicinoids from *Capsicum frutescens* on a lab- and pilot-plant scale. *Ultrasonics Sonochemistry*. 2008;**15**:1075-1079
- [107] Deng X-Y. Optimization of ultrasonic-assisted extraction procedure of capsaicinoids from chili peppers using orthogonal array experimental design. *African Journal of Biotechnology*. 2012;**11**:13153-13161
- [108] Duarte C, Moldão-Martins M, Gouveia AF, da Costa SB, Leitão AE, Bernardo-Gil MG. Supercritical fluid extraction of red pepper (*Capsicum frutescens* L.). *Journal of Supercritical Fluids*. 2004;**30**:155-161
- [109] Vázquez-Espinosa M, De Peredo AVG, Ferreiro-González M, Barroso CG, Palma M, Barbero GF, et al. Optimizing and comparing ultrasound- and microwave-assisted extraction methods applied to the extraction of antioxidant capsinoids in peppers. *Agronomy*. 2019;**9**(633):1-18
- [110] Sganzerla M, Coutinho JP, de Melo AMT, Godoy HT. Fast method for capsaicinoids analysis from *Capsicum chinense* fruits. *Food Research International*. 2014;**64**:718-725
- [111] de Aguiar AC, dos Santos P, Coutinho JP, Barbero GF, Godoy HT, Martínez J. Supercritical fluid extraction and low pressure extraction of Biquinho pepper (*Capsicum chinense*). *LWT—Food Science and Technology*. 2014;**59**: 1239-1246
- [112] Dias ALB, Arroio Sergio CS, Santos P, Barbero GF, Rezende CA, Martínez J. Ultrasound-assisted extraction of bioactive compounds from dedo de moça pepper (*Capsicum baccatum* L.): Effects on the vegetable matrix and mathematical modeling. *Journal of Food Engineering*. 2017;**198**: 36-44
- [113] Uquiche E, del Valle JM, Ortiz J. Supercritical carbon dioxide extraction of red pepper (*Capsicum annuum* L.) oleoresin. *Journal of Food Engineering*. 2004;**65**:55-66
- [114] Sricharoen P, Lamaiphan N, Patthawaro P, Limchoowong N, Techawongstien S, Chanthai S. Phytochemicals in capsicum oleoresin from different varieties of hot chilli peppers with their antidiabetic and antioxidant activities due to some phenolic compounds. *Ultrasonics Sonochemistry*. 2017;**38**:629-639
- [115] Cacace JE, Mazza G. Mass transfer process during extraction of phenolic compounds from milled berries. *Journal of Food Engineering*. 2003;**59**:379-389

- [116] Fernández-Ronco MP, Gracia I, de Lucas A, Rodríguez JF. Extraction of *Capsicum annuum* oleoresin by maceration and ultrasound-assisted extraction: Influence of parameters and process modeling. *Journal of Food Process Engineering*. 2013;**36**:343-352
- [117] Yilmaz E, Sevgi Arsunar E, Aydeniz B, Güneşer O. Cold pressed capia pepperseed (*Capsicum annuum* L.) oils: Composition, aroma, and sensory properties. *European Journal of Lipid Science and Technology*. 2015;**33**(8): 926-932
- [118] Melgar-Lalanne G, Hernández-Álvarez AJ, Jiménez-Fernández M, Azuara E. Oleoresins from *Capsicum* spp.: Extraction methods and bioactivity. *Food and Bioprocess Technology*. 2017;**10**:51-76
- [119] Kalil SJ, Moraes CC, Sala L, Burkert CAV. Bioproduct extraction from microbial cells by conventional and nonconventional techniques. In: *Food Bioconversion*. Vol. 2. London, UK: Elsevier Ltd; 2017. pp. 179-206
- [120] Ventura SPM, Nobre BP, Ertekin F, Hayes M, García-Vaquero M, Vieira F, et al. Extraction of value-added compounds from microalgae. In: *Microalgae-Based Biofuels and Bioproducts: From Feedstock Cultivation to End-Products*. Kidlington, UK: Elsevier Ltd; 2017. pp. 461-483
- [121] Kim S, Chojnacka K. In: Kim S, Chojnacka K, editors. *Marine Algae Extracts*. Weinheim: Wiley-VCH; 2015. ISBN: 9783527333271
- [122] Kapoore R, Butler T, Pandhal J, Vaidyanathan S. Microwave-assisted extraction for microalgae: From biofuels to biorefinery. *Biology (Basel)*. 2018;**7**:18
- [123] Lavilla I, Bendicho C. Fundamentals of ultrasound-assisted extraction. In: *Water Extraction of Bioactive Compounds: From Plants to Drug Development*. Kidlington, UK: Elsevier Ltd; 2017. pp. 291-316
- [124] Gerde JA, Montalbo-Lomboy M, Yao L, Grewell D, Wang T. Evaluation of microalgae cell disruption by ultrasonic treatment. *Bioresource Technology*. 2012;**125**:175-181
- [125] Baby KC, Ranganathan TV. Enzyme-assisted extraction of bioingredients. *Chemical Weekly*. 2013; **1**:213-224
- [126] Alam MA, Sarker MZI, Ghafoor K, Happy RA, Ferdosh S. Bioactive compounds and extraction techniques. In: *Recovering Bioactive Compounds from Agricultural Wastes*. Chennai, India: John Wiley & Sons Ltd; 2017. pp. 33-53
- [127] Nadar SS, Rao P, Rathod VK. Enzyme assisted extraction of biomolecules as an approach to novel extraction technology: A review. *Food Research International*. 2018;**108**: 309-330
- [128] Rastogi NK, Raghavarao KSMS, Balasubramaniam VM, Niranjan K, Knorr D. Opportunities and challenges in high pressure processing of foods. *Critical Reviews in Food Science and Nutrition*. 2007;**47**:69-112
- [129] Daher D, Le Gourrierc S, Pérez-Lamela C. Effect of high pressure processing on the microbial inactivation in fruit preparations and other vegetable based beverages. *Agriculture*. 2017;**7**(9): 1-18
- [130] Tadapaneni RK, Daryaei H, Krishnamurthy K, Edirisinghe I, Burton-Freeman BM. High-pressure processing of berry and other fruit products: Implications for bioactive compounds and food safety. *Journal of Agricultural and Food Chemistry*. 2014; **62**:3877-3885

- [131] Huang HW, Hsu CP, Yang BB, Wang CY. Advances in the extraction of natural ingredients by high pressure extraction technology. *Trends in Food Science and Technology*. 2013;**33**:54-62
- [132] Torres JA, Velazquez G. Commercial opportunities and research challenges in the high pressure processing of foods. *Journal of Food Engineering*. 2005;**67**:95-112
- [133] Dixon C, Wilken LR. Green microalgae biomolecule separations and recovery. *Bioresources and Bioprocessing*. 2018;**5**(14):1-24
- [134] Poojary MM, Barba FJ, Aliakbarian B, Donsì F, Pataro G, Dias DA, et al. Innovative alternative technologies to extract carotenoids from microalgae and seaweeds. *Marine Drugs*. 2016;**14**(11):1-34
- [135] Jubeau S, Marchal L, Pruvost J, Jaouen P, Legrand J, Fleurence J. High pressure disruption: A two-step treatment for selective extraction of intracellular components from the microalga *Porphyridium cruentum*. *Journal of Applied Phycology*. 2013;**25**:983-989
- [136] İşlek C, Altuner EM, Alpas H. The effect of high hydrostatic pressure on the physiological and biochemical properties of pepper (*Capsicum annuum* L.) seedlings. *High Pressure Research*. 2015;**35**:396-404
- [137] Hernández-Carrión M, Hernando I, Quiles A. High hydrostatic pressure treatment as an alternative to pasteurization to maintain bioactive compound content and texture in red sweet pepper. *Innovative Food Science and Emerging Technologies*. 2014;**26**:76-85
- [138] Pereira CG, Meireles MAA. Supercritical fluid extraction of bioactive compounds: Fundamentals, applications and economic perspectives. *Food and Bioprocess Technology*. 2010;**3**:340-372
- [139] Ehlers D, Czech E, Quirin KW, Weber R. Distribution of aflatoxins between extract and extraction residue of paprika using supercritical carbon dioxide. *Phytochemical Analysis*. 2006;**17**:114-120
- [140] Škerget M, Knez Ž. Solubility of binary solid mixture β -carotene-capsaicin in dense CO₂. *Journal of Agricultural and Food Chemistry*. 1997;**45**:2066-2069
- [141] Ricci A, Parpinello GP, Versari A. Recent advances and applications of pulsed electric fields (PEF) to improve polyphenol extraction and color release during red winemaking. *Beverages*. 2018;**4**(1):1-12
- [142] Joannes C, Sipaut CS, Dayou J, Yasir SM, Mansa RF. The potential of using pulsed electric field (PEF) technology as the cell disruption method to extract lipid from microalgae for biodiesel production. *International Journal of Renewable Energy Resources*. 2015;**5**:598-621
- [143] Ade-Omowaye BIO, Rastogi NK, Angersbach A, Knorr D. Combined effects of pulsed electric field pretreatment and partial osmotic dehydration on air drying behaviour of red bell pepper. *Journal of Food Engineering*. 2003;**60**:89-98
- [144] Won YC, Min SC, Lee DU. Accelerated drying and improved color properties of red pepper by pretreatment of pulsed electric fields. *Drying Technology*. 2015;**117**:1016-1026
- [145] Ade-Omowaye BIO, Angersbach A, Taiwo KA, Knorr D. The use of pulsed electric fields in producing juice from paprika (*Capsicum annuum* L.). *Journal of Food Processing and Preservation*. 2001;**25**:353-365
- [146] Campos-Vega R, Oomah BD, Vergara-Castaneda HA. Vegetable byproducts. In: *Food Wastes and Byproducts: Nutraceutical and Health Potential*. Pondicherry, India: John Wiley & Sons; 2019. pp. 254-255