We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Pore-Forming Proteins: Fluorescent Dyes to Study the Channel Functionality and Biophysical Properties

Guilherme Teixeira and Robson Faria

Abstract

Large conductance channels, when activated transiently, alter the plasma membrane permeability permitting the passage of molecules with the size of until 1000 Da. These pore-forming proteins are found in the plasma membrane of mammals and invertebrate cells such as the P2X7 receptor, transient receptor potential vanilloid 1 (TRPV1), transient potential melastin receptor 8 (TRPM8), pannexin-1, and hemichannel proteins. Other proteins may be secreted for organisms creating nonselective pathways to large molecules and proteins. Some examples are α -hemolysin and gramicidin. In both cases, an efficient manner to evaluate the functionality is using fluorescent dyes. Thus, we discuss the uses of the fluorescent dyes as tools to study the functionality and some biophysical properties of the proteins described above, among others.

Keywords: fluorescent dyes, ion channels, pore-forming proteins

1. Introduction

Techniques involving fluorescence have been used and improved in studies involving cells, the visualization, and research of their diverse biochemical structures and functions. Detection of proteins, recognition of organelles, nucleic acids, membrane permeabilization, and ions present in cells are widely explored through the feasibility of fluorescent dyes according to their physical-chemical characteristics such as excitation and emission lengths, electrical charges, chemical structures, and molecular weights [1, 2].

Ion channel-forming ionotropic receptors are extremely important proteins in maintaining the potential of cell membranes. In some cases, they are able to form pores (high conductance channels) in the plasma membrane, allowing the passage of certain ions of varying charges and different physical characteristics according to the concentration and electrochemical gradient. Thus, there are a large number of high conductance ion channels capable of opening pores, changing the permeability of the plasma membrane. As examples, we can mention the P2X7 purinergic receptor, transient receptor potential vanilloid (TRPV), voltage-dependent anion channel (VDAC), and others. When stimulated, they can allow ions to pass through

the formed pores and molecules with up to 1000 Da to the intracellular medium [3–7]. Thus, certain fluorescent dyes can be used to observe these pores of high ionic conductance, thus being able to further explore about the biophysical and chemical characteristics of these membrane proteins.

2. Pore-forming ionic channels in mammals

The formation of pores in the plasma membrane is a phenomenon often caused by defense proteins in mammalian cells as an immune response to infections by pathogens, tumors, or danger signs. As a response, several proteins are activated, forming pores such as perforin, gasdermin, complement system, mixed lineage kinase domain-like pseudokinase (MLKL), and granulysin, thus mediating cytotoxic and antimicrobial immune activities aimed at preventing and protecting the body from harmful agents [8, 9].

In addition to cellular defense proteins, ion channels are another class of proteins involved in the formation of pores in the plasma membrane, being responsible for the transit of ions such as selective to Ca^{2+} , N^+ , K^+ , and Cl^- and acting in a physiological and pathophysiological way. There are a large number of ion channels addressed in the literature related to the pore opening. Among them, the P2X7 receptor has been highlighted. The P2X7 receptor, which is the most studied among P2X, is a cationic channel activated by adenosine-5'-triphosphate (ATP) [10, 11]. This channel is responsible for several intracellular processes, such as apoptosis, NLRP3 inflammasome (one of the IL-1 β study pathways), inflammation, and pain. When activated, P2X7 leads to an increase in ion flow that can intensify with a decrease in the influx of cations from divalent charges such as Ca^{2+} [12]. Electrophysiology (EP) studies addressed in 2005 by Faria et al. [4] about the high pore conductance of the P2X7 receptor tested on 2BH4 and peritoneal macrophage cells observed unitary conductance values corresponding to 400 pS. In addition to these data, experiments using high-molecular-weight cationic and anionic fluorescent dyes also confirm the high pore conductance. Regarding the permeabilization through the pore, fluorophores tested were able to penetrate the cell regardless of its load, ethidium bromide (1^+), propidium iodide (2^+), and Lucifer Yellow (2^-).

However, other channels in the literature also correlate with the pore of the P2X7 receptor showing high conductance and permeability to fluorophores through the membrane. As highlights are the TRP receptors as TRPV 1–4, with notoriety for TRPV1, which is the most studied subtype of TRPs. This subtype is expressed in several cells of the organism such as the liver, lung, and nociceptive neurons. TRPV1 is a nonselective cationic channel with high permeability for Ca^{2+} , with a unitary conductance of 35–77 pS [13, 14]; however, the unitary conductance of the pore formed was not record until now. Although the ionic channel may be activated for distinct stimuli (capsaicin, low pH, and heat), this receptor only permits the fluorescent dye uptake after capsaicin activation and low pH conditions. Other TRP subtypes related to high conductance channel formation are the transient potential ankyrin type-1 (TRPA1) cationic channels with a conductance of 87–100 pS and transient potential melastin receptor 8 (TRPM8) with a conductance of 40–83 pS, according to [14, 15]. Pannexin-1 [16, 17], connexin hemichannel [18], calcium homeostasis modulator 1 (CALHM1) [19], voltage-dependent anion channel (VDAC1 highlighted) [6, 20], maxi anion [21], P2X (P2X2 and P2X4) [11], and maitotoxin-induced pore [22] are some other different types of channels with unitary conductance values greater than 200 pS and permeable to molecules of until 1000 Da [4, 10, 23]. More details about these pore-forming proteins and some techniques used to record their functionality are shown in **Table 1**.

Pore-forming proteins	EF	PLP	FC	FPR	CA	CPA
P2X7R	[4]	[24]	[25]	[26]	[26]	[4]
TRPA1	[27]	[28]	N/A	[27]	[27]	[27]
Maxi anion	[17]	N/A	N/A	[17]	[17]	[17]
TRPV1	[15]	[59]	N/A	[71]	[24]	[15]
Plasma VDAC	[29]	[30]	N/A	N/A	[29]	[29]
Connexin	[31]	[32]	[33]	[31]	[31]	[31]
Pannexin	[34]	[16]	[35]	[17]	[17]	[17]
Maitotoxin	[22]	[36]	N/A	[22]	[37]	[37]
CALHM1	[31]	N/A	[38]	[31]	[31]	[31]
TRPM8	[39]	[40]	[41]	N/A	[40]	[42]

Table taken from Ref. [49] and adapted.

Table 1.
Pore-forming proteins and some assessment techniques as electrophysiology (EP), planar lipid bilayer (PLP), flow cytometry (FC), fluorescent plate reader (FPR), colorimetric assay (CA), and cell permeabilization assay (CPA) for functional detection of the high conductance.

Although they have similarities to conductance, these different channels have particularities related to biophysical properties, having different functionalities and mechanisms. Some channels exhibit a distinct selectivity for certain ions, some having preference for anions such as Cl^- , PO_4^{3-} , and glutamate in the case of VDAC1 and maxi anion or cations like Ca^{2+} for CALHM1 [6, 19, 43].

These channels are the subject of studies for their involvement in several cellular processes, such as intracellular signaling response, release of cytokines, inflammation, cell death, pain, tumors, and metabolism regulation. Interestingly, the same functions are performed for these large conductance channels with distinct cell types or tissues. We can highlight the role of TRPs (TRPV1 and TRPM8), P2X7 receptor, connexin 43, and pannexin-1 [11, 44–48].

Based on the study of these channels, several molecular and cellular biology techniques are employed to obtain the best understanding of the functions that these receptors perform. Among these assays, the cell death assay, flow cytometry, intracellular calcium quantification, cytokine assay, and cell permeabilization stand out. However, for the best performance of these techniques, fluorescent dyes are used according to their physical-chemical characteristics and molecular mass, serving as a strategy to obtain better results and performance of experimental tests [49]. For additional information about some of the techniques mentioned above, more details involving their respective methodologies can be found in the reviews published below [50–52].

3. Fluorescent dyes and their use in the study of pores by ion channels

Fluorescence is a kind of luminescence in which the electron can capture energy by entering an excited state for a short period and returning to its normal state by emitting light. The first discoveries involving this form of light were in the middle of the sixteenth century by physician and botanist Nicolás Monardes through a wood (*Lignum nephriticum*) from Mexico used to treat kidney diseases [53, 54]. Over the years, the term “fluorescence” was coined by physicist George Gabriel Stokes, in addition to presenting a list of fluorescent substances, confirming new

advances in the area. With the advancement of science and the years, the first fluorescent dye was synthesized, fluorescein, an anionic dye produced by German chemist Adolf von Baeyer, a molecule that would be used in several laboratory methods in the future [49, 55].

Fluorescent dyes are widely used in various specialties within science, among them; the staining of different structures in tissues and cells stands out. A vast number of laboratory techniques of molecular and cellular biology in broad fields of research use these molecules, based on their chemical and physical properties according to the method worked, thus being an excellent resource in the investigation and visualization of the topic addressed [56, 57]. Among many techniques, the opening function of the high conductance pores mediated through ion channels can be evaluated in experimental tests for the capture of intracellular dye [4, 49]. Many dyes are impermeable to the membrane due to their molecular weight and physical characteristics such as positive charge (cationic dyes) and negative charge (anionic dyes). However, due to the opening of the pores with a subsequent increase in permeability, it is possible to investigate these dyes through the capture by fluorescence reading. The interaction of fluorescent dyes in the intracellular medium occurs according to the electrical charge of the target structure to be marked. Several structures that make up the intracellular environment, such as ions, nucleic acids, and organelles, each are having a specific electrical character, thus enabling the connection with fluorophores after permeating through the high conductance pores [1, 58].

In 1987, one of the first works was carried out with the use of fluorescent dyes to determine pore permeability, in which there were significant results about the subject. Steinberg et al. [58] in a study using macrophages of the J774 strain found that the presence of extracellular ATP was able to induce increased cation flow through an unidentified membrane receptor. In order to better analyze the P2X7 receptor permeability, five fluorescent anionic dyes impermeable to the membrane with varying were tested. The authors tested Lucifer Yellow (457 Da, 2^-) being the Ca^{2+} pentavalent Fura-2 indicator (832 Da, 5^-), 6-carboxyfluorescein (376 Da, 2^-), trypan blue, and Evans blue with similar charge and molecular weight (960.81 Da, 4^-). The first three dyes showed permeability through the influence of ATP. However, trypan blue and Evans blue were not permeable to the cell and may suggest a specific conductance by the receptor through the different molecular weights of the fluorophores used by the authors. This data led to the exploration and better understanding of P2X pores, mainly from the P2X7 receptor [11, 12]. However, the data about the P2X7 receptor pore permeability to cationic fluorescent dyes with charge higher 2^+ (propidium iodide) was not investigated.

In another study, Faria et al. [59] evaluated the dye uptake involving proteins P2X7, TRPV1, TRPA1, and maxi anion in astrocytes. The selectivity of these pores for cationic dyes (ethidium bromide and propidium iodide) and anionics (fluorescein and calcein) was measured. The TRPV1 and TRPA1 receptors were selective for cationic ones allowing their permeability and maxi anion for anionic ones, suggesting a possible preference of these pores concerning the electric charge of the fluorophore. This aspect did not occur with the P2X7 receptor, already reported, and another study about the subject [4]. However, the data about the maximal size of cationic dyes able to permeate for TRPV1 and TRPA1 channels or anionic for maxi anion channels was not investigated. Additionally, the permeability to compounds with the higher charge was not discussed.

One of the points to be noted regarding the capture of dyes in the investigation of pore-forming proteins is the low diversity of fluorophores used in these tests. In

most experiments, “standard” dyes are used, among which are the cationic ethidium bromide (394 Da, 1⁺), propidium iodide (668.39 Da, 2⁺), YO-PRO-1 (629 Da, 2⁺), and DAPI (350.25 Da, 2⁺) which mostly interact with nucleic acids. Anionics such as Lucifer Yellow (457 Da, 2⁻) Fluo-4 (736.63 Da, 4⁻), Fura-2 (832 Da, 5⁻), and fluorescein (332.31 Da, 1⁻) are usually ion markers or intracellular structures [27, 59, 60]. A study by McCoy et al. [42] on the permeabilization of large molecules mediated by the TRPM8 receptor, the cationic dye PO-PRO-3 (605.30 Da, 2⁺), was used to measure the dilation of the pore. However, it is not common to use fluorophores other than those mentioned above, having a great scenario to be explored in ionic conductance tests. Another impacting factor regarding dye testing is related to the electrical charge in which the molecule stands out. As seen previously, mainly in cationic dyes, the charges vary between the charge of 1⁺ and 2⁺, a fact little investigated, and the search for tests in dyes with more tri or tetravalent charges is relevant in the evaluation of these pores. However, pores similar to dyes also have charges, suggesting varied possibilities for molecular interaction of pores with the physicochemical properties of fluorophores, a fact that is not thoroughly investigated in scientific researches [49].

We consider it relevant to emphasize that some of the pore-forming proteins mentioned have no reports of the use of fluorescent dyes such as pl-VDAC. Other proteins possess basically the same ones studied at the P2X7 receptor, such as pannexin-1, maitotoxin, and maxi anion [22, 34, 59].

The only exception would be connexins, such as connexin 43, which is due to its prominent regulatory function in several vital processes, such as cardiac synchronism. However, these proteins have too much variability of homotypic and heterotypic channels with unitary conductance and permeability by size and charge of the variable molecule. Thus, the biophysical characterization of these channels can be models to generate protocols for the study of the dyes above in the other high unit conductance channels [61].

There is a plurality of fluorophores described in the literature still used in the evaluation of pore formation functionality [62], given a large field to be explored and researched regarding the use of these dyes in order to understand more clearly some specific characteristics of these pores. The following tables (**Tables 2 and 3**) contain a total of 24 different examples of fluorescent dyes with varied molecular weights, structures, and different charges (mono-, di-, tri-, or tetravalent), cutoff, within their physical-chemical characteristics, in which can be tested in future experiments to deepen the understanding about the biophysical properties of these high ion conductance channels. We hope that the data contained in these tables can contribute to the purpose of studies of the pore-forming channels, which may serve as a tactic and add discoveries to the scientific field, following more knowledge of cell physiology.

A detail to note is that the table corresponding to cationic lacks more examples of tetravalent dyes. The proteins addressed in the previous topic are reported to allow the permeabilization of molecules of up to 1000 Da. However, there are other types of tetravalent fluorophores found in the literature that exceed the permeable molecular weight limit for these pores [2]. BoBo 1 (1202.66 Da), BoBo 3 (1254.73 Da), ToTo-1 (1302.78 Da), and ToTo 3 (1354.85 Da) are some other examples of charged dyes (4⁺) [62]. Thus, the lack of other viable tetravalent dyes to the corresponding permeability values of the described proteins prevents the exploration and evaluation of the pores with these molecules. Therefore, we consider it necessary to investigate other fluorescent dyes with this type of charge, aiming at possible future experiments on the molecular pore interaction with this level of net charge [56, 63].

Anionic dyes	MW	Charge	Molecular structure
PromoFluor 840	788.96	1 ⁻	<p>The chemical structure of PromoFluor 840 is a complex molecule. It features a central xanthene core. One ring of the xanthene is substituted with a diethylamino group (-N(CH₂CH₃)₂) and a trimethylsilyl group (-Si(CH₃)₃). The other ring is substituted with a carboxylic acid group (-COOH) and a sulfonate group (-SO₃⁻). The central carbon of the xanthene is also substituted with a sulfonate group (-SO₃⁻) and a sodium ion (Na⁺).</p>
Yakima Yellow	654.30	1 ⁻	<p>The chemical structure of Yakima Yellow is a xanthene derivative. It features a central xanthene core. One ring is substituted with a methyl group (-CH₃) and a chlorine atom (-Cl). The other ring is substituted with a chlorine atom (-Cl) and a hydroxyl group (-OH). The central carbon of the xanthene is substituted with a carboxylic acid group (-COOH) and a sodium ion (Na⁺).</p>
5-Carboxyfluorescein	376.32	1 ⁻	<p>The chemical structure of 5-Carboxyfluorescein is a xanthene derivative. It features a central xanthene core. One ring is substituted with a carboxylic acid group (-COOH). The other ring is substituted with a hydroxyl group (-OH). The central carbon of the xanthene is substituted with a hydroxyl group (-OH).</p>
PromoFluor 680	852.96	2 ⁻	<p>The chemical structure of PromoFluor 680 is a complex molecule. It features a central xanthene core. One ring of the xanthene is substituted with a diethylamino group (-N(CH₂CH₃)₂) and a trimethylsilyl group (-Si(CH₃)₃). The other ring is substituted with a carboxylic acid group (-COOH) and a sulfonate group (-SO₃⁻). The central carbon of the xanthene is also substituted with a sulfonate group (-SO₃⁻) and two sodium ions (2Na⁺).</p>
JOE	602.34	2 ⁻	<p>The chemical structure of JOE is a xanthene derivative. It features a central xanthene core. One ring is substituted with a chlorine atom (-Cl) and a hydroxyl group (-OH). The other ring is substituted with a chlorine atom (-Cl) and a methoxy group (-OCH₃). The central carbon of the xanthene is substituted with a carboxylic acid group (-COOH) and a sodium ion (Na⁺).</p>

Anionic dyes	MW	Charge	Molecular structure
Dichlorofluorescein	401.2	2 ⁻
PromoFluor 405	695.59	3 ⁻
Methoxyphenetrisulfonic acid trisulfonate (MPTS)	538.4	3 ⁻
8-Aminopyrene-1,3,6-trisulfonic acid (APTS)	523.38	3 ⁻
Tetrakis(4sulfophenyl) porphine (TSPP)	935	4 ⁻

Anionic dyes	MW	Charge	Molecular structure
Tetrakis(4carboxyphenyl) porphine (TCPP)	790.8	4 ⁻
Quin-2	693.87	4 ⁻

Table 2.
Fluorescent anionic dyes.

Cationic dyes	MW	Charge	Molecular structure
QSY9 carboxylic acid succinimidyl ester	951.43	1 ⁺
PyMPO	564.4	1 ⁺
Hexidium iodide	497.42	1 ⁺
TO-PRO-1	645.38	2 ⁺

Cationic dyes	MW	Charge	Molecular structure
JO-PRO-1	630.31	2 ⁺
BO-PRO-1	595.32	2 ⁺
EvaGreen	509.73	2 ⁺
Hoechst 769121	596.96	3 ⁺
Hoechst 33342	561.93	3 ⁺
Hoechst 34580	560.96	3 ⁺

Cationic dyes	MW	Charge	Molecular structure
Hoechst 33258	533.88	3 ⁺
EthD-1	856.77	4 ⁺

Table 3.
Fluorescent cationic dyes.

4. Conclusion

Scientific advances with methods involving fluorescence, such as fluorimetry and colorimetry, have led to the significant discoveries in different horizons of biological knowledge, with the application of fluorophores in these methodologies having an immense importance in the purpose of evaluating different topics involving the study of cells. Studies on the permeabilization of the plasma membrane through high conductance pore-forming proteins are a topic widely addressed in laboratory research, with the use of fluorescent dyes being a useful device in the evaluation and investigation of the functionality of ionotropic receptors. Through the tables presented above, several anionic and cationic dyes with different charge values (mono-, di-, tri-, and tetravalent) can be used by researchers in numerous experiments, aiming to check these pores through their physical-chemical properties and conductance. We hope that this review will motivate further research on the applicability, mechanisms, and biophysical properties of these high ion conductance channels, aiming at new knowledge and discoveries for the cellular biological perspective.

Acknowledgements

We are grateful to the Coordination of Superior Level Staff Improvement (CAPES), Finance Code 001, Oswaldo Cruz Institute (IOC), and Postgraduation of Sciences and Biotechnology (PPBI), and Federal Fluminense University (UFF).

Conflict of interest

The authors declare no conflict of interest.

Notes/thanks/other declarations

We are grateful to the Coordination of Superior Level Staff Improvement (CAPES), Finance Code 001, National Council for Scientific and Technological

Development, Foundation Carlos Chagas Filho for Research Support of the State of Rio de Janeiro, Oswaldo Cruz Institute (IOC), Postgraduation and Cellular and Molecular Biology (BCM) from IOC, and Postgraduation of Sciences and Biotechnology (PPBI) from Federal Fluminense University (UFF).

IntechOpen

IntechOpen

Author details

Guilherme Teixeira and Robson Faria*
Laboratory of Toxoplasmosis and Other Protozoans, Oswaldo Cruz Institute (IOC),
Fiocruz, Rio de Janeiro, Brazil

*Address all correspondence to: robson.xavier@gmail.com

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. Distributed under the terms of the Creative Commons Attribution - NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited.

References

- [1] Giepmans BNG, Adams SR, Ellisman MH, Tsien RY. The fluorescent toolbox for assessing protein location and function. *Science*. 2006;**312**:217-224
- [2] Petit JM, Denis-Gay M, Ratinaud MH. Assessment of fluorochromes for cellular structure and function studies by flow cytometry. *Biology of the Cell*. 1993;**78**:1-13
- [3] Adams DJ. Ionic channels in vascular endothelial cells. *Trends in Cardiovascular Medicine*. 1994;**4**:18-26
- [4] Faria RX, DeFarias FP, Alves LA. Are second messengers crucial for opening the pore associated with P2X 7 receptor? *American Journal of Physiology-Cell Physiology*. 2005;**288**:C260-C271
- [5] Hille B. Ionic selectivity of Na and K channels of nerve membranes. *Membranes (Basel)*. 1975;**3**:255-323
- [6] Shoshan-Barmatz V, Mizrachi D. VDAC1: From structure to cancer therapy. *Frontiers in Oncology*. 2012;**2**:1-34
- [7] Tominaga M, Tominaga T. Structure and function of TRPV1. *Pflügers Archiv - European Journal of Physiology*. 2005;**451**:143-150
- [8] Brogden KA, Ackermann M, McCray PB, Tack BF. Antimicrobial peptides in animals and their role in host defences. *International Journal of Antimicrobial Agents*. 2003;**22**:465-478
- [9] Liu X, Lieberman J. Knocking ‘em Dead: Pore-Forming Proteins in Immune Defense. *Annual Review of Immunology*. 2020;**38**:455-485
- [10] Alves LA, Ferreira LB, Pacheco PF, Mendivelso EAC, Teixeira PCN, Faria RX. Pore forming channels as a drug delivery system for photodynamic therapy in cancer associated with nanoscintillators. *Oncotarget*. 2018;**8**:25342-25354
- [11] Burnstock G. Purine and pyrimidine receptors. *Cellular and Molecular Life Sciences*. 2007;**64**:1471-1483
- [12] North RA. Molecular physiology of P2X receptors. *Physiological Reviews*. 2002;**82**:1013-1067
- [13] Cui M, Honore P, Zhong C, Gauvin D, Mikusa J, Hernandez G, et al. TRPV1 receptors in the CNS play a key role in broad-spectrum analgesia of TRPV1 antagonists. *The Journal of Neuroscience*. 2006;**26**:9385-9393
- [14] Ferreira LGB, Faria RX. TRPing on the pore phenomenon: What do we know about transient receptor potential ion channel-related pore dilation up to now? *Journal of Bioenergetics and Biomembranes*. 2016;**48**:1-12
- [15] Diver MM, Cheng Y, Julius D. Structural insights into TRPM8 inhibition and desensitization. *Science*. 2019;**365**:1434-1440
- [16] Mou L, Ke M, Song M, Shan Y, Xiao Q, Liu Q, et al. Structural basis for gating mechanism of Pannexin 1 channel. *Cell Research*. 2020;**30**:452-454
- [17] Pelegrin P, Surprenant A. Pannexin-1 mediates large pore formation and interleukin-1 β release by the ATP-gated P2X7 receptor. *The EMBO Journal*. 2006;**25**:5071-5082
- [18] Stout CE, Costantin JL, Naus CCG, Charles AC. Intercellular calcium signaling in astrocytes via ATP release through connexin hemichannels. *The Journal of Biological Chemistry*. 2002;**277**:10482-10488
- [19] Ma Z, Siebert AP, Cheung KH, Lee RJ, Johnson B, Cohen AS, et al.

- Calcium homeostasis modulator 1 (CALHM1) is the pore-forming subunit of an ion channel that mediates extracellular Ca^{2+} regulation of neuronal excitability. *Proceedings of the National Academy of Sciences of the United States of America*. 2012;**109**:E1963-E1971
- [20] Sabirov RZ, Dutta AK, Okada Y. Volume-dependent ATP-conductive large-conductance anion channel as a pathway for swelling-induced ATP release. *The Journal of General Physiology*. 2001;**118**:251-266
- [21] Islam MR, Uramoto H, Okada T, Sabirov RZ, Okada Y. Maxi-anion channel and pannexin 1 hemichannel constitute separate pathways for swelling-induced ATP release in murine L929 fibrosarcoma cells. *American Journal of Physiology-Cell Physiology*. 2012;**303**:C924-C935
- [22] Schilling WP, Sinkins WG, Estacion M. Maitotoxin activates a nonselective cation channel and a P2Z/P2X7-like cytolytic pore in human skin fibroblasts. *American Journal of Physiology-Cell Physiology*. 1999;**277**:C755-C765
- [23] Alves LA, De Melo Reis RA, De Souza CAM, De Freitas MS, Teixeira PCN, Neto Moreira Ferreira D, et al. The P2X7 receptor: Shifting from a low- to a high-conductance channel—An enigmatic phenomenon? *Biochimica et Biophysica Acta - Biomembranes*. 2014;**1838**:2578-2587
- [24] De Souza CAM, Teixeira PCN, Faria RX, Krylova O, Pohl P, Alves LA. A consensus segment in the M2 domain of the hP2X 7 receptor shows ion channel activity in planar lipid bilayers and in biological membranes. *Biochimica et Biophysica Acta, Biomembranes*. 2012;**1818**:64-71
- [25] Jursik C, Sluyter R, Georgiou JG, Fuller SJ, Wiley JS, Gu BJ. A quantitative method for routine measurement of cell surface P2X7 receptor function in leucocyte subsets by two-colour time-resolved flow cytometry. *Journal of Immunological Methods*. 2007;**325**:67-77
- [26] Faria RX, de Jesus Hiller N, Salles JP, Resende JALC, Diogo RT, von Ranke NL, et al. Arylboronic acids inhibit P2X7 receptor function and the acute inflammatory response. *Journal of Bioenergetics and Biomembranes*. 2019;**51**:277-290
- [27] Banke TG, Chaplan SR, Wickenden AD. Dynamic changes in the TRPA1 selectivity filter lead to progressive but reversible pore dilation. *American Journal of Physiology-Cell Physiology*. 2010;**298**:C1457-C1468
- [28] Sinica V, Zimova L, Barvikova K, Macikova L, Barvik I, Vlachova V. Human and mouse TRPA1 are heat and cold sensors differentially tuned by voltage. *Cell*. 2019;**9**:57-80
- [29] Elinder F, Akanda N, Tofighi R, Shimizu S, Tsujimoto Y, Orrenius S, et al. Opening of plasma membrane voltage-dependent anion channels (VDAC) precedes caspase activation in neuronal apoptosis induced by toxic stimuli. *Cell Death and Differentiation*. 2005;**12**:1134-1140
- [30] Böhm R, Amodeo GF, Murlidaran S, Chavali S, Wagner G, Winterhalter M, et al. The structural basis for low conductance in the membrane protein VDAC upon β -NADH binding and voltage gating. *Structure*. 2020;**28**:206-214
- [31] Gaete PS, Lillo MA, Lopez WI, Liu Y, Harris AL, Contreras JE. Novel biophysical properties of molecular permeation in CALHM1 and connexin channels. *Biophysical Journal*. 2020;**118**:270a
- [32] Carnarius C, Kreir M, Krick M, Methfessel C, Moehrle V, Valerius O,

- et al. Green fluorescent protein changes the conductance of connexin 43 (Cx43) hemichannels reconstituted in planar lipid bilayers. *The Journal of Biological Chemistry*. 2012;**287**:2877-2886
- [33] Xu C, Fan Z, Shan W, Hao Y, Ma J, Huang Q, et al. Cyclic stretch influenced expression of membrane connexin 43 in human periodontal ligament cell. *Archives of Oral Biology*. 2012;**57**:1602-1608
- [34] Iglesias R, Locovei S, Roque A, Alberto AP, Dahl G, Spray DC, et al. P2X7 receptor-pannexin1 complex: Pharmacology and signaling. *American Journal of Physiology-Cell Physiology*. 2008;**295**:C752-C760
- [35] Chekeni FB, Elliott MR, Sandilos JK, Walk SF, Kinchen JM, Lazarowski ER, et al. Pannexin 1 channels mediate “find-me” signal release and membrane permeability during apoptosis. *Nature*. 2010;**467**:863-867
- [36] Treviño CL, De La Vega-Beltrán JL, Nishigaki T, Felix R, Darszon A. Maitotoxin potently promotes Ca^{2+} influx in mouse spermatogenic cells and sperm, and induces the acrosome reaction. *Journal of Cellular Physiology*. 2006;**206**:449-456
- [37] Caillaud A, Yasumoto T, Diogène J. Detection and quantification of maitotoxin-like compounds using a neuroblastoma (Neuro-2a) cell based assay. Application to the screening of maitotoxin-like compounds in *Gambierdiscus* spp. *Toxicon*. 2010;**56**:36-44
- [38] Moreno-Ortega AJ, Buendia I, Mouhid L, Egea J, Lucea S, Ruiz-Nuño A, et al. CALHM1 and its polymorphism P86L differentially control Ca^{2+} homeostasis, mitogen-activated protein kinase signaling, and cell vulnerability upon exposure to amyloid β . *Aging Cell*. 2015;**14**:1094-1102
- [39] Kühn FJP, Knop G, Lückhoff A. The transmembrane segment S6 determines cation versus anion selectivity of TRPM2 and TRPM8. *The Journal of Biological Chemistry*. 2007;**282**:27598-27609
- [40] Asuthkar S, Velpula KK, Elustondo PA, Demirkhanyan L, Zakharian E. TRPM8 channel as a novel molecular target in androgen-regulated prostate cancer cells. *Oncotarget*. 2015;**6**:17221-17236
- [41] Zeng J, Wu Y, Zhuang S, Qin L, Hua S, Mungur R, et al. Identification of the role of TRPM8 in glioblastoma and its effect on proliferation, apoptosis and invasion of the U251 human glioblastoma cell line. *Oncology Reports*. 2019;**42**:1517-1526
- [42] McCoy DD, Palkar R, Yang Y, Ongun S, McKemy DD. Cellular permeation of large molecules mediated by TRPM8 channels. *Neuroscience Letters*. 2017;**639**:59-67
- [43] Sabirov RZ, Merzlyak PG. Plasmalemmal VDAC controversies and maxi-anion channel puzzle. *Biochimica et Biophysica Acta, Biomembranes*. 2012;**1818**:1570-1580
- [44] Hofmann F, Navarrete M, Álvarez J, Guerrero I, Gleisner MA, Tittarelli A, et al. Cx43-gap junctions accumulate at the cytotoxic immunological synapse enabling cytotoxic T lymphocyte melanoma cell killing. *International Journal of Molecular Sciences*. 2019;**20**:1-16
- [45] North RA. P2X receptors. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*. 2016;**371**:1-7
- [46] Samivel R, Kim DW, Son HR, Rhee YH, Kim EH, Kim JH, et al. The role of TRPV1 in the CD4+ T cell-mediated inflammatory response of allergic rhinitis. *Oncotarget*. 2016;**7**:148-160

- [47] Valdebenito S, Barreto A, Eugenin EA. The role of connexin and pannexin containing channels in the innate and acquired immune response. *Biochimica et Biophysica Acta, Biomembranes*. 2018;**1860**:154-165
- [48] Zhang L, Barritt GJ. TRPM8 in prostate cancer cells: A potential diagnostic and prognostic marker with a secretory function? *Endocrine-Related Cancer*. 2006;**13**:27-38
- [49] Ferreira L, Pereira L, Faria R. Fluorescent dyes as a reliable tool in P2X7 receptor-associated pore studies. *Journal of Bioenergetics and Biomembranes*. 2015;**47**:283-307
- [50] Gee KR, Brown KA, Chen WNU, Bishop-Stewart J, Gray D, Johnson I. Chemical and physiological characterization of fluo-4 Ca^{2+} -indicator dyes. *Cell Calcium*. 2000;**97**:97-106
- [51] Kepp O, Galluzzi L, Lipinski M, Yuan J, Kroemer G. Cell death assays for drug discovery. *Nature Reviews Drug Discovery*. 2011;**10**:221-237
- [52] Picot J, Guerin CL, Le Van Kim C, Boulanger CM. Flow cytometry: Retrospective, fundamentals and recent instrumentation. *Cytotechnology*. 2012;**64**:109-130
- [53] Lakowicz JR. Principles of Fluorescence Spectroscopy. 3rd ed. Boston, MA: Springer; 2006
- [54] Valeur B, Berberan-Santos MN. A brief history of fluorescence and phosphorescence before the emergence of quantum theory. *Journal of Chemical Education*. 2011;**88**:731-738
- [55] Baeyer A. Ueber eine neue Klasse von Farbstoffen. *Berichte der Dtsch Chem Gesellschaft*. 1871;**4**:555-558
- [56] Khadria AS, Senes A. Fluorophores, environments, and quantification techniques in the analysis of transmembrane helix interaction using FRET. *Biopolymers*. 2015;**104**:247-264
- [57] Lemasters JJ, Ramshesh VK. Imaging of mitochondrial polarization and depolarization with cationic fluorophores. *Methods in Cell Biology*. 2007;**80**:283-295
- [58] Steinberg TH, Newman AS, Swanson JA, Silverstein SC. ATP4-permeabilizes the plasma membrane of mouse macrophages to fluorescent dyes. *The Journal of Biological Chemistry*. 1987;**262**:8884-8888
- [59] Faria RX, Reis RAM, Ferreira LGB, Cezar-de-Mello PFT, Moraes MO. P2X7R large pore is partially blocked by pore forming proteins antagonists in astrocytes. *Journal of Bioenergetics and Biomembranes*. 2016;**48**:309-324
- [60] Rissiek B, Stabernack J, Cordes M, Duan Y, Behr S, Menzel S, et al. Astrocytes and microglia are resistant to NAD^{+} -mediated cell death along the ARTC2/P2X7 axis. *Frontiers in Molecular Neuroscience*. 2020;**12**:1-8
- [61] Johnson RG, Le HC, Evenson K, Loberg SW, Myslajek TM, Prabhu A, et al. Connexin hemichannels: Methods for dye uptake and leakage. *The Journal of Membrane Biology*. 2016;**249**:719-741
- [62] Sabnis RW. Handbook of Fluorescent Dyes and Probes. Hoboken, New Jersey: John Wiley & Sons; 2015
- [63] Stockert JC, Blazquez-Castro A. Fluorescence Microscopy in Life Sciences. Sharjah, UAE: Bentham Science Publishers; 2017