We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

An Insight into the Responses of Early-Maturing *Brassica napus* to Different Low-Temperature Stresses

Xin He

Abstract

Rapeseed (*Brassica napus* L.) is an important oil crop worldwide, responds to vernalization, and shows an excellent tolerance to cold stresses during vegetative stage. The winter-type and semi-winter-type rapeseed were typical winter biennial plants in Europe and China. In recent years, more and more early-maturing semi-winter rapeseed varieties were planted across China. Unfortunately, the early-maturing rapeseed varieties with low cold tolerance have higher risk of freeze injury in cold winter and spring. The molecular mechanisms for coping with different low-temperature stress conditions in rapeseed recently had gained more attention and development. The present review gives an insight into the responses of early-maturing *B. napus* to different low-temperature stresses (chilling, freezing, cold-acclimation, and vernalization), and the strategies to improve tolerance against low-temperature stresses are also discussed.

Keywords: *Brassica napus*, low-temperature, early-maturing

1. Introduction

Low-temperature is a major environmental stress that adversely affects plant growth and development, limiting the productivity and regional distribution of crops [1, 2]. Rapeseed is an important oil crop worldwide, with planting area of 37.58 million hectares producing 75.00 million tons of oilseeds in 2018 (<http://www.fao.org/faostat/>). Based on vernalization requirement, rapeseed is divided into three main ecotypes, i.e., winter, semi-winter and spring types [3–5]. The winter type rapeseed is mainly grown in Europe and is sown in late summer, which requires strong vernalization and flowerings in spring, exhibiting a classical winter annual and with excellent cold tolerance during vegetative stage [3, 6]. The semi-winter type rapeseed is mainly grown in China only needs moderate or weak vernalization to promote flowering in spring, and with weak cold tolerance [3, 7]. The semi-winter type rapeseed excessive exposure to low temperature stress in winter will lead to plant damage at vegetative stage and finally cause yield loss [8]. Yangtze River basin is the major region for planting semi-winter rapeseed in China, which accounts for at least 90% of the nation's total production [9]. The semi-winter rapeseed is usually sown in late September and early October shortly after the harvest of rice, and harvested in May before the cropping of rice in this area [10].

However, in recent years, due to the delay of rice harvest which leads to the postpone of rapeseed sowing until late October or early November, therefore, more and more early-maturing semi-winter rapeseed varieties were planted across Yangtze River basin. Unfortunately, the early-maturing rapeseed varieties with low cold tolerance have higher risk of freeze injury in cold winter and spring [11]. Hence, it is vital to compare early-maturing rapeseed varieties tolerant to cold and evaluate molecular mechanisms that adapt to different low-temperature stress conditions.

2. Morphophysiological mechanism of rapeseed in responses to low-temperature stress

Cold (low-temperature) stress included chilling stress ($>0^{\circ}\text{C}$) and freezing stress ($<0^{\circ}\text{C}$) [12]. Chilling stress ($0\text{--}15^{\circ}\text{C}$) causes the membrane to rigidify, destabilizes protein complexes and impairs photosynthesis, eventually made plant stop growing, whereas freezing stress ($<0^{\circ}\text{C}$) causes intracellular and extracellular ice crystal formation, and results in mechanical injury, and plant death [13–15].

Despite the fact that winter and semi-winter rapeseed is an overwintering oil crop, cold stress can still affect rapeseed development and ultimately lead to a decrease in production [8, 11]. The suitable temperature scope is $10\sim 20^{\circ}\text{C}$ for the growth of winter and semi-winter rapeseed. The rapeseed flower number was reduced below 10°C and the rapeseed flowering was arrested when the temperature decreased to 5°C . The rapeseed growth was arrested below 3°C and rapeseed leaves was injured below 0°C [8]. The delay of rapeseed sowing results in poor germination [16], decreased seedling biomass [17, 18], delay of floral initiation and floral bud differentiation processes [17, 19], and decreased flower number, effective pod number, pod length, and seed yield [17, 20, 21] due to low-temperature stress. In January 2008, South China was exposed to an extremely ice-frozen weather, which caused serious injuries to winter rape, affected 77.8% of the overall winter rape area in China and resulted in 10.9% yield losses [22]. Due more and more early-maturing semi-winter rapeseed varieties were planted across Yangtze River basin, rapeseed faces increased risks from continuous low temperature overcast and rainy weather in March. Continuous low temperature overcast and rainy weather during the rapeseed flowering stage or after flowering decreased the ratio of effectual silique, seeds per silique and oil content [23]. In March and April 2010, the middle and lower reaches of the Yangtze River region were exposed to continuous low temperature overcast and rainy weather, which resulted in 10–20% yield losses [23].

To date, many studies have investigated the morphological and physiological changes of low-temperature stressed rapeseeds. Leaves are the main organ to perceive low temperature stress and transmit stress signal in plants [24]. The morphological changes (dehydrated and wilting) of leaves became increasingly evident with the decrease of temperature, due to the total water content in leaves of rapeseed decreased [25, 26].

In winter rapeseed, prolonged cold acclimation led to increased thickness of young leaf blades and leaf cell walls, modified dimensions of mesophyll cells, numerous invaginations of plasma membranes and large phenolic deposits in chloroplasts, large vesicles or cytoplasm/tonoplast interfaces [27, 28]. Unlike cold acclimation, transient freezing treatment reduced the thickness of leaf cell walls and phenolic aggregates, caused reversible disorganization of the cytoplasm and chloroplasts swelling [27, 28]. Obvious gaps existed in the chloroplast grana and starch grains increased in quantity and volume [25]. In general, cold-tolerant winter rapeseed usually grows slowly, having small thick creeping deep-green waxy leaves and large root system.

Low temperature-induced thermodynamic constraints on carbon metabolism was the primary reason for lower photosynthetic activity in plants [24]. Photosynthetic efficiency is a good indicator for Low temperature tolerance in plants [10]. Just like in other crop plants, a marked reduction of photosynthetic activity is observed in rapeseed leaves when treated with low temperature [24, 29]. Though the photosynthetic activities were reduced both in the cold-stressed leaves of cold-tolerant and cold-sensitive rapeseed cultivars, the chlorophyll a, chlorophyll b and photosynthetic activities in the young leaves of cold-tolerant cultivar all were higher than that in cold-sensitive cultivar [24].

Simultaneously, low-temperature stress caused the overproduction of reactive oxygen species (ROS), elevated H_2O_2 level and increased malondialdehyde (MDA) content in plants, which leads to a necrosis of plants. Plants possess an effective antioxidant system includes superoxide dismutase (SOD), ascorbate peroxidase (APX), peroxidase (POD) and catalase (CAT) enzymes, whose combined activities play an important role in elimination of destructive effects of ROS [24, 30]. Furthermore, under natural cold stress in field, the proline, soluble sugar, soluble protein, MDA contents and SOD, POD, CAT activities changed obviously in functional leaves of rapeseed. CAT and SOD activity reached the highest when temperature dropped to 5 and 3°C, respectively. The proline and soluble sugar contents increased when mean daily temperature decreased to 5°C and reached the maximum when temperature was below 0°C. The contents of soluble protein and MDA showed a trend to decrease at first and then increase when mean daily temperature dropped to 10, 5 and 0°C [30]. The SOD and APX activities were both increased by low temperature in the young leaves of cold-tolerant rapeseed cultivar. However, the APX activity was decreased by low temperature in the young leaves of cold-sensitive rapeseed cultivar. While, in the cold-stressed mature leaves, both cold-tolerant and cold-sensitive rapeseed cultivars represented similar antioxidant capacities [24].

Under chilling and freezing stress, the increment of proline accumulation, soluble sugar and protein contents were enhanced in cold-tolerant cultivar compared with cold-sensitive cultivar [24, 26]. Leaf abscisic acid (ABA) was enhanced in cold-tolerant cultivar under chilling and freezing stress [26].

3. Molecular mechanisms influencing responses to different low-temperature stresses in rapeseed

Plants showed increased freezing tolerance during exposure to chilling and low nonfreezing temperatures in a phenomenon known as cold acclimation [31]. The molecular mechanism of cold acclimation and cold tolerance in *Arabidopsis* and winter cereals has been extensively studied. Cold acclimation is a very complex trait involving an array of physiological and biochemical modifications, and these altered processes involved changes in gene expression patterns via phytohormone and the ICE (Inducer of CBF Expressions)-CBF (C-repeat binding factors)-COR (cold-responsive) signaling pathway [13, 32, 33].

3.1 ICE-CBF-COR signaling

In most plant species, CBF transcription factors could bind directly to the promoters of *COR* genes and induce the expression of *COR* genes [34–36]. The *COR* genes protected plant cells against cold-induced damage, repaired cold-rigidified membranes and stabilized cellular osmotic potential by encoding cryoprotective proteins and key enzymes for osmolyte biosynthesis [37]. In *Arabidopsis*, the basic

helix-loop-helix transcription factor ICE1/2 were induced by cold stress, could bind directly to the promoters of CBF and induced CBF expression under low-temperature stress [38–40].

Similar as other plants, the expression of CBF and COR genes were induced by chilling and freezing stresses in different ecotypes rapeseed with different cold tolerance [11, 36, 41–45]. CBFs (*BnaAnng34260D/BnaCnng49280D/BnaC03g71900D/BnaC07g39680D*), *Kin1* and COR15 all were upregulated in both winter and semi-winter ecotype rapeseeds after cold stress (4°C for 7 days), while *BnaA08g30910D* (a CBF-like gene) and *BnCOR25* were not varied in expression in any cold stressed rapeseeds [41]. Furthermore, COR15A/B, COR413-PM1 and nine CBF1/2/4 genes all were strongly upregulated in winter and spring ecotype rapeseeds after cold stress (4°C for 12 h) [42]. While *ICE1* and COR413-PM2 was downregulated in both winter and spring ecotypes after cold stress, two *CBF3* genes were not induced by cold stress [42]. Most of *COR15A* and *COR15B* were significantly induced in both cold-tolerant and cold-sensitive semi-winter early-maturing rapeseeds after chilling (4°C for 12 h) and freezing (–4°C for 12 h) stress, regardless of cold acclimation (4°C for 14 days) [11]. Ten of 12 *CBF* genes were strongly induced by freezing stress more than chilling stress, regardless of cold acclimation [11]. Unlike other CBFs, *BnaC03g71900D* was induced by chilling stress more than freezing stress and *BnaA03g13620D* was induced by freezing stress but suppressed by cold acclimation [11]. However, two *ICE1* genes were not induced by chilling and freezing stress [11], as opposed to *AtICE1* [38]. Inexplicably, no CBF genes were identified in responsive to freezing stress in freezing-tolerant rapeseed as reported by Pu [25].

BnCOR25 were significantly induced by cold and osmotic stress treatment in rapeseed, overexpression of *BnCOR25* in *Arabidopsis* enhances plant tolerance to cold stress [46]. Overexpression of two rapeseed CBF-like transcription factors *BnCBF5* and *BnCBF17* in spring rapeseed resulted in increased constitutive freezing tolerance, increased photochemical efficiency and photosynthetic capacity [29]. However, constitutively overexpressing *BNCBF5/17* in rapeseed resulted in various degrees of dwarf habit and longer time to flower [29]. The multi-gene (*NCED3*, *ABAR*, *CBF3*, *LOS5*, and *ICE1*) transgenic rapeseed plants exhibited pronounced growth advantage under both normal growth and stress conditions [47].

3.2 ABA signaling

Absciscic acid (ABA) is a vital plant hormone that plays a key role in stress resistance during plant growth and development [48–50]. It was reported that ABA levels are increased after cold stress in plants and exogenous application of ABA can induce plant cold tolerance [11, 51, 52]. *OST1/SnRK2E*, a serine-threonine protein kinase in ABA core signaling pathway, acted upstream of CBFs to positively regulate freezing tolerance via phosphorylating *ICE1* to prevent its 26S proteasome-mediated degradation by *HOS1* [53]. *OST1* phosphorylated basic transcription factors 3 (*BTF3*) and *BTF3*-like factors, and facilitated their interactions with CBFs to promote CBF stability under cold stress [54].

27 ABA biosynthesis genes (nine-*cis*-epoxycarotenoid dioxygenase (*NCED3/4/5/9*), ABA DEFICIENT 1/2 (*ABA1/2*), abscisic aldehyde oxidase 1/2/3 (*AAO1/2/3*) and carotenoid cleavage dioxygenase 1 (*CCD1*)) were regulated by cold stress in winter and/or spring rapeseed. Additionally, many genes involved in ABA signal transduction, such as ABA INSENSITIVE 1/5 (*ABI1/5*), ABA-responsive element binding protein 3 (*AREB3*), ABA responsive element-binding factor 1/2/3 (*ABF1/2/3*), highly ABA-induced PP2C gene 1/2/3 (*HAI1/2/3*), OPEN STOMATA 1 (*OST1*), *PYL1*-like 4/6/7/10 (*PYL4/6/7/10*), regulatory component of ABA receptor 1/3 (*RCAR1/3*), *SNF1*-related protein kinase 2.2/2.5/2.7/2.10 (*SnRK2.2/2.5/2.7/2.10*),

all were differentially expressed in winter and/or spring rapeseed after cold stress [42]. Furthermore, in freezing-treated (-2°C) leaves of cold-tolerant winter rapeseed line 2016TS(G)10, one *PYL* gene and one *ABI5*-like gene were up-regulated, while 13 *PP2C* and 4 *ABI5*-like genes were down-regulated [25]. 72.8% ABA signaling genes (94/129) were regulated by chilling and/or freezing in both cold-tolerant and cold-sensitive rapeseed plants but they presented different expression profiles [11]. The ABA receptors *PYL5/7* genes were both induced by cold and freezing stress, while the *PYL1/9* genes were suppressed. The *PYR1/4* genes were only induced by chilling stress but not by freezing stress. The *PYL6* were induced by chilling stress and freezing stress following cold acclimation. ABA co-receptor *ABI1* and *HAB1* were suppressed by all low-temperature treatments, while *HAB2* was upregulated. The *SnRK2B* and *SnRK2D* were induced by all low-temperature treatments, while *SnRK2C* was suppressed. *SnRK2F* and one *ABI5* were both induced only by freezing treatment. The *OST1* was induced only in cold-tolerant but not in cold-sensitive rapeseed [11]. While 13 *ABI5*-like genes have a complex expression pattern in response to different low-temperature conditions, including up-regulated, down-regulated and no changed. Exogenous application of ABA significantly improved the rapeseed seedlings freezing tolerance [11]. Overexpression of *BnaABI3* leads to improved embryo degreening following frost exposure and enhanced pod strength in rapeseed [55].

3.3 Ca^{2+} signaling

Calcium (Ca^{2+}) is an important second messenger of signal transduction in the plant stress responses, plant growth and development. Ca^{2+} signaling were detected and transmitted by calmodulin/calmodulin-like proteins (CaM/CML), calcium-dependent protein kinase (CDPK) and calcineurin B-like proteins (CBLs) [56, 57]. The level of cytosolic Ca^{2+} was transiently increased in plants under cold stress [57–59]. In rice, *COLD1* interacts with the G-protein α subunit and activates the Ca^{2+} channel, results the increment of expression of CBF under low-temperature stress [60]. In *Arabidopsis*, overexpression of CaM3 repressed the expression of *COR* genes (*RD29A*, *KIN1* and *KIN2*) [61]. CaM-binding transcription activators (CAMTAs) bind to the promotor of CBF2, promoting CBF2 expression and plant freezing tolerance [62, 63].

88 of 129 *CaM/CML* genes were regulated by cold stress in semi-winter rapeseed cultivar ZS11 [64]. 91 of 129 *CaM/CML* were regulated by chilling and/or freezing stress in semi-winter early-maturing rapeseed, and most of them were strongly induced by freezing stress [11, 64]. Additionally, 22 rapeseed CDPK genes were up-regulated by freezing treatments [11]. Furthermore, there were 91 genes involved in Ca^{2+} signaling (35 CDPK, 38 *CaM*, 16 *CIPK* and 2 Ca^{2+} -ATPase) were differentially expressed in winter rapeseed after cold stress, while 79 genes (44 CDPK, 19 *CaM*, 15 *CIPK* and 1 Ca^{2+} -ATPase) were differentially expressed in spring rapeseed after cold stress [42]. In Zhang's report, there were 5 CBL, 7 *CIPK* and 7 CDPK genes were regulated by cold treatment [65, 66].

4. Future directions

Rapeseed is one of the most important oil crops in the world and China and is affected by chilling and freezing stress. In recently years, several studies have tried to identify the main signaling pathways and genes responsible for low-temperature stress (chilling and/or freezing; cold acclimation and/or cold shock) in different rapeseeds (winter, semi-winter and spring type; cold-sensitive and cold-tolerant;

late maturing and early maturing) based on transcriptomics, metabolomics, lipidomics, and QTL analyses [11, 25, 41, 42, 45, 67–69]. Though there were so many candidate genes involved in the response to low-temperature stress have been identified, only few genes' functions in cold tolerant have been tested and verified in rapeseed [10, 29, 55, 70, 71]. It is a pity that constitutive overexpression of rapeseed BnCBF5 and BnCBF17 resulted in various degrees of dwarf habit and longer time to flower, though which resulted in increased freezing tolerance remarkably in spring rapeseed “Westar” [29]. There is still much work to be performed to understand rapeseed plants' responses to low-temperature stress and breed cold-tolerant rapeseed.

Genome editing is an efficient approach for crop improvement either by loss or gain of gene function and several different strategies have been developed [72]. Though there were a few studies using CRISPR/Cas9 system for editing genes associated with plant/pod development, fatty acid synthesis and biotic stress response [72], no application of CRISPR-Cas9 for editing genes involved in chilling and freezing tolerant in rapeseed. It is expected that the newly emerging genome editing system will make a contribution to future gene function research and molecular design breeding in cold-tolerant rapeseed.

Author details

Xin He^{1,2,3}

1 Southern Regional Collaborative Innovation Center for Grain and Oil Crops in China, Hunan Agricultural University, Changsha, Hunan, China

2 Oil Crops Research, Hunan Agricultural University, Changsha, Hunan, China

3 Hunan Branch of National Oilseed Crops Improvement Center, Changsha, Hunan, China

*Address all correspondence to: hexinzsh@126.com

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Liu J, Shi Y, Yang S. Insights into the regulation of C-repeat binding factors in plant cold signaling. *Journal of Integrative Plant Biology*. 2018;**60**(9):780-795
- [2] Ding Y, Shi Y, Yang S. Advances and challenges in uncovering cold tolerance regulatory mechanisms in plants. *The New Phytologist*. 2019;**222**(4):1690-1704
- [3] Lu K, Wei L, Li X, Wang Y, Wu J, Liu M, et al. Whole-genome resequencing reveals *Brassica napus* origin and genetic loci involved in its improvement. *Nature Communications*. 2019;**10**(1):1154
- [4] Chalhoub B, Denoeud F, Liu S, Parkin IA, Tang H, Wang X, et al. Plant genetics. Early allopolyploid evolution in the post-Neolithic *Brassica napus* oilseed genome. *Science*. 2014;**345**(6199):950-953
- [5] Wei D, Cui Y, He Y, Xiong Q, Qian L, Tong C, et al. A genome-wide survey with different rapeseed ecotypes uncovers footprints of domestication and breeding. *Journal of Experimental Botany*. 2017;**68**(17):4791-4801
- [6] O'Neill CM, Lu X, Calderwood A, Tudor EH, Robinson P, Wells R, et al. Vernalization and floral transition in autumn drive winter annual life history in oilseed rape. *Current Biology: CB*. 2019;**29**(24):4300-4306 e4302
- [7] Wu D, Liang Z, Yan T, Xu Y, Xuan L, Tang J, et al. Whole-genome resequencing of a worldwide collection of rapeseed accessions reveals the genetic basis of ecotype divergence. *Molecular Plant*. 2019;**12**(1):30-43
- [8] Lei Y, Tariq S, Yong C, Yan LÜ, Xue-Kun Z, Xi-ling Z. Physiological and molecular responses to cold stress in rapeseed (*Brassica napus* L.). *Journal of Integrative Agriculture*. 2019;**12**:60345-60347
- [9] Tian Z, Ji YH, Sun LX, Xu XL, Fan DL, Zhong HL, et al. Changes in production potentials of rapeseed in the Yangtze River Basin of China under climate change: A multi-model ensemble approach. *Journal of Geographical Sciences*. 2018;**28**(11):1700-1714
- [10] Huang Y, Hussain MA, Luo D, Xu H, Zeng C, Havlickova L, et al. A *Brassica napus* reductase gene dissected by associative transcriptomics enhances plant adaption to freezing stress. *Frontiers in Plant Science*. 2020;**11**:971
- [11] Xin H, Xianchao N, Pan X, Wei L, Min Y, Yu K, et al. Comparative transcriptome analyses revealed conserved and novel responses to cold and freezing stress in *Brassica napus* L. G3. 2019;**9**(8):2723-2737
- [12] Guy C. Molecular responses of plants to cold shock and cold acclimation. *Journal of Molecular Microbiology and Biotechnology*. 1999;**1**(2):231-242
- [13] Shi Y, Ding Y, Yang S. Molecular regulation of CBF signaling in cold acclimation. *Trends in Plant Science*. 2018;**23**(7):623-637
- [14] Kocsy G, Galiba G, Brunold C. Role of glutathione in adaptation and signalling during chilling and cold acclimation in plants. *Physiologia Plantarum*. 2001;**113**(2):158-164
- [15] Sanghera GS, Wani SH, Hussain W, Singh NB. Engineering cold stress tolerance in crop plants. *Current Genomics*. 2011;**12**(1):30-43
- [16] Luo T, Xian M, Zhang C, Zhang C, Hu L, Xu Z. Associating transcriptional regulation for rapid germination of

- rapeseed (*Brassica napus* L.) under low temperature stress through weighted gene co-expression network analysis. *Scientific Reports*. 2019;**9**(1):55
- [17] Zhang Y, Zhang D, Yu H, Lin B, Fu Y, Hua S. Floral initiation in response to planting date reveals the key role of floral meristem differentiation prior to budding in canola (*Brassica napus* L.). *Frontiers in Plant Science*. 2016;**7**:1369
- [18] Zhang SJ, Li L, Zhang CL. Effects of sowing date and planting density on the seed yield and oil content of winter oilseed rape. *The Journal of Applied Ecology*. 2012;**23**(5):1326-1332
- [19] Luo T, Zhang J, Khan MN, Liu JH, Xu ZH, Hu LY. Temperature variation caused by sowing dates significantly affects floral initiation and floral bud differentiation processes in rapeseed (*Brassica napus* L.). *Plant Science*. 2018;**271**:40-51
- [20] Hua SJ, Lin BG, Hussain N, Zhang YF, Yu HS, Ren Y, et al. Delayed planting affects seed yield, biomass production, and carbohydrate allocation in canola (*Brassica napus*). *International Journal of Agriculture and Biology*. 2014;**16**(4):671-680
- [21] Ozer H. Sowing date and nitrogen rate effects on growth, yield and yield components of two summer rapeseed cultivars. *European Journal of Agronomy*. 2003;**19**(3):453-463
- [22] Xue-Kun Z, Chun-Lei Z, Xing L, Han-Zhong W. Investigation on 2008 low temperature and freeze injure on winter rape along Yangtze River. *Chinese Journal of Oil Crop Sciences*. 2008;**30**:122-126
- [23] Tu Y-Q, X-L D. Effects of continuous low temperature overcast and rainy weather on yield and oil content of *Brassica napus* during flowering stage. *Chinese Journal of Oil Crop Sciences*. 2011;**33**:470-475
- [24] Do Ru A, Cak Rlar H. Is leaf age a predictor for cold tolerance in winter oilseed rape plants? *Functional Plant Biology: FPB*. 2020;**47**(3):250-262
- [25] Pu Y, Liu L, Wu J, Zhao Y, Bai J, Ma L, et al. Transcriptome profile analysis of winter rapeseed (*Brassica napus* L.) in response to freezing stress, reveal potentially connected events to freezing stress. *International Journal of Molecular Sciences*. 2019;**20**(11):2771
- [26] Yan L, Shah T, Cheng Y, Lu Y, Zhang XK, Zou XL. Physiological and molecular responses to cold stress in rapeseed (*Brassica napus* L.). *Journal of Integrative Agriculture*. 2019;**18**(12):2742-2752
- [27] Stefanowska M, Kuras M, Kacperska A. Low temperature-induced modifications in cell ultrastructure and localization of phenolics in winter oilseed rape (*Brassica napus* L. var. *oleifera* L.) leaves. *Annals of Botany*. 2002;**90**(5):637-645
- [28] Stefanowska M, Kuraś M, Kubacka-Zebalska M, Kacperska A. Low temperature affects pattern of leaf growth and structure of cell walls in winter oilseed rape (*Brassica napus* L., var. *oleifera* L.). *Annals of Botany*. 1999;**84**(3):313-319
- [29] Savitch LV, Allard G, Seki M, Robert LS, Tinker NA, Huner NP, et al. The effect of overexpression of two *Brassica* CBF/DREB1-like transcription factors on photosynthetic capacity and freezing tolerance in *Brassica napus*. *Plant & Cell Physiology*. 2005;**46**(9):1525-1539
- [30] Xiao-Hong Z, Liang-Jie F, Te-Wu Y, Zheng-Hua X, Li-Yong H. Effects of chilling stress on physiological characteristics of rapeseed seedlings in winter. *Plant Physiology*. 2015;**51**:737-746
- [31] Thomashow MF. PLANT COLD ACCLIMATION: Freezing tolerance

genes and regulatory mechanisms. Annual Review of Plant Physiology and Plant Molecular Biology. 1999;**50**:571-599

[32] Miura K, Furumoto T. Cold signaling and cold response in plants. International Journal of Molecular Sciences. 2013;**14**(3):5312-5337

[33] Ritonga FN, Chen S. Physiological and molecular mechanism involved in cold stress tolerance in plants. Plants. 2020;**9**(5):560

[34] Jaglo-Ottosen KR, Gilmour SJ, Zarka DG, Schabenberger O, Thomashow MF. *Arabidopsis* CBF1 overexpression induces COR genes and enhances freezing tolerance. Science. 1998;**280**(5360):104-106

[35] Liu Q, Kasuga M, Sakuma Y, Abe H, Miura S, Yamaguchi-Shinozaki K, et al. Two transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought- and low-temperature-responsive gene expression, respectively, in *Arabidopsis*. The Plant Cell. 1998;**10**(8):1391-1406

[36] Jaglo KR, Kleff S, Amundsen KL, Zhang X, Haake V, Zhang JZ, et al. Components of the *Arabidopsis* C-repeat/dehydration-responsive element binding factor cold-response pathway are conserved in *Brassica napus* and other plant species. Plant Physiology. 2001;**127**(3):910-917

[37] Chinnusamy V, Zhu J, Zhu JK. Cold stress regulation of gene expression in plants. Trends in Plant Science. 2007;**12**(10):444-451

[38] Chinnusamy V, Ohta M, Kanrar S, Lee BH, Hong X, Agarwal M, et al. ICE1: A regulator of cold-induced transcriptome and freezing tolerance in *Arabidopsis*. Genes & Development. 2003;**17**(8):1043-1054

[39] Kurbidaeva A, Ezhova T, Novokreshchenova M. *Arabidopsis thaliana* ICE2 gene: Phylogeny, structural evolution and functional diversification from ICE1. Plant Science: An International Journal of Experimental Plant Biology. 2014;**229**:10-22

[40] Kim YS, Lee M, Lee JH, Lee HJ, Park CM. The unified ICE-CBF pathway provides a transcriptional feedback control of freezing tolerance during cold acclimation in *Arabidopsis*. Plant Molecular Biology. 2015;**89**(1-2):187-201

[41] Ke L, Lei W, Yang W, Wang J, Gao J, Cheng J, et al. Genome-wide identification of cold responsive transcription factors in *Brassica napus* L. BMC Plant Biology. 2020;**20**(1):62

[42] Jian H, Xie L, Wang Y, Cao Y, Wan M, Lv D, et al. Characterization of cold stress responses in different rapeseed ecotypes based on metabolomics and transcriptomics analyses. PeerJ. 2020;**8**:e8704

[43] Gao MJ, Allard G, Byass L, Flanagan AM, Singh J. Regulation and characterization of four CBF transcription factors from *Brassica napus*. Plant Molecular Biology. 2002;**49**(5):459-471

[44] Weretilnyk E, Orr W, White TC, Iu B, Singh J. Characterization of three related low-temperature-regulated cDNAs from winter *Brassica napus*. Plant Physiology. 1993;**101**(1):171-177

[45] Du C, Hu K, Xian S, Liu C, Fan J, Tu J, et al. Dynamic transcriptome analysis reveals AP2/ERF transcription factors responsible for cold stress in rapeseed (*Brassica napus* L.). Molecular Genetics and Genomics: MGG. 2016;**291**(3):1053-1067

[46] Chen L, Zhong H, Ren F, Guo QQ, Hu XP, Li XB. A novel cold-regulated gene, COR25, of *Brassica napus* is

involved in plant response and tolerance to cold stress. *Plant Cell Reports*. 2011;**30**(4):463-471

[47] Wang Z, Yang C, Chen H, Wang P, Wang P, Song C, et al. Multi-gene co-expression can improve comprehensive resistance to multiple abiotic stresses in *Brassica napus* L. *Plant Science*. 2018;**274**:410-419

[48] Vishwakarma K, Upadhyay N, Kumar N, Yadav G, Singh J, Mishra RK, et al. Absciscic acid signaling and abiotic stress tolerance in plants: A review on current knowledge and future prospects. *Frontiers in Plant Science*. 2017;**8**:161

[49] Fahad S, Nie L, Chen Y, Wu C, Xiong D, Saud S, et al. Crop plant hormones and environmental stress. In: Lichtfouse E, editor. *Sustainable Agriculture Reviews*. Vol. 15. Cham: Springer International Publishing; 2015. pp. 371-400

[50] Fahad S, Hussain S, Matloob A, Khan FA, Khaliq A, Saud S, et al. Phytohormones and plant responses to salinity stress: A review. *Plant Growth Regulation*. 2015;**75**(2):391-404

[51] Shinkawa R, Morishita A, Amikura K, Machida R, Murakawa H, Kuchitsu K, et al. Absciscic acid induced freezing tolerance in chilling-sensitive suspension cultures and seedlings of rice. *BMC Research Notes*. 2013;**6**(1):351

[52] Huang X, Shi H, Hu Z, Liu A, Amombo E, Chen L, et al. ABA is involved in regulation of cold stress response in Bermudagrass. *Frontiers in Plant Science*. 2017;**8**:1613

[53] Ding Y, Li H, Zhang X, Xie Q, Gong Z, Yang S. OST1 kinase modulates freezing tolerance by enhancing ICE1 stability in *Arabidopsis*. *Developmental Cell*. 2015;**32**(3):278-289

[54] Ding Y, Jia Y, Shi Y, Zhang X, Song C, Gong Z, et al. OST1-mediated

BTF3L phosphorylation positively regulates CBFs during plant cold responses. *The EMBO Journal*. 2018;**37**(8):e98228

[55] Perkins M, Skori L, Hickerson NMN, Jamshed M, Samuel MA. Genetic manipulation of ABI3 confers frost-tolerant seed degreening in canola. *Plant Biotechnology Journal*. 2020;**18**(3):602-604

[56] McAinsh MR, Pittman JK. Shaping the calcium signature. *The New Phytologist*. 2009;**181**(2):275-294

[57] Dodd AN, Kudla J, Sanders D. The language of calcium signaling. *Annual Review of Plant Biology*. 2010;**61**:593-620

[58] Knight H, Trewavas AJ, Knight MR. Cold calcium signaling in *Arabidopsis* involves two cellular pools and a change in calcium signature after acclimation. *Plant Cell*. 1996;**8**(3):489-503

[59] Knight MR, Campbell AK, Smith SM, Trewavas AJ. Transgenic plant aequorin reports the effects of touch and cold-shock and elicitors on cytoplasmic calcium. *Nature*. 1991;**352**(6335):524-526

[60] Ma Y, Dai X, Xu Y, Luo W, Zheng X, Zeng D, et al. COLD1 confers chilling tolerance in rice. *Cell*. 2015;**160**(6):1209-1221

[61] Townley HE, Knight MR. Calmodulin as a potential negative regulator of *Arabidopsis* COR gene expression. *Plant Physiology*. 2002;**128**(4):1169

[62] Doherty CJ, Van Buskirk HA, Myers SJ, Thomashow MF. Roles for *Arabidopsis* CAMTA transcription factors in cold-regulated gene expression and freezing tolerance. *Plant Cell*. 2009;**21**(3):972-984

- [63] Kidokoro S, Yoneda K, Takasaki H, Takahashi F, Shinozaki K, Yamaguchi-Shinozaki K. Different cold-signaling pathways function in the responses to rapid and gradual decreases in temperature. *The Plant Cell*. 2017;**29**(4):760
- [64] He X, Liu W, Li W, Liu Y, Wang W, Pan Xie YK, et al. Genome-wide identification and expression analysis of CaM/CML genes in *Brassica napus* under abiotic stress. *Journal of Plant Physiology*. 2020:153251. DOI: 10.1016/j.jplph.2020.153251
- [65] Zhang H, Yang B, Liu W-Z, Li H, Wang L, Wang B, et al. Identification and characterization of CBL and CIPK gene families in canola (*Brassica napus* L.). *BMC Plant Biology*. 2014;**14**:8-8
- [66] Zhang H, Liu W-Z, Zhang Y, Deng M, Niu F, Yang B, et al. Identification, expression and interaction analyses of calcium-dependent protein kinase (CPK) genes in canola (*Brassica napus* L.). *BMC Genomics*. 2014;**15**:211-211
- [67] Edrisi Maryan K, Samizadeh Lahiji H, Farrokhi N, Hasani Komeleh H. Analysis of *Brassica napus* dehydrins and their co-expression regulatory networks in relation to cold stress. *Gene Expression Patterns: GEP*. 2019;**31**:7-17
- [68] Tasseva G, de Virville JD, Cantrel C, Moreau F, Zachowski A. Changes in the endoplasmic reticulum lipid properties in response to low temperature in *Brassica napus*. *Plant Physiology and Biochemistry: PPB*. 2004;**42**(10):811-822
- [69] Huang Z, Zhao N, Qin M, Xu A. Mapping of quantitative trait loci related to cold resistance in *Brassica napus* L. *Journal of Plant Physiology*. 2018;**231**:147-154
- [70] Nokhrina K, Ray H, Bock C, Georges F. Metabolomic shifts in *Brassica napus* lines with enhanced BnPLC2 expression impact their response to low temperature stress and plant pathogens. *GM Crops & Food*. 2014;**5**(2):120-131
- [71] Dahal K, Gadapati W, Savitch LV, Singh J, Huner NP. Cold acclimation and BnCBF17-over-expression enhance photosynthetic performance and energy conversion efficiency during long-term growth of *Brassica napus* under elevated CO₂ conditions. *Planta*. 2012;**236**(5):1639-1652
- [72] Lohani N, Jain D, Singh MB, Bhalla PL. Engineering multiple abiotic stress tolerance in canola, *Brassica napus*. *Frontiers in Plant Science*. 2020;**11**:3