We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Terpenes in Essential Oils: Bioactivity and Applications

Paco Noriega

Abstract

Secondary metabolites from plant organisms have always been excellent options for the pharmaceutical, cosmetic, and food industries. Essential oils are a type of metabolites found in vegetables, and their chemical composition is diverse; however, monoterpenes and sesquiterpenes are inside the most abundant molecules. These terpenes have a diverse chemical composition that range from a simple molecule with carbon and hydrogen to more complex molecules with oxygenated organic groups, such as alcohols, aldehydes, ketones, and ethers. Many of these molecules with 10 and 15 carbon atoms have an especially important biological activity, being important the antimicrobial, antifungal, antioxidant, anti-inflammatory, insecticide, analgesic, anticancer, cytotoxic, among others. Some of these substances are potentially toxic, and hence, they should be handled with caution, especially when they are pure. They are easily obtained by different methods, and their industrial value grows every year, with a market of several million dollars. This chapter seeks to provide a better understanding of this type of bioactive molecules, with an emphasis in those whose information is remarkable in the scientific literature and whose value for health and human well-being makes them extremely important.

Keywords: terpenes, essential oils, bioactivity, chemical analysis

1. Introduction

Terpenes are chemical molecules synthesized from isoprene, 2-methyl-1,3 butadiene which are polymerized, thus obtaining one of nature's most diversified families of secondary metabolites.

The chemical diversity of terpenes is determined by the polymerization capacity of isoprene; because of this their classification is linked to the addition of five carbons to the basic molecular unit. The biosynthesis of the chemical precursors of isoprene, dimethylallyl pyrophosphate (DMAPP) and isopentenyl pyrophosphate (IPP) is produced by two diversified metabolic routes, the mevalonate route (MEV) and the 2C-Methyl-D-erythriol-4-phosphate (MEP) route [1].

DMAPP and IPP are hemiterpenes and are responsible of forming the various subclasses of compounds that make up the terpenes. Additionally, these isoprene polymers can be linear or can form rings and adhere to their structure oxygen and nitrogen atoms. The approximate number of known terpenes is close to 55,000 compounds [2].

Traditionally they are classified as [3]:

Hemiterpenes. These are constituted by five carbon atoms and are the basic units of the terpenes, the best-known example is 2-methyl-1,3 butadiene or isoprene.

Monoterpenes. These are constituted by 10 carbon atoms, resulting from the union of two units of isoprene, which are abundant in essential oils. Some important substances are: pinene, myrcene, limonene, thujene, etc.

Sesquiterpenes. These are formed by 15 carbon atoms, which are the result of the junction of three units of isoprene, some examples are: bisabolene, zingiberene, germacrene, caryophyllene, etc.

Diterpenes. These are formed by 20 carbon atoms or four units of isoprene; some important compounds are retinol, taxol and phytol.

Triterpenes. Squalene and several phytosterols such as sitosterol stand out among the terpenes containing 30 carbon atoms or six units of isoprene.

Tetraterpenes. These are constituted by 40 carbon atoms and eight units of isoprene, many of them are dyes like carotenes, among these the most important are carotene, lycopene and bixin.

Polyterpenes. These are composed of more than 40 carbon atoms; they are often found in gums and latex of various plant species.

2. Essential oils

Essential oils are common secondary metabolites in vegetables. From 10 to 200 compounds can be found in an essential oil, and their main characteristic is their ability to evaporate at room temperature. The chemical variability in an oil is significant; however, its components can be classified into three large groups (**Figure 1**).

Terpenes are the majority group, being monoterpenes and sesquiterpenes the most abundant. These can be present as hydrocarbons, consisting of carbon and

Figure 1.
Main molecules of essential oils.

hydrogen, or can have various functional groups such as alcohols, thiols, aldehydes, ketones, and ethers.

The second group of importance is aromatic compounds, many of them with an important biological activity such as derivatives of cinnamaldehyde, thymol, anethole or carvacrol.

There is a third miscellaneous group in a lower proportion that groups various molecules such as hydrocarbons, aldehydes, ketones, esters, etc. Examples of these substances are isovaleraldehyde or dodecanal.

Essential oils are usually found in low concentrations in plant organisms, ranging from 0.1 to 1%. They can exceed this value as is the case of clove oil with up to 10%, and are present in all plant organs and leaves: *Mentha piperita*, *Origanum majorana*, *Thymus vulgaris*; flowers: *Rosa damascena*, *Matricaria chamomilla*, *Lavandula officinale*; stems: *Cinnamomum verum*, *Ocotea quixos*, *Santalum album*; roots: *Valeriana officinale*; fruits: *Citrus bergamia*; rhizomes: *Zingiber officinale*, *Curcuma longa*; and seeds: *Pimpinella anisum*, *Syzygium aromaticum* and *Cuminum cyminum*.

The extraction processes are diverse, depending on the part of the plant used; the simplest and most widespread is the extraction by distillation with steam current, which does not require expensive equipment. Other methods are mechanical extraction used mainly to obtain oil from citrus pericarps, extraction using solvents which is useful when components can be affected by high temperatures and extraction using a supercritical CO₂ current, which does not need high temperatures while maintaining the chemistry of molecules, but it is very expensive to implement.

About 4000 species have been investigated by their ability to produce essential oils, but only about 30 are marketed massively globally; their main use is intended for the cosmetic industry and aromatherapy, although several of the compounds from essences could be valuable to the pharmaceutical industry. There are certainly still species whose essential oils have not been analyzed in their chemical composition or in their bioactivity, which could be interesting as a source of new secondary metabolites.

3. Chemical analysis

Since they are volatile metabolites, their low boiling points make it possible to have them as steam in a remarkably simple way; for this reason the ideal analysis is gas chromatography with GC/MS mass spectrometry.

The use of capillary columns has made it possible to have defined separations in essential oils that exceed 100 compounds, usually chromatographic separation is made in nonpolar columns with 95% dimethylpolysiloxane, due to the fact that several components of an essential oil contain polar groups such as hydroxyl (OH); the realization of these components using columns of intermediate polarity has been made. Both assays result in a complete chemical inquiry of molecules and are complementary. The correct structural elucidation is performed by combining several analyses such as comparison with spectrum databases and the theoretical and experimental determination of the retention rates of the compounds. For this purpose, there are databases, being the most used the "Identification of essential oil components by gas chromatography/mass spectrometry," with approximately 4000 compounds from essential oils [4].

The GC/MS technique is limited in the fact that it is ineffective in evaluating stereoisomers, in such cases it is necessary to use chiral columns or techniques such as nuclear magnetic resonance imaging.

A more thorough investigation of the chemical identity of the molecules of an essential oil can be done with an equipment that couples gas chromatography with spectrophotometric techniques, such as nuclear magnetic resonance imaging and infrared spectroscopy. It is also possible to analyze NMR or IR spectra in previously isolated molecules by column or thin layer chromatography [5].

4. Monoterpenes with therapeutic importance

Several monoterpenes have a diverse and useful biological activity for treating diseases and ailments; some have valuable aromatic characteristics in cosmetics and perfumery. Those molecules that have relevant information and studies are analyzed to verify their use as phytotherapeutic elements (**Figure 2**).

Pinenes. These have alpha and beta isomers; their formula is $C_{10}H_{16}$ and they are common in essential oils from conifers, although they can be found in many other species such as rosemary and lavender [6–8]; oils with high concentrations of pinenes generally have antimicrobial activity [8, 9]. Traditionally many plants containing pinene-rich essential oils are used in respiratory system disease [9].

1-8 cineol (Eucalyptol). Oxygenated monoterpene has a $C_{10}H_{18}O$ formula whose functional group is an ether that is present in many varieties of eucalyptus. Among its most noteworthy properties are analgesic, anti-inflammatory and antimicrobial [10]. Plants with eucalyptol-rich essential oils are used for expectorant and decongestant properties of the respiratory system [11].

Limonene. It is a monoterpene whose formula is $C_{10}H_{16}$; it has two optical isomers R-limonene or D-limonene and S-limonene or L-limonene, which stand out by the insecticide [12, 13] and antimicrobial properties [14].

Myrcene. It is a monoterpene whose formula is $C_{10}H_{16}$; it is the main component of *Cannabis sativa* essential oil [15]. Several studies highlight its analgesic-sedative [16, 17] and anti-inflammatory activity [18].

Linalool. It is a hydroxylated monoterpene with $C_{10}H_{18}O$ formula, its pleasant aroma makes it widely used in perfumery. Its action on the central nervous system is evidenced by its sedative, anxiolytic, analgesic and anti-inflammatory

Figure 2.
Monoterpene molecules with therapeutic importance.

properties [19, 20]. Its antimicrobial and antioxidant properties have been evaluated with good results [21, 22].

Citral. It is an oxygenated monoterpene containing a group of aldehyde; its formula is $C_{10}H_{16}O$. There are two isomers known as neral (cis isomer) and geranial (trans isomer) [23], which are abundant in species such as *Backhousia citriodora* [24] and *Cymbopogon citratus* [25]. Its antimicrobial [25] and insect repellent action is noteworthy [26].

Camphor. It is an oxygenated monoterpene whose functional group is a ketone; its formula is $C_{10}H_{16}O$ and it is present in two optical isomers R and S, which are abundant in the species *Cinnamomum camphora* [27]. Traditionally camphor has been used in traditional Asian medicine, and it is known to have digestive effects, but its most important use is related to its analgesic and antiseptic effect, being very popular its inclusion in topical formulations such as liniments and creams [28, 29].

Menthol. It is a hydroxylated monoterpene, with a $C_{10}H_{20}O$ formula, which has seven isomers that are very common in mint varieties such as Peppermint. It is one of the most used compounds in the food, cosmetic, pharmaceutical industries, and pesticides, among others. Its aromatic properties are very well known [30]; however, its most noticeable and known effect is that of analgesia at the topical level [31, 32].

Terpineol. It is a hydroxylated monoterpene with a $C_{10}H_{18}O$ formula. It is known by having five isomers (α , β , γ , δ and 4-terpineol) [33], which are abundant in the essential oil of tea tree (*Malaleuca alternifolia*) [34]. The outstanding properties are antihypertensive, anticancer, antioxidant, antimicrobial, antifungal and sedative [33, 35].

Citronellol. It is a hydroxylated monoterpene with a $C_{10}H_{20}O$ formula. There are two enantiomers (+)-citronellol and (–)-citronellol [36]. The first is quite common in citronella oil, and the second is abundant in rose oil [37], which is used in perfumery. Its properties are insecticide [38], analgesic and anti-inflammatory [39], and antioxidant [40].

5. Sesquiterpenes with therapeutic importance

Several molecules with interesting properties can be found in C_{15} sesquiterpene (**Figure 3**). From a therapeutic view, there is evidence that validates its biological activity, highlighting anti-inflammatory, analgesic and anticancer trials.

Bisabolol. These are isomers, out of which stand (–)- α -Bisabolol, (–)-epi- α -Bisabolol, (+)- α -Bisabolol and (+)-epi- α -Bisabolol which are abundant in the species *Matricaria camomilla* [41], and in other species such as *Salvia runcinata* [42]. The most well-known effects in the molecule are analgesic and anti-inflammatory [43], antimicrobial and antioxidant properties; for this reason, the molecule is widely used in the cosmetic industry [44].

β -Caryophyllene. It has a $C_{15}H_{24}$ formula. It is one of the most abundant sesquiterpenes in essential oils. Various bioactivity studies have been carried out in this molecule with good results, such as analgesic [45, 46], anti-inflammatory [47] and anticancer [48].

Chamazulene. With a $C_{14}H_{16}$ formula, it is a molecule derived from the sesquiterpene matricina, which is one of the few aromatic molecules that have a blue coloration. It is found in *Matricaria camomilla*, being its most known property the anti-inflammatory [49, 50]. Several studies highlight its antioxidant effects [51, 52].

Caryophyllene oxide. It is an oxygenated sesquiterpene with a $C_{15}H_{24}O$ formula; it has properties similar to those of caryophyllene, such as analgesic and anti-inflammatory [53].

Figure 3.
Sesquiterpene molecules with therapeutic purposes.

Germacrene. It belongs to the sesquiterpenes family, and it has three double links in its structure. There are five types of germacrenes: A, B, C, D, E. Recent studies mention its antioxidant potential [5, 54].

Artemisinic acid. It has a $C_{15}H_{22}O_2$ formula, and it is one of the most interesting sesquiterpenes for health due to its antimalarial properties [55]. It is abundant in the species *Artemisia annua*, and it is generally found as a sesquiterpenic lactone [56].

Patchoulene. It is a sesquiterpene with a $C_{15}H_{24}$ formula. It is common to find its isomers α , β , α , and δ in essential oils. It is attributed to various types of bioactivity, the most relevant being those found in β -patchoulene as anti-inflammatory [57], antigastritis [58, 59], and cosmetic [60].

Humulene. Also known as α -caryophyllene, its formula is $C_{15}H_{24}$. It is named after the essential oil of the species *Humulus lupulus* [61]. It has anti-inflammatory [62, 63] and anticancer properties [64].

Bergamotene. It is a sesquiterpene with a $C_{15}H_{24}$ formula. It has four isomers α -cis, β -cis, α -trans and β -trans. It is found in several citric species such as *Citrus bergamia* [65]. One of the properties of this molecule is to act as a pheromone [66, 67].

Farnesene. It has a $C_{15}H_{24}$ formula. It is a molecule found in several essential oils, and it is a precursor to many other sesquiterpenes since its open-chain structure and its 4-double bonds contribute to this action, as well as in the possibility of having a wide variety of isomers between geometrics and stereoisomers. Its cytotoxic and genotoxic [68], insecticide [69] and neuroprotective effects [70, 71] have been evaluated.

Eudesmol. Hydroxylated sesquiterpene with a $C_{15}H_{26}O$ formula is a very interesting molecule by the multiple positive bioactivity assays, highlighting antimicrobial and antifungal [72], anticancer [73, 74] and antiangiogenic [75].

6. Toxicity

Most of the terpenes present in essential oils have some degree of toxicity, which is not detected when consuming aromatic species directly because in most cases

the oil yield is low. Many commonly used essential oil components are potentially dermal irritating with restrictions on application concentrations [76, 77]. There are also some terpenes whose toxicity is much more dangerous, such as pulegone which causes liver damage and seizures [78], and thujone that can cause dementia by being neurotoxic [79].

7. Conclusion

This brief review has shown the chemical and biological importance of low molecular weight and volatile terpenes. For this reason, components of secondary metabolites are known as essential oils. The abundance of these molecules is much higher than the one presented in this chapter, since the information presented covers those whose scientific evidence and industrial importance are references in this family of metabolites. There is still much research to be carried out on the hundreds of molecules from which there is still little or no information. There are still aromatic species whose essential oils have not yet been described and that could be a source of new monoterpenes and sesquiterpenes that are beneficial to humans.

Author details

Paco Noriega
Salesian Polytechnic University, Quito, Ecuador

*Address all correspondence to: pnoriega@ups.edu.ec

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Natanya C. Natural Products in Chemical Biology. Hoboken, New Jersey: John Wiley & Sons; 2012. pp. 127-129
- [2] Guimarães AG, Serafini MR, Quintans-Júnior LJ. Terpenes and derivatives as a new perspective for pain treatment: A patent review. *Expert Opinion on Therapeutic Patents*. 2014;**24**(3):243-265
- [3] Perveen S, Al-Taweel A. Introductory chapter: Terpenes and terpenoids. In: *Terpenes and Terpenoids*. London: IntechOpen; 2018. pp. 1-7
- [4] Adams RP. Identification of Essential Oils by Ion Trap Mass Spectroscopy. San Diego, California: Academic Press; 2012
- [5] Noriega P, Guerrini A, Sacchetti G, Grandini A, Ankuash E, Manfredini S. Chemical composition and biological activity of five essential oils from the Ecuadorian Amazon rain forest. *Molecules*. 2019;**24**(8):1637
- [6] Jiang Y, Wu N, Fu YJ, Wang W, Luo M, Zhao CJ, et al. Chemical composition and antimicrobial activity of the essential oil of rosemary. *Environmental Toxicology and Pharmacology*. 2011;**32**(1):63-68
- [7] Zheljazkov VD, Astatkie T, Hristov AN. Lavender and hyssop productivity, oil content, and bioactivity as a function of harvest time and drying. *Industrial Crops and Products*. 2012;**36**(1):222-228
- [8] Silva ACRD, Lopes PM, Azevedo MMBD, Costa DCM, Alviano CS, Alviano DS. Biological activities of α -pinene and β -pinene enantiomers. *Molecules*. 2012;**17**(6):6305-6316
- [9] Rivera PFN, Paredes EA, Gómez ED, Lueckhoff A, Almeida GA, Suarez SE. Composición química y actividad antimicrobiana del aceite esencial de los rizomas de *Renealmia thyrsoides* (Ruiz & Pav) Poepp. & Eddl (shiwanku muyu). *Revista Cubana de Plantas Medicinales*. 2017;**22**(2)
- [10] Bhowal M, Gopal M. Eucalyptol: Safety and pharmacological profile. *Journal of Pharmaceutical Sciences*. 2015;**5**:125-131
- [11] Adnan M. Bioactive potential of essential oil extracted from the leaves of *Eucalyptus globulus* (Myrtaceae). *Journal of Pharmacognosy and Phytochemistry*. 2019;**8**(1):213-216
- [12] Karr LL, Costas JR. Insecticidal properties of d-limonene. *Journal of Pesticide Science*. 1988;**13**(2):287-290
- [13] Malacrinò A, Campolo O, Laudani F, Palmeri V. Fumigant and repellent activity of limonene enantiomers against *Tribolium confusum* du Val. *Neotropical Entomology*. 2016;**45**(5):597-603
- [14] Espina L, Gelaw TK, de Lamo-Castellví S, Pagán R, García-Gonzalo D. Mechanism of bacterial inactivation by (+)-limonene and its potential use in food preservation combined processes. *PLoS One*. 2013;**8**(2):e56769
- [15] Gulluni N, Re T, Loiacono I, Lanzo G, Gori L, Macchi C, et al. Cannabis essential oil: A preliminary study for the evaluation of the brain effects. *Evidence-Based Complementary and Alternative Medicine*. 2018;**(1)**:1-11
- [16] Mirghaed AT, Ghelichpour M, Hoseini SM. Myrcene and linalool as new anesthetic and sedative agents in common carp, *Cyprinus carpio* - Comparison with eugenol. *Aquaculture*. 2016;**464**:165-170

- [17] Rao VSN, Menezes AMS, Viana GSB. Effect of myrcene on nociception in mice. *The Journal of Pharmacy and Pharmacology*. 1990;**42**(12):877-878
- [18] Rufino AT, Ribeiro M, Sousa C, Judas F, Salgueiro L, Cavaleiro C, et al. Evaluation of the anti-inflammatory, anti-catabolic and pro-anabolic effects of E-caryophyllene, myrcene and limonene in a cell model of osteoarthritis. *European Journal of Pharmacology*. 2015;**750**:141-150
- [19] Aprotosoiaie AC, Hăncianu M, Costache II, Miron A. Linalool: A review on a key odorant molecule with valuable biological properties. *Flavour and Fragrance Journal*. 2014;**29**(4):193-219
- [20] Pereira I, Severino P, Santos AC, Silva AM, Souto EB. Linalool bioactive properties and potential applicability in drug delivery systems. *Colloids and Surfaces, B: Biointerfaces*. 2018;**171**:566-578
- [21] Herman A, Tambor K, Herman A. Linalool affects the antimicrobial efficacy of essential oils. *Current Microbiology*. 2016;**72**(2):165-172
- [22] Hu J, Liu S, Deng W. Dual responsive linalool capsules with high loading ratio for excellent antioxidant and antibacterial efficiency. *Colloids and Surfaces, B: Biointerfaces*. 2020;**190**:110978
- [23] Sacks J, Greenley E, Leo G, Willey P, Gallis D, Mangravite JA. Separation and analysis of citral isomers: An undergraduate organic laboratory experiment. *Journal of Chemical Education*. 1983;**60**(5):434
- [24] Southwell IA, Russell M, Smith RL, Archer DW. *Backhousia citriodora* F. Muell. (Myrtaceae), a superior source of citral. *Journal of Essential Oil Research*. 2000;**12**(6):735-741
- [25] Onawunmi GO, Yisak WA, Ogunlana EO. Antibacterial constituents in the essential oil of *Cymbopogon citratus* (DC.) Stapf. *Journal of Ethnopharmacology*. 1984;**12**(3):279-286
- [26] Hao H, Wei J, Dai J, Du J. Host-seeking and blood-feeding behavior of *Aedes albopictus* (Diptera: Culicidae) exposed to vapors of geraniol, citral, citronellal, eugenol, or anisaldehyde. *Journal of Medical Entomology*. 2014;**45**(3):533-539
- [27] Pragadheesh VS, Saroj A, Yadav A, Chanotiya CS, Alam M, Samad A. Chemical characterization and antifungal activity of *Cinnamomum camphora* essential oil. *Industrial Crops and Products*. 2013;**49**:628-633
- [28] Green BG. Sensory characteristics of camphor. *The Journal of Investigative Dermatology*. 1990;**94**(5):662-666
- [29] Craven R. The comfort of camphor. *Nature Reviews. Neuroscience*. 2005;**6**(11):826-826
- [30] Kamatou GP, Vermaak I, Viljoen AM, Lawrence BM. Menthol: A simple monoterpene with remarkable biological properties. *Phytochemistry*. 2013;**96**:15-25
- [31] Galeotti N, Mannelli LDC, Mazzanti G, Bartolini A, Ghelardini C. Menthol: A natural analgesic compound. *Neuroscience Letters*. 2002;**322**(3):145-148
- [32] Pan R, Tian Y, Gao R, Li H, Zhao X, Barrett JE, et al. Central mechanisms of menthol-induced analgesia. *The Journal of Pharmacology and Experimental Therapeutics*. 2012;**343**(3):661-672
- [33] Khaleel C, Tabanca N, Buchbauer G. α -Terpineol, a natural monoterpene: A review of its biological properties. *Open Chemistry*. 2018;**16**(1):349-361
- [34] Papadopoulos CJ, Carson CF, Chang BJ, Riley TV. Role of the

MexAB-OprM efflux pump of *Pseudomonas aeruginosa* in tolerance to tea tree (*Melaleuca alternifolia*) oil and its monoterpene components terpinen-4-ol, 1, 8-cineole, and α -terpineol. *Applied and Environmental Microbiology*. 2008;**74**(6):1932-1935

[35] Kong Q, Zhang L, An P, Qi J, Yu X, Lu J, et al. Antifungal mechanisms of α -terpineol and terpene-4-alcohol as the critical components of *Melaleuca alternifolia* oil in the inhibition of rot disease caused by *Aspergillus ochraceus* in postharvest grapes. *Journal of Applied Microbiology*. 2019;**126**(4):1161-1174

[36] Ravid U, Putievsky E, Katzir I, Ikan R, Weinstein V. Determination of the enantiomeric composition of citronellol in essential oils by chiral GC analysis on a modified γ -cyclodextrin phase. *Flavour and Fragrance Journal*. 1992;**7**(4):235-238

[37] Katsukawa M, Nakata R, Koeji S, Hori K, Takahashi S, Inoue H. Citronellol and geraniol, components of rose oil, activate peroxisome proliferator-activated receptor α and γ and suppress cyclooxygenase-2 expression. *Bioscience, Biotechnology, and Biochemistry*. 2011;**75**(5):1010-1012

[38] Brari J, Thakur DR. Insecticidal potential properties of citronellol derived ionic liquid against two major stored grain insect pests. *Journal of Entomology and Zoology Studies*. 2016;**4**(3):365-370

[39] Brito RG, Guimarães AG, Quintans JS, Santos MR, De Sousa DP, Badaue-Passos D, et al. Citronellol, a monoterpene alcohol, reduces nociceptive and inflammatory activities in rodents. *Journal of Natural Medicines*. 2012;**66**(4):637-644

[40] Jagdale AD, Kamble SP, Nalawade ML, Arvindekar AU. Citronellol: A potential antioxidant and aldose reductase inhibitor from *Cymbopogon citratus*.

International Journal of Pharmacy and Pharmaceutical Sciences. 2015;**7**(3):203-209

[41] Isaac O, Thiemer K. Biochemical studies on camomile components/III. In vitro studies about the antipeptic activity of (–)- α -bisabolol (author's transl). *Arzneimittel-Forschung*. 1975;**25**:1352-1354

[42] Viljoen AM, Gono-Bwalya AB, Kamatou GPP, Baser KHC, Demirci B. The essential oil composition and chemotaxonomy of *Salvia stenophylla* and its allies *S. repens* and *S. runcinata*. *Journal of Essential Oil Research*. 2006;**18**:37-45

[43] Rocha NFM, Rios ERV, Carvalho AMR, Cerqueira GS, de Araújo LA, Leal LKAM, et al. Antinociceptive and anti-inflammatory activities of (–)- α -bisabolol in rodents. *Naunyn-Schmiedeberg's Archives of Pharmacology*. 2011;**384**(6):525-533

[44] Kamatou GP, Viljoen AM. A review of the application and pharmacological properties of α -bisabolol and α -bisabolol-rich oils. *Journal of the American Oil Chemists' Society*. 2010;**87**(1):1-7

[45] Ghelardini C, Galeotti N, Mannelli LDC, Mazzanti G, Bartolini A. Local anaesthetic activity of β -caryophyllene. *Il Farmaco*. 2001;**56**(5-7):387-389

[46] Fidyk K, Fiedorowicz A, Strzdała L, Szumny A. β -Caryophyllene and β -caryophyllene oxide—Natural compounds of anticancer and analgesic properties. *Cancer Medicine*. 2016;**5**(10):3007-3017

[47] Tambe Y, Tsujiuchi H, Honda G, Ikeshiro Y, Tanaka S. Gastric cytoprotection of the non-steroidal anti-inflammatory sesquiterpene, β -caryophyllene. *Planta Medica*. 1996;**62**(05):469-470

- [48] Legault J, Pichette A. Potentiating effect of β -caryophyllene on anticancer activity of α -humulene, isocaryophyllene and paclitaxel. *The Journal of Pharmacy and Pharmacology*. 2007;**59**(12):1643-1647
- [49] Ma D, He J, He D. Chamazulene reverses osteoarthritic inflammation through regulation of matrix metalloproteinases (MMPs) and NF- κ B pathway in in-vitro and in-vivo models. *Bioscience, Biotechnology, and Biochemistry*. 2020;**84**(2):402-410
- [50] Flemming M, Kraus B, Rasclé A, Jürgenliemk G, Fuchs S, Fürst R, et al. Revisited anti-inflammatory activity of matricine in vitro: Comparison with chamazulene. *Fitoterapia*. 2015;**106**:122-128
- [51] Rekka EA, Kourounakis AP, Kourounakis PN. Investigation of the effect of chamazulene on lipid peroxidation and free radical processes. *Research Communications in Molecular Pathology and Pharmacology*. 1996;**92**(3):361-364
- [52] Capuzzo A, Occhipinti A, Maffei ME. Antioxidant and radical scavenging activities of chamazulene. *Natural Product Research*. 2014;**28**(24):2321-2323
- [53] Chavan MJ, Wakte PS, Shinde DB. Analgesic and anti-inflammatory activity of caryophyllene oxide from *Annona squamosa* L. bark. *Phytomedicine*. 2010;**17**(2):149-151
- [54] Casiglia S, Bruno M, Bramucci M, Quassinti L, Lupidi G, Fiorini D, et al. *Kundmannia sicula* (L.) DC: A rich source of germacrene D. *Journal of Essential Oil Research*. 2017;**29**(6):437-442
- [55] Ro DK, Ouellet M, Paradise EM, Burd H, Eng D, Paddon CJ, et al. Induction of multiple pleiotropic drug resistance genes in yeast engineered to produce an increased level of anti-malarial drug precursor, artemisinic acid. *BMC Biotechnology*. 2008;**8**(1):83
- [56] Ram M, Gupta MM, Naqvi AA, Kumar S. Effect of planting time on the yield of essential oil and artemisinin in *Artemisia annua* under subtropical conditions. *Journal of Essential Oil Research*. 1997;**9**(2):193-197
- [57] Zhang Z, Chen X, Chen H, Wang L, Liang J, Luo D, et al. Anti-inflammatory activity of β -patchoulene isolated from patchouli oil in mice. *European Journal of Pharmacology*. 2016;**781**:229-238
- [58] Liu Y, Liang J, Wu J, Chen H, Zhang Z, Yang H, et al. Transformation of patchouli alcohol to β -patchoulene by gastric juice: β -patchoulene is more effective in preventing ethanol-induced gastric injury. *Scientific Reports*. 2017;**7**(1):1-13
- [59] Liang J, Dou Y, Wu X, Li H, Wu J, Huang Q, et al. Prophylactic efficacy of patchoulene epoxide against ethanol-induced gastric ulcer in rats: Influence on oxidative stress, inflammation and apoptosis. *Chemico-Biological Interactions*. 2018;**283**:30-37
- [60] Api AM, Belsito D, Botelho D, Browne D, Bruze M, Burton GA, et al. RIFM fragrance ingredient safety assessment β -Patchoulene, CAS Registry Number 514-51-2. *Food and Chemical Toxicology*. 2018;**115**:S256-S263
- [61] Nance MR, Setzer WN. Volatile components of aroma hops (*Humulus lupulus* L.) commonly used in beer brewing. *Journal of Brewing and Distilling*. 2011;**2**(2):16-22
- [62] Rogerio AP, Andrade EL, Leite DF, Figueiredo CP, Calixto JB. Preventive and therapeutic anti-inflammatory properties of the sesquiterpene α -humulene in experimental airways allergic inflammation. *British Journal of Pharmacology*. 2009;**158**(4):1074-1087

- [63] Fernandes ES, Passos GF, Medeiros R, da Cunha FM, Ferreira J, Campos MM, et al. Anti-inflammatory effects of compounds α -humulene and (–)-trans-caryophyllene isolated from the essential oil of *Cordia verbenacea*. *European Journal of Pharmacology*. 2007;**569**(3):228-236
- [64] El Hadri A, del Rio MG, Sanz J, Coloma AG, Idaomar M, Ozonas BR, et al. Cytotoxic activity of α -humulene and transcaryophyllene from *Salvia officinalis* in animal and human tumor cells. *Anales de la Real Academia Nacional de Farmacia*. 2010;**76**(3):343-356
- [65] Navarra M, Mannucci C, Delbò M, Calapai G. *Citrus bergamia* essential oil: From basic research to clinical application. *Frontiers in Pharmacology*. 2015;**6**:36
- [66] Cônsoli FL, Williams HJ, Vinson SB, Matthews RW, Cooperband MF. Trans-bergamotenes—Male pheromone of the ectoparasitoid *Melittobia digitata*. *Journal of Chemical Ecology*. 2002;**28**(8):1675-1689
- [67] Haber AI, Sims JW, Mescher MC, De Moraes CM, Carr DE. A key floral scent component (β -trans-bergamotene) drives pollinator preferences independently of pollen rewards in seep monkeyflower. *Functional Ecology*. 2019;**33**(2):218-228
- [68] Çelik K, Toğar B, Türkeş H, Taşpınar N. In vitro cytotoxic, genotoxic, and oxidative effects of acyclic sesquiterpene farnesene. *Turkish Journal of Biology*. 2014;**38**(2):253-259
- [69] Sun Y, Qiao H, Ling Y, Yang S, Rui C, Pelosi P, et al. New analogues of (E)- β -farnesene with insecticidal activity and binding affinity to aphid odorant-binding proteins. *Journal of Agricultural and Food Chemistry*. 2011;**59**(6):2456-2461
- [70] Arslan ME, Türkeş H, Mardinoğlu A. In vitro neuroprotective effects of farnesene sesquiterpene on alzheimer's disease model of differentiated neuroblastoma cell line. *The International Journal of Neuroscience*. 2020;**130**:1-10
- [71] Turkez H, Sozio P, Geyikoglu F, Tatar A, Hacimuftuoglu A, Di Stefano A. Neuroprotective effects of farnesene against hydrogen peroxide-induced neurotoxicity in vitro. *Cellular and Molecular Neurobiology*. 2014;**34**(1):101-111
- [72] Noriega P, Ballesteros J, De la Cruz A, Veloz T. Chemical composition and preliminary antimicrobial activity of the hydroxylated sesquiterpenes in the essential oil from *Piper barbatum* Kunth leaves. *Plants*. 2020;**9**(2):211
- [73] Plengsuriyakarn T, Karbwang J, Na-Bangchang K. Anticancer activity using positron emission tomography-computed tomography and pharmacokinetics of β -eudesmol in human cholangiocarcinoma xenografted nude mouse model. *Clinical and Experimental Pharmacology and Physiology*. 2015;**42**(3):293-304
- [74] Miyazawa M, Shimamura H, Nakamura SI, Kameoka H. Antimutagenic activity of (+)- β -eudesmol and paeonol from *Dioscorea japonica*. *Journal of Agricultural and Food Chemistry*. 1996;**44**(7):1647-1650
- [75] Tsuneki H, Ma EL, Kobayashi S, Sekizaki N, Maekawa K, Sasaoka T, et al. Antiangiogenic activity of β -eudesmol in vitro and in vivo. *European Journal of Pharmacology*. 2005;**512**(2-3):105-115
- [76] Mekonnen A, Tesfaye S, Christos SG, Dires K, Zenebe T, Zegeye N, et al. Evaluation of skin irritation and acute and subacute oral toxicity of *Lavandula angustifolia* essential oils in rabbit and mice. *Journal of Toxicology*. 2019;(1):1-8

[77] Lee CJ, Chen LW, Chen LG, Chang TL, Huang CW, Huang MC, et al. Correlations of the components of tea tree oil with its antibacterial effects and skin irritation. *Journal of Food and Drug Analysis*. 2013;**21**(2):169-176

[78] Gordon WP, Forte AJ, McMurtry RJ, Gal J, Nelson SD. Hepatotoxicity and pulmonary toxicity of pennyroyal oil and its constituent terpenes in the mouse. *Toxicology and Applied Pharmacology*. 1982;**65**(3):413-424

[79] Pelkonen O, Abass K, Wiesner J. Thujone and thujone-containing herbal medicinal and botanical products: Toxicological assessment. *Regulatory Toxicology and Pharmacology*. 2013;**65**(1):100-107