We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Elevated CO₂ Concentration Improves Heat-Tolerant Ability in Crops

Ayman EL Sabagh, Akbar Hossain, Mohammad Sohidel Islam, Muhammad Aamir Iqbal, Ali Raza, Çetin Karademir, Emine Karademir, Abdul Rehman, Md Atikur Rahman, Rajesh Kumar Singhal, Analía Llanes, Muhammad Ali Raza, Muhammad Mubeen, Wajid Nasim, Celaledin Barutçular, Ram Swaroop Meena and Hirofumi Saneoka

Abstract

The rising concentration of atmospheric carbon dioxide (aCO₂) and increasing temperature are the main reasons for climate change, which are significantly affecting crop production systems in this world. However, the elevated carbon dioxide (CO₂) concentration can improve the growth and development of crop plants by increasing photosynthetic rate (higher availability of photoassimilates). The combined effects of elevated CO₂ (eCO₂) and temperature on crop growth and carbon metabolism are not adequately recognized, while both eCO₂ and temperature triggered noteworthy changes in crop production. Therefore, to increase crop yields, it is important to identify the physiological mechanisms and genetic traits of crop plants which play a vital role in stress tolerance under the prevailing conditions. The eCO₂ and temperature stress effects on physiological aspects as well as biochemical profile to characterize genotypes that differ in their response to stress conditions. The aim of this review is directed the open-top cavities to regulate the properties like physiological, biochemical, and yield of crops under increasing aCO₂, and temperature. Overall, the extent of the effect of eCO₂ and temperature response to biochemical components and antioxidants remains unclear, and therefore further studies are required to promote an unperturbed production system.

Keywords: elevated CO₂, heat stress, physio-biochemical mechanisms

1. Introduction

Climate change in the form of increasing temperature and increasingly variable rainfall patterns threatens the production of the crop [1, 2]. Therefore, elevated CO₂ (eCO₂) absorption may indorse plant growth, whereas increased temperature

is repressive for C_3 plants. Both CO_2 and temperature caused significant changes in crop productivity. The collaborative properties of eCO_2 and increased temperatures on the crop growth and carbon metabolism are not well known.

The fluctuating climatic surroundings are predictable to upsurge the atmospheric CO_2 (aCO_2) meditation, temperatures and modify the rainfall outline. The aCO_2 meditation is prophesied to range 550 ppm by 2050, and possibly surpass 700 ppm by the end of the present century [3]. These fluctuations are expected to disturb the creation and output of cultivated crops, and stimulus the upcoming food safety. The influence analysis of climate alteration on worldwide food construction reveals a 0.5% failure by 2020 and 2.3% by 2050 [4]. The progress of climate arranged germplasm to counterbalance these wounded is of the highest reputation [5].

The eCO_2 is significant abiotic stress and has a noteworthy fertilization encouragement on crops. Widespread preceding educations have described that eCO_2 meaningfully enhanced the water use efficiency (WUE), reduced transpiration frequency, abridged maize growth rate, and augmented plant height, leaf number, leaf area, growth frequency, and overall yield [6]. Furthermore, the cumulative of aCO_2 disturbs precipitation equilibrium, which can alter the periodic precipitation circulation [7]. It has been predicted that this result would carry about a 10% upsurge or decline in water capitals in several areas [8]. The raised temperature, i.e., heat stress (HS) damage growth and physiological ailments, and consequentially reduce yield [9]. Increased temperature due to eCO_2 has a primary effect on the food grain invention reliant on the places. With the rising temperature by 1.0–2.0°C in tropical and subtropical states and the food grain manufacture in India is predictable to decline up to 30% [9].

The C_4 grass maize (*Zea mays* L.) is the third most vital food crop worldwide in the relation of the invention, and its claim is prophesied to rise by 45% from 1997 to 2020 [10]. Educations with maize retort to dual the ambient CO_2 presented variable possessions on growth fluctuating from no inspiration of yield [11] to 50% stimulation [12]. The growth and productivity of maize are expected to be pretentious by raised aCO_2 and temperature. Raised temperature severely disturbs the growth, and yield of maize plants [13]. There is unpredictable information on the properties of eCO_2 on the vintage of maize changing from slight positive consequence [14], no consequence [15] to rice harvest by 50% [16].

Record of the experimentations on the influences of eCO_2 and temperature on the crop yield, though, used measured atmosphere amenities like phytotron and plant growth cavities or crop growth reproduction models [13, 17]. Determination of the impact of raised CO_2 on the photosynthesis tolerance to severe HS is vital to expect the plant replies for universal warming since photosynthesis is sensitive to severe HS and aCO_2 upsurgences slightly [18, 19]. Further, flowering is a critical element for plant generative achievement and seed-set. The increase in temperature and eCO_2 is the main climate revolution issues that might influence plant suitability and associated flowering actions. Resolving the influence of these ecological issues on the flowering actions like time of days to anthesis and flowering (duration from germination till flowering) is serious to appreciate the acclimatization of crops in altering climate [20].

2. Interaction of eCO_2 with high-temperature stress and other factors to climate change

The impacts of eCO_2 and stress factors on crops have been made using controlled atmosphere amenities like plant growth cavities or crop growth reproduction

models in many studies [13]. The eCO₂ contributes to global warming, causing alterations in the precipitations, water scarcity and changes at temperatures in several regions affecting the growth and development of crop plants [3]. However, the interactive effects of eCO₂ and environmental stress conditions on the crop growth and carbon metabolism are not well predictable. The interactions between eCO₂ and stress factors are critical to photosynthesis performance. It has been reported how stomata react to eCO₂ levels, but the effects on photosynthesis performance of other environmental factors are poorly understood [21].

Numerous studies showed the combining impacts of eCO₂ and drought and revealed that the machines are different from singular eCO₂ and drought. The impact of combined eCO₂ and drought are varying with crop stage such as during vegetative stage, and it restricted the shoot development, decreased leaf area, diminished mobilization of nutrients due to weak root growth, reduced stomatal closure, transpiration, and relative water contents (RWC). However, it enhances the resource use efficiencies of the plant, including WUE, light use efficiency (LUE), and nutrient use efficiency (NUE) at a certain level [22, 23]. Similarly, eCO₂ and drought affect reproductive growth severely such as the impacts on the pollen abortion, pollination, flower formation, panicle length, panicle weight, seed formation, seed size and, yield potential of important agriculture crops [22].

Altering climate, counting eCO₂, increasing temperatures, changing precipitation designs have influenced terrestrial environment assembly and function, carbon and water balance, and finally production of crops [24, 25]. Several experiments have described the biological replies to CO₂ enhancement and their communication with ecological alteration at different levels [26]. The temperature has an important role in plant growth and development and regulates the several functions and enzymatic reactions in plants. Although, the increased value of temperature causes several abnormalities in plants such as reduces the chlorophyll contents, leaf growth, fresh and dry biomass, photosynthesis, and stomata limitations, inhibits the functions of several temperature-sensitive enzymes such as ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO). Impacts of eCO₂ and temperature combined stresses are very destructive at the reproductive phase. They may cause reduced pollination, spike sterility, less filling of grain, reduction of grain size and number, test weight, and yield potential of major crops [22].

It has been reported that eCO₂ promotes an initial stimulation of photosynthesis by an increase of substrate or RuBisCO carboxylation activity and self-consciousness of RuBisCO oxygenation, which might ultimately underwrite for advanced biomass in cereal crops [27, 28]. Nevertheless, growth responses over the long-standing under eCO₂ conditions include a reduction of photosynthesis measurements and several regulatory mechanisms to avoid potential damage by this condition. A regulatory mechanism to maintain the growth and expansion of plants consists of the equilibrium among manufacture and removal of reactive oxygen species (ROS) at the intracellular level. This balance is continued by both enzymatic and non-enzymatic antioxidant defense systems [29–31].

The enzymatic machinery involves numerous antioxidant enzymes, such as superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPX), guaiacol peroxidase (POX), peroxiredoxins (Prxs), and enzymes of the ascorbate-glutathione (AsAGSH) cycle. The AsAGSH includes different enzymes like ascorbate peroxidase (APX), monodehydroascorbate reductase (MDHAR), dehydroascorbate reductase (DHAR), and glutathione reductase (GR) [30, 32]. Modifications of this enzymatic antioxidant component have been reported in studies of cereal crops under eCO₂ conditions with contradictory results. Thus, the response of these antioxidants components is unclear with studies that find to increase [33], decreases [34], or no consistent alterations [35].

Besides, plant cells have non-enzymatic components that involve ascorbate and glutathione (GSH) along with phenolic acids, flavonoids, carotenoids anthocyanins, and phenolic composites. The free hydroxyl groups on the phenolic rings or the chromanol rings of these non-enzymatic compounds are responsible for their antioxidant properties [36]. The ring hydrogen atom can be given to free radicals, dropping and counteracting ROS. The phenolic compounds can lose a hydrogen atom which develops a free radical that is directly non-reactive by character delocalization in the entire ring assembly [28, 37]. Several studies reported only an increase in some of the individual phenolic compounds in cereal crops under high CO₂ conditions [38].

Furthermore, crop growth responses to eCO₂ rely on the tissue category, developmental stage as well as strength and duration of these conditions which also depend on the diversity of apparatuses of construction and purification of ROS, and the result of free radicals on antioxidants [39]. Several studies have reported a high production of hydrogen peroxide (H₂O₂) after exposure of crops to eCO₂, and the concentration of H₂O₂ is dependent on the duration of these conditions. Besides, H₂O₂ production differed among various cellular compartments [40]. The outcome of elevated aCO₂ meditation on growth and various antioxidant actions is superior in C₃ plants to C₄ plants [41, 42]. For example, rice as a C₃ plant is further sensitive to the variations of the aCO₂. However, there are inconsistent studies on the impacts of eCO₂ on the antioxidant responses and yield of rice changing from the reduction of the growth responses [43] to increment of antioxidants components and enhancement of growth [44, 45].

Kumar et al. [46] reported that rice plants under eCO₂ conditions showed modifications in electrolyte leakage, leaf water potential, proline, CAT, and POD activity as compared to ambient CO₂, which assisted the plant to battle contrary effects of stressful environments. Thus, these authors suggest that undesirable possessions on rice yield subsequent from abiotic stress conditions may be moderated by the eCO₂ meditations [46]. In agreement, leaves of a susceptible wheat cultivar (*Triticum aestivum* L.cv. Yitpi) infected with Barley yellow dwarf virus-PAV (*PadiAvenae* virus) and grown under eCO₂ presented that the eCO₂ conditions may decrease the oxidative stress caused by virus infection [47]. Nevertheless, more evidence for direct communicating possessions of eCO₂ and biotic and abiotic stress conditions in cereal crops is necessary.

3. eCO₂ mitigates oxidative stress in plants

Various environmental stresses induce the production of ROS, which triggers oxidative stress in plants [30, 48, 49]. The most common ROS are O₂^{•-}, [•]OH, and H₂O₂. In response to stressful conditions, H₂O₂ is mainly synthesized by photorespiration, beta (β)-oxidation, or due to the activity of nicotinamide adenine dinucleotide phosphate (NADPH) oxidase [30, 50]. The eCO₂ can efficiently reduce the ROS level by increasing RuBisCO carboxylation along with reducing photorespiratory H₂O₂ production. Several reports have provided indications regarding the eCO₂ influences on the mitigation of abiotic stress in plants [30, 51, 52]. Elevated CO₂ known to be induced plant growth by supplying additional Carbon (C) sources, subsequently alleviates abiotic stress in plants. Although the elevated CO₂ mediated particular physiological and molecular mechanisms related to abiotic stress alleviation are still to be explored. In the physiological aspect, supplying extra C by eCO₂ leads to induce stomatal closing, improves WUE that protect drought stress in plants [53]. However, abiotic stress-induced ROS (e.g., O₂^{•-}, [•]OH, and H₂O₂) and cellular oxidative damages (e.g., protein oxidation, lipid peroxidation) are involved in non-stomatal factors with metabolic changes [30, 54].

A systematic study of recently published articles addressed the two major hypotheses such as enhancement of antioxidant (antioxidant hypothesis) and reduction of stress impact (relaxation hypothesis) by reducing ROS generation in the plant under stressful conditions [30, 51, 55]. Several reports have been found in favor of the relaxation hypothesis in plants in response to eCO₂ under stressful conditions. The ROS level was found to be reduced by eCO₂ in plants under drought, and heat stresses through increasing RuBisCO carboxylation as well as reducing the level of photorespiratory H₂O₂ [51]. In the same study, glycine/serine (Gly/Ser) ration, glycolate oxidase (GO), and hydroxypyruvate reductase (HPR) level were evaluated as an indicator of photorespiration, which was found to be decreased in response to eCO₂ under drought and heat stresses. In barley, these all parameters were found at a lower level in response to eCO₂ [56]. Moreover, reduced photorespiration is correlated to the decreased level of NADPH oxidase. Therefore, a combined effect of lower photorespiration and NADPH oxidase responses may lead to reduce H₂O₂ in plants.

According to the antioxidant approach, the availability of additional C by eCO₂ enhances antioxidant molecules, which increase ROS scavenging activity as well as protects grapevine and tomato plants from abiotic stress induced-oxidative damages [57, 58]. More specifically, higher C availability due to eCO₂ may enhance the supply of defense molecules, which improve protection against oxidative injury (antioxidant hypothesis) under stressful conditions in plant cells. However, changes in antioxidant levels are not specific, or C₃ or C₄ based metabolism, or not for a particular group of species. It has been reported in C₄ plants that photorespiration mildly active, in which eCO₂ reduces ROS level as well as oxidative injury without alteration of antioxidants level. It suggests a distinct non-stomatal process that except antioxidant defense or reduces photorespiration processes. Besides, several reports have provided the evidence regarding eCO₂ reduces NADPH oxidase activity and ROS formation in mitochondria and chloroplast in plants [59, 60], but the activity of beta (β)-oxidation is still to be explored.

The ascorbate-glutathione (ASC-GSH) cycle is one of the major mechanisms for stress-induced H₂O₂ regulation. However, only limited reports have been found concerning eCO₂ mediated changes of ASC-GSH cycle components in plants under stress conditions. For example, HS alleviated through enhancement of DHAR, MDHAR, APX, and GR in tomato [58]. Similarly, GR and APX were found to be increased by eCO₂ in wheat under ozone stress [61]. Also, responses of ASC-GSH cycle components varied based on the plant species and experimental set-up. Therefore, additional studies are needed concerning the eCO₂ mediated oxidative stress alleviation as well as enhancement of ASC-GSH cycle components in plants under abiotic stresses.

4. eCO₂ improves photosynthesis under high temperature

Several researchers on the consequences of eCO₂ and stress factors on crops have been made using monitored situation amenities like plant growth chambers, free-air concentration enrichment (FACE) experiment, open-top chamber (OTC) or stimulated crop growth models [13]. The eCO₂ contributes to global warming causing alterations in the precipitations patterns, water scarcity, flood, and changes at extreme temperatures in several regions affecting the growth and development of plants [3]. However, the interactive effects of eCO₂ and environmental stress conditions on the development of crops and metabolism are not well documented. It has been shown the average reduction of stomatal conductance (20–30%), stomatal density (5–7%), stomatal developments, and increment in WUE (8–18%)

under eCO₂ conditions. However, these changes vary with the crop species, developmental stages, nature of stressors and duration, surrounding environments, and plant attributes [62]. Likewise, the interactions between an eCO₂ with stress factors are crucial to understanding the photosynthesis performance. Therefore, there is a considerable deviation in the light-saturated photosynthetic assimilation rate under eCO₂ rely upon the plant type, plant functional traits, micro or surrounding environment, and resource availability. For instance, the stimulation in photosynthetic assimilation rate under eCO₂ is varied from 30 to 80% (strong stimulations in C₃ species as compared to C₄) in FACE experiment or pot conditions but diminished in field conditions because of the integration of other multiple stressors such as drought, heat, flood and nutrient deficiencies.

In this regard, several studies were conducted to observe the combining effects of eCO₂ and drought and reveals that the growth mechanisms are distinct to the singular eCO₂ or drought. The combined stress resulted in the longer retention time of dissolved organic carbon (accumulation of soil organic C), induce invertase and catalase activity in the soil, and ameliorate stress conditions via improving plant physiological traits and activates feedback mechanisms [63]. Further, it limits the activity of some antioxidant enzymes such as proline and MDA content and stimulates others such as SOD, CAT, and GPX [46, 64]. Absciscic acid, calcium-dependent protein kinase and glutathione S-transferase (GST) play an important role in the amelioration of drought stress responses by inducing signaling mechanisms under the combined form. Conclusively, it is suggested that drought and HS generate ROS, and affect the antioxidant defense mechanism of plants, which might be ameliorated by the eCO₂ via stimulation of antioxidant defense enzymes [51, 62]. Similarly, eCO₂ combined with drought and HS regulates the sugars (starch, sucrose) and amino acids (alanine, pyruvate, arginine, glutamate) and secondary metabolites (coumaric acid, salicylic acid) metabolism, protective proteins and readjusted the metabolic, redox, and osmotic equilibrium of plants under combined eCO₂ and drought [65, 66].

The impact of combined eCO₂ and drought are varied with crop stage such as during the vegetative phase decreased the shoot elongation and leaf area, diminished mobilization of nutrients due to weak root growth, reduced stomatal conductance by increasing stomatal resistance and stomatal movements, plant hydraulic conductance, aquaporins, and reduce transpiration and RWC. However, at certain levels, it enhances the biomass allocations to the reproductive part, improves resource use efficiencies of plants including WUE, LUE, and NUE [67]. Similarly, eCO₂ and drought affect reproductive growth severely such as its impacts on the assimilate partitioning, pollen abortion, pollination, flower formation, panicle length, panicle weight, productive tiller number, seed formation, seed size, and yield potential of important crops [22, 68]. Besides the yield potential, eCO₂ decreased the grain quality via affecting macro- and micro-nutrients content such as phosphorus (P), Sulphur (S), and Iron (Fe), Zinc (Zn) contents of dryland legumes, which further associated with yield dilutions [23].

Several findings have stated the biological responses to CO₂ enhancement and their communication with conservation alteration at diverse levels [26]. In these aspects, the temperature has an important role in plant growth and development and regulates the several functions and enzymatic reactions in plants. Although, the increased value of temperature cause several abnormalities in plants such as change the emission of volatile organic compounds, reduce nitrogen uptake, chlorophyll contents, leaf growth, fresh and dry biomass, photosynthesis, and stomata limitations (by membrane damage and photosystem II (PSII) activity), enhance the activity of mitochondrial electron transport, stress proteins and plant growth regulators, limitation of several temperature-sensitive enzymes such as RuBisCO.

The combined effects of eCO₂ and temperature stresses are very destructive at the reproductive phase. They may cause changes in flowering time, pollination, spike sterility, less filling of grain, reduction of grain size and number, test weight, grain quality, and yield potential of major crops [20, 22, 69].

Besides the drought and HS (major), other stressors also interact with the eCO₂ under field conditions and influence the plant growth and development. For instance, eCO₂ and salt stress conditions influence the nitrogen metabolism, water balance, photosynthetic inhibitions, nutrient deficiency or toxicity, stomatal conductance, carbohydrate metabolism, phenolic enrichments, and generation of secondary metabolites [63, 70]. Similarly, limited nitrogen supply under eCO₂ modifies the C/N ratio, nitrogen metabolism, protein supply, protein structure, gene expression, sugar metabolism, and decreases antioxidant enzyme, amino acid synthesis, photosynthetic pigments, and elevated ROS, which influence the redox equilibrium and leads to early senescence in plants [71]. Likewise, under N limitations, the photosynthetic rate is more affected in C₃ species (because of more N requirement for RuBisCO synthesis), and eCO₂ could help in mitigation of N limitations by reducing photorespiration, elevating starch level, increase chloroplast size, higher stomatal resistance, mitochondrial respiration, metabolites and dilution of chlorophyll concentrations [62]. Therefore, the eCO₂ and stressors impacts have differed than singular stress, and up to a certain level of eCO₂ try to recover plants via inducing defense machinery, feedback mechanisms, activating secondary messenger signaling, and expression of stress proteins.

5. eCO₂ improves yield under high temperature

By the end of the 21st century, CO₂ is expected to rise from the current level 370 $\mu\text{mol}\cdot\text{mol}^{-1}$ to 540–970 $\mu\text{mol}\cdot\text{mol}^{-1}$, and about to grasp 550 $\mu\text{mol}\cdot\text{mol}^{-1}$ near 2050 and 750 $\mu\text{mol}\cdot\text{mol}^{-1}$ in 2100. In the meantime, Earth's global temperature will rise by about 2–4°C [72]. The increased concentration of CO₂ in the atmosphere would increase the temperature of Earth that is why global heating will develop, the most important aspect of upcoming climate variation. The relations between the temperature and CO₂ will have an intense effect on global agricultural production and the Earth's environment [73]. The rise in the atmospheric temperature and CO₂ would also accelerate the procedure of growth in plants [58, 74].

Climate change impacts on crop growth are becoming global concerns. They are particularly important for food supply and sustainable agricultural development [75, 76]. CO₂ concentration and temperature are two key factors affecting crop growth, development and yield [77]. Combined or individual possessions of temperature increase and eCO₂ meditation change on crop growth and yield during the recent decades have been observed [78]. For instance, the modeled improvements in soybean absorption of CO₂ with an increase in the growing season temperature, and aCO₂ hindered the photorespiration by 23–48%, which depends on the future climatic conditions [79].

The growth and distribution of crops are reduced by environmental factors like CO₂ and temperature. The production of the biomass of modern C₃ plants was decreased by 50% when it was grown at a low concentration of CO₂ (180–220 ppm), while the other conditions were optimal. Crops need the almost dual amount of water at 2°C increase in temperatures at a higher elevation of agricultural plains. Elevated CO₂ concentration increases the yield of the crop once the substrate for the photosynthesis process of leaf and the incline of CO₂ absorption of air increases. C₃ plants are more benefitted at eCO₂ than C₄ plants [80]. However, the doubling of CO₂ does not deteriorate the adverse effects of high temperature on the reproductive

growth of crops or fiber quality. Therefore, increased CO₂ concentration is associated with higher temperatures, crop yield, and quality that reduce particularly in areas where current temperatures are near to optimal [81].

Elevated CO₂ resulted in major changes in morpho-physiological restrictions. Besides, eCO₂ along with atmospheric temperatures during the phenological stages of rice cultivars showed contrasting results of the time of flowering and maturation such as eCO₂ in combination with the lower atmospheric temperature that stopped flowering in the CR-1014 cultivar while with the higher temperature increased grain yield in the Naveen cultivar [82].

Growth and photosynthesis of C₃ crops are enhanced when it is grown at a high level of CO₂, although, the degree of stimulation differs with temperature among cultivars as well as species. The probable decline in the transpiration process due to the partial closure of stomata in the eCO₂ level is largely invalid by the energy balance between the crop and its environment, which could result in total water use in similar climate conditions. The yield of seeds is increased by an increase in the CO₂ under the ideal temperature. On the other hand, at supra-optimal temperature, the yield of seeds is decreased under both raised and ambient CO₂. The yield of kidney bean decreased in that region where temperatures are at or above optimal conditions in combination with increased CO₂ concentration [83].

The effects of HS on the grain and biomass yield of plants depend on the duration and magnitude of HS. HS at the vegetative phase decreased the grain and biomass yield mostly by increasing plant growth and dropping the time obtainable to capture photons, and also by dropping the rate of photosynthesis [84]. At the anthesis or flowering phase, HS decreases the amount of grain due to pollen abortion. In contrast, at the grain-filling phase, HS decreases the heaviness of grain by restraining the translocation of assimilates, and marginally the period of grain-filling [85, 86].

Notably, eCO₂ may lessen the harmful influence of heat stress on the grain and biomass yield by inspiration of photosynthesis, defense of the photosynthetic devices from HS injury, and improvement in the water status of plants owing to reduced transpiration. Moreover, high levels of hexoses and sucrose in plants with eCO₂ are related to increase fertile florets and dry spike mass [87], and osmotic modification [88], which can develop heat stress tolerance [89]. It is hypothesized that HS at anthesis has a drastic impact on the grain yield and plant biomass, but it has a less impact eCO₂ than ambient CO₂ [90].

6. Conclusion

It is very important to know that the impacts of climate change on crop growth are becoming global concerns. Interactive effects of eCO₂ and environmental stress conditions on crop growth and carbon metabolism are not well predictable. The influence of eCO₂ with the connection of temperature is considerable on crops under stress environments. The CO₂ concentration improves the productivity of crops because of improved carbon exchange rates, and superior vegetative and reproductive growth. In contrast, crop productivity is decreased with increased temperatures. Hence, there is a need to develop genotypes that are different intolerant to various environments or to identify genotypes that perform better under predicted climate change. In this review, cereal genotypes have been characterized by differing responses to eCO₂ and HS and identified the mechanisms of tolerance to HS. It can promote the crop potential to assist the breeding program for the development of new genotypes tolerance to HS.

Conflicts of interest

The authors declare no conflicts of interest.

Disclosure statement

Authors declare that no conflict of interest could arise.

Author details

Ayman EL Sabagh^{1,2*}, Akbar Hossain³, Mohammad Sohidul Islam⁴,
Muhammad Aamir Iqbal⁵, Ali Raza⁶, Çetin Karademir², Emine Karademir²,
Abdul Rehman⁷, Md Atikur Rahman⁸, Rajesh Kumar Singhal⁹, Analía Llanes¹⁰,
Muhammad Ali Raza¹¹, Muhammad Mubeen¹², Wajid Nasim¹³,
Celaledin Barutçular¹⁴, Ram Swaroop Meena¹⁵ and Hirofumi Saneoka¹⁶

1 Department of Agronomy, Faculty of Agriculture, Kafrelsheikh University,
Kafrelsheikh, Egypt

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

2 Department of Field Crops, Faculty of Agriculture, Siirt University, Turkey

3 Bangladesh Wheat and Maize Research Institute, Dinajpur, Bangladesh

4 Department of Agronomy, Hajee Mohammad Danesh Science and Technology University, Dinajpur, Bangladesh

5 Department of Agronomy, University of Poonch Rawalakot Azadkashmir, Pakistan

6 Key Lab of Biology and Genetic Improvement of Oil Crops, Oil Crops Research Institute, Chinese Academy of Agricultural Sciences (CAAS), Wuhan, China

7 Department of Crop Sciences and Biotechnology, Dankook University, 119, Dandae-ro, Dongnam-gu, Cheonan-si, Korea

8 Grassland and Forage Division, National Institute of Animal Science, Rural Development Administration, Cheonan, Korea

9 ICAR-Indian Grassland and Fodder Research Institute, Jhansi, India

10 Plant Physiology Laboratory, Universidad Nacional de Río Cuarto (UNRC)-Instituto Nacional de Investigaciones Agrobiotecnológicas (INIAB), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Río Cuarto, Argentina

11 College of Agronomy, Sichuan Agricultural University, Chengdu, P.R. China

12 Department of Environmental Sciences, COMSATS University Islamabad, Vehari Campus, Pakistan

13 Department of Agronomy, University College of Agriculture and Environmental Sciences, The Islamia University of Bahawalpur (IUB), Bahawalpur, Pakistan

14 Department of Field Crops, Faculty of Agriculture, Cukurova University, Turkey

15 Department of Agronomy, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, UP, India

16 Graduate School of Integrated Sciences for Life, Hiroshima University, 1-4-4 Kagamiyama, Higashi-Hiroshima, Japan

*Address all correspondence to: aymanelsabagh@agr.kfs.edu.eg

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Raza A, Razzaq A, Mehmood SS, Zou X, Zhang X, Lv Y, Xu J. Impact of climate change on crops adaptation and strategies to tackle its outcome: A review. *Plants*. 2019;8:34
- [2] Hossain A, Farooq M, EL Sabagh A, Hasanuzzaman M, Erman M, Islam T. Morphological, Physiobiochemical and Molecular Adaptability of Legumes of Fabaceae to Drought Stress, with Special Reference to *Medicago sativa* L. In: Hasanuzzaman M, Araújo S, Gill S. (eds) *The Plant Family Fabaceae*. Springer, Singapore; 2020. pp. 289-317. https://doi.org/10.1007/978-981-15-4752-2_11.
- [3] IPCC. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, Pachauri RK, Meyer LA (eds.)]. IPCC, Geneva, Switzerland, 2014. p.151
- [4] Calzadilla A, Zhu T, Rehdanz K, Tol RSJ, Ringler C. Economy wide impacts of climate change on agriculture in Sub-Saharan Africa. *Ecological Economics*. 2013;93:150-165
- [5] Cairns JE, Sonder K, Zaidi PH, Verhulst N, Mahuku G, et al. Maize Production in a Changing Climate: Impacts, Adaptation, and Mitigation Strategies. *Advances in Agronomy*. 2012;114:1-65
- [6] Ziska L. Observed changes in soyabean growth and seed yield from *Abutilon theophrasti* competition as a function of carbon dioxide concentration. *Weed Research*. 2013;53:140-145
- [7] Easterling DR, Evans JL, Groisman PY, Karl TR, Kunkel KE, et al. Observed variability and trends in extreme climate events: a brief review. *Bulletin of American Meteorological Society*. 2000;81:417-425
- [8] Wallace JS. Increasing agricultural water use efficiency to meet future food production. *Agriculture, Ecosystems & Environment*. 2000;82:105-119
- [9] Johkan M, Oda M, Maruo T, Shinohara Y. Crop Production and Global Warming Impacts, Case Studies on the Economy, Human Health, and on Urban and Natural Environments. 2011;9:139-152. <http://www.intechopen.com>.
- [10] Young KJ, Long SP. Crop ecosystem responses to climatic change: maize and sorghum. In: Reddy KR, Hodges HF (Eds.). *Climate change and global crop productivity*, CABI International, Oxon, United Kingdom; 2000. pp. 107-131
- [11] Hunt R, Hand D, Hannah M, Neal A. Response to CO₂ enrichment in 27 herbaceous species. *Functional Ecology*. 1991;5:410-421
- [12] Rogers HH, Dahlman RC. Crop responses to CO₂ enrichment. CO₂ and biosphere. *Advances in vegetation Science*. 1993;14:117-131
- [13] Pathak H, Aggarwal PK, Singh SD. Climate change impacts, adaptations and mitigation in agriculture: methodology for assessment and application, Indian Agricultural Research Institute, New Delhi, India, 2012;pp. 1-302
- [14] Leakey A, Uribeblarrea M, Ainsworth E, Naidu S, Rogers A, et al. Photosynthesis, productivity and yield of maize are not affected by open-air elevation of CO₂ concentration in the absence of drought. *Plant Physiology*. 2006;140:779-790
- [15] Kim SH, Gitz DC, Sicher RC, Baker JT, Timlin DJ. Temperature

dependence of growth development, and photosynthesis in maize under elevated CO₂. *Environment and Experimental Botany*. 2007;**61**:224-236

[16] Vanaja M, Maheswari M, Jyothi Lakshmi N, Sathish P, Yadav SK, Salini K, Vagheera P, Vijay Kumar G, Razak A. Variability in growth and yield response of maize genotypes at eCO₂ concentration. *Advances in Plants & Agriculture Research*. 2015;**2**:42. DOI: 10.15406. apar.2015.02.00042

[17] Meena RS, Kumar V, Yadav GS, Mitran T. Response and interaction of *Bradyrhizobiumjaponicum* and *Arbuscularmycorrhizalfungi* in the soybean rhizosphere: A review. *Plant Growth Regulation*. 2018;**84**:207-223

[18] Wang D, Heckathorn SA, Barua D, Joshi P, Hamilton EW, LaCroix JJ. Effects of eCO₂ on the tolerance of photosynthesis to acute heat stress in C₃, C₄, and CAM species. *American Journal of Botany*. 2008;**95**:165-176. <https://doi.org/10.3732/ajb.95.2.165>.

[19] Meena RS, Lal R, Yadav GS. Long-term impact of topsoil depth and amendments on carbon and nitrogen budgets in the surface layer of an Alfisol in Central Ohio. *Catena*, 2020;**194**:104752.<https://doi.org/10.1016/j.catena.2020.104752>

[20] Jagadish SVK, Bahuguna RN, Djanaguiraman M, Gamuyao R, Prasad PV, Craufurd PQ. Implications of high temperature and elevated CO₂ on flowering time in plants. *Frontiers in Plant Science*. 2016;**7**:111. <http://dx.doi.org/10.3389/fpls.2016.00913>.

[21] Xu Z, Jiang Y, Jia B, Zhou G. Elevated-CO₂ response of stomata and its dependence on environmental factors. *Frontiers in Plant Science*. 2016;**7**:657. <https://doi.org/10.3389/fpls.2016.00657>

[22] Kadam NN, Xiao G, Melgar RJ, Bahuguna RN, Quinones C,

Tamilselvan A, Prasad PVV, Jagadish KS. Agronomic and physiological responses to high temperature, drought, and elevated CO₂ interactions in cereals. *Advances in Agronomy*. 2014;**127**:111-156

[23] Salim N, Raza A. Nutrient use efficiency (NUE) for sustainable wheat production: a review. *Journal of Plant Nutrition*. 2020;**43**:297-315

[24] Lavania D, Dhingra A, Siddiqui MH, Al-Whaibi MH, Grover A. Current status of the production of high temperature tolerant transgenic crops for cultivation in warmer climates. *Plant Physiology and Biochemistry*. 2015;**86**:100-108. Doi:10.1016/j.plaphy.2014.11.019

[25] Raza A, Ashraf F, Zou X, Zhang X, Tosif H. Plant Adaptation and Tolerance to Environmental Stresses: Mechanisms and Perspectives. In *Plant Ecophysiology and Adaptation under Climate Change: Mechanisms and Perspectives I*. Springer, Singapore, 2020;pp. 117-145

[26] Rodrigues WP, Martins MQ, Fortunato AS, Rodrigues AP, Semedo JN, Simões-Costa MC, et al. Long-term elevated air [CO₂] strengthens photosynthetic functioning and mitigates the impact of supra-optimal temperatures in tropical *Coffea arabica* and *C. canephora* species. *Global Change Biology*. 2016;**22**:415-431. Doi:10.1111/gcb.13088

[27] Aranjuelo I, Sanz-Sáez Á, Jauregui I, Irigoyen JJ, Araus JL, Sánchez-Díaz M, Erice G. Harvest index, a parameter conditioning responsiveness of wheat plants to elevated CO₂. *Journal of Experimental Botany*. 2013;**64**:1879-1892

[28] Goufo P, Trindade H. Rice antioxidants: phenolic acids, flavonoids, anthocyanins, proanthocyanidins, tocopherols, tocotrienols, γ -oryzanol,

and phytic acid. Food Science & Nutrition. 2014;2:75-104

[29] Horvat D, Šimić G, Drezner G, Lalić A, Ledenčan T, Tucak M, ... Zdunić Z. Phenolic Acid Profiles and Antioxidant Activity of Major Cereal Crops. Antioxidants. 2020;9:527

[30] Hasanuzzaman M, Bhuyan MHM, Zulfiqar F, Raza A, Mohsin SM, Mahmud JA, ... Fotopoulos V. Reactive Oxygen Species and Antioxidant Defense in Plants under Abiotic Stress: Revisiting the Crucial Role of a Universal Defense Regulator. Antioxidants. 2020;9:681

[31] EL Sabagh A, Hossain A, Islam MS, Fahad S, Ratnasekera D, Meena RS, Wasaya A, Yasir TA, Ikram M, Mubeen M, Fatima M, Nasim W, Çığ A, Çığ F, Erman M, Hasanuzzaman M. Nitrogen Fixation of Legumes Under the Family Fabaceae: Adverse Effect of Abiotic Stresses and Mitigation Strategies. In: Hasanuzzaman M, Araújo S, Gill S. (eds) The Plant Family Fabaceae. Springer, Singapore; 2020. pp. 75-111. https://doi.org/10.1007/978-981-15-4752-2_4.

[32] Mittler R, Vanderauwera S, Gollery M, Van Breusegem F. Reactive oxygen gene network of plants. Trends in Plant Science. 2004;9:490-498

[33] diToppo LS, Marabottini R, Badiani M, Raschi A. Antioxidant status in herbaceous plants growing under elevated CO₂ in mini-FACE rings. Journal of Plant Physiology. 2002;159:1005-1013

[34] Pérez-López U, Robredo A, Lacuesta M, Sgherri C, Mena-Petite A, Navari-Izzo F, Muñoz-Rueda A. Lipoic acid and redox status in barley plants subjected to salinity and elevated CO₂. Physiologia Plantarum. 2010;139:256-268

[35] Tausz-Posch S, Borowiak K, Dempsey RW, Norton RM, Seneweera S,

Fitzgerald GJ, Tausz M. The effect of elevated CO₂ on photochemistry and antioxidative defence capacity in wheat depends on environmental growing conditions-A FACE study. Environmental and Experimental Botany. 2013;88:81-92

[36] Landete JM. Dietary intake of natural antioxidants: vitamins and polyphenols. Critical Reviews in Food Science and Nutrition. 2013;53:706-721

[37] Visioli F, Lastra CADL, Andres-Lacueva C, Aviram M, Calhau C, Cassano A, ...Llorach R. Polyphenols and human health: a prospectus. Critical Reviews in Food Science and Nutrition. 2011;51:524-546

[38] Cho JY, Lee HJ, Kim GA, Kim GD, Lee YS, Shin SC, Park KY, Moon JA. Quantitative analyses of individual g-oryzanol (sterylferulates) in conventional and organic brown rice (*Oryza sativa* L.). Journal of Cereal Science. 2012;55:337e343

[39] Zinta G, AbdElgawad H, Domagalska MA, Vergauwen L, Knapen D, Nijs I, ... Asard H. Physiological, biochemical, and genome-wide transcriptional analysis reveals that elevated CO₂ mitigates the impact of combined heat wave and drought stress in *Arabidopsis thaliana* at multiple organizational levels. Global Change Biology. 2014;20:3670-3685

[40] Slesak I, Libik M, Karpinska B, Karpinski S, Misalski Z. The role of hydrogen peroxide in regulation of plant metabolism and cellular signalling in response to environmental stresses. Acta Biochimica Polonica. 2007;54:39-50

[41] Hymus GJ, Baker NR, Long SP. Growth in elevated CO₂ can both increase and decrease photochemistry and photoinhibition of photosynthesis in a predictable manner. Dactylis glomerata grown in two levels

of nitrogen nutrition. *Plant Physiology*. 2001;**127**:1204-1211

[42] Cunliffe J, Charles M, Jones G, Osborne CP. Reduced plant water status under sub-ambient p CO₂ limits plant productivity in the wild progenitors of C₃ and C₄ cereals. *Annals of Botany*. 2016;**118**:1163-1173

[43] Satapathy BS, Duary B, Saha S, Pun KB, Singh T. Effect of weed management practices on yield and yield attributes of wet direct seeded rice under lowland ecosystem of Assam. *ORYZA-An International Journal on Rice*. 2017;**54**:29-36

[44] Bhattacharyya P, Roy KS, Neogi S, Dash PK, Nayak AK, Mohanty S, ... Rao KS. Impact of elevated CO₂ and temperature on soil C and N dynamics in relation to CH₄ and N₂O emissions from tropical flooded rice (*Oryza sativa* L.). *Science of the Total Environment*. 2013;**461**:601-611

[45] Roy S, Banerjee A, Mawkhlieng B, Misra AK, Pattanayak A, Harish GD, ... Bansal KC. Genetic diversity and population structure in aromatic and quality rice (*Oryza sativa* L.) landraces from North-Eastern India. *PloS One*. 2015;**10**:e0129607

[46] Kumar A, Nayak AK, Sah RP, Sanghamitra P, Das BS. Effects of elevated CO₂ concentration on water productivity and antioxidant enzyme activities of rice (*Oryza sativa* L.) under water deficit stress. *Field Crops Research*. 2017;**212**:61-72

[47] Vandegheer RK, Powell KS, Tausz M. Barley yellow dwarf virus infection and elevated CO₂ alter the antioxidants ascorbate and glutathione in wheat. *Journal of Plant Physiology*. 2016;**199**:96-99

[48] Sharma P, Jha AB, Dubey RS, Pessarakli M. Reactive oxygen species, oxidative damage, and antioxidative

defense mechanism in plants under stressful conditions. *Journal of Botany*. 2012;**2012**:217037:<http://doi.org/10.1155/2012/217037>.

[49] EL Sabagh A, Hossain A, Islam MS, Barutcular C, et al. Drought and Heat Stress in Cotton (*Gossypium hirsutum* L.): Consequences and Their Possible Mitigation Strategies. In: Hasanuzzaman M. (eds) *Agronomic Crops*. Springer, Singapore. 2020a. https://doi.org/10.1007/978-981-15-0025-1_30

[50] Apel K, Hirt H. Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annual Review of Plant Biology*. 2004;**55**:373-399

[51] AbdElgawad H, Farfan-Vignolo ER, De Vos D, Asard H. Elevated CO₂ mitigates drought and temperature-induced oxidative stress differently in grasses and legumes. *Plant Science*. 2015;**231**:1-10. [10.1016/j.plantsci.2014.11.001](https://doi.org/10.1016/j.plantsci.2014.11.001)

[52] AbdElgawad H, Zinta G, Beemster GTS, Janssens IA, Asard H. Future climate CO₂ levels mitigate stress impact on plants: increased defense or decreased challenge? *Frontiers in Plant Science*. 2016;**7**:556. Doi: [10.3389/fpls.2016.00556](https://doi.org/10.3389/fpls.2016.00556).

[53] Xu Z, Shimizu H, Yagasaki Y, Ito S, Zheng Y, Zhou G. Interactive effects of elevated CO₂, drought, and warming on plants. *Journal of Plant Growth Regulation*. 2013;**32**:692-707

[54] Ghannoum O. C₄ photosynthesis and water stress. *Annals of Botany*. 2009;**103**:635-644

[55] Yassin M, El Sabagh A, Mekawy AMM, Islam MS, Hossain A, Barutcular C, Alharby H, Bamagoos A, Liu L, Ueda A, Saneoka H Comparative performance of two bread wheat (*Triticum aestivum* L.) genotypes under salinity stress. *Applied Ecology and*

Environmental Research. 2019;**17**:5029-5041. https://doi.org/10.15666/aer/1702_50295041

[56] Fair P, Tew J, Cresswell C. Enzyme activities associated with carbon dioxide exchange in illuminated leaves of *Hordeum vulgare* L. II. Effects of external concentrations of carbon dioxide and oxygen. Annals of Botany. 1973;**37**:1035-1039

[57] Salazar-Parra C, Aguirreolea J, Sánchez-Díaz M, Irigoyen JJ, Morales F. Climate change (elevated CO₂, elevated temperature and moderate drought) triggers the antioxidant enzymes' response of grapevine cv. Tempranillo, avoiding oxidative damage. Physiologia Plantarum. 2012;**144**:99-110

[58] Li W, Zhu Q, Wang Y, Wang S, Chen X, Zhang D, Wang W. The relationships between physiological and biochemical indexes and the yield characteristics of rice under high temperature stress. 2013;**29**:

[59] Booker FL, Reid CD, Brunschön-Harti S, Fiscus EL, Miller JE. Photosynthesis and photorespiration in soybean [*Glycine max* (L.) Merr.] chronically exposed to elevated carbon dioxide and ozone. Journal of Experimental Botany. 1997;**48**:1843-1852

[60] Lin J-S, Wang G-X. Doubled CO₂ could improve the drought tolerance better in sensitive cultivars than in tolerant cultivars in spring wheat. Plant Science. 2002;**163**:627-637

[61] Rao MV, Hale BA, Ormrod DP. Amelioration of ozone-induced oxidative damage in wheat plants grown under high carbon dioxide (Role of antioxidant enzymes), Plant Physiology. 1995;**109**:421-432

[62] Xu Z, Jiang Y, Zhou G. Response and adaptation of photosynthesis, respiration, and antioxidant systems to

elevated CO₂ with environmental stress in plants. Frontiers in Plant Science. 2015;**6**:701

[63] Dietzen CA, Larsen KS, Ambus PL, Michelsen A, Arndal MF, Beier C, Reinsch S, Schmidt IK. Accumulation of soil carbon under elevated CO₂ unaffected by warming and drought. Global Change Biology. 2019;**25**:2970-2977

[64] Li S, Li Y, Gao Y, He X, Zhang D, Liu B, Li Q. Effects of CO₂ enrichment on non-structural carbohydrate metabolism in leaves of cucumber seedlings under salt stress. Scientia Horticulturae. 2020;**265**:109275

[65] Zinta G, AbdElgawad H, Peshev D, Weedon JT, Van den Ende W, Nijs I, Janssens IA, Beemster GT, Han A. Dynamics of Metabolic Responses to Combined Heat and Drought Spells in *Arabidopsis thaliana* under Ambient and Rising Atmospheric CO₂. Journal of Experimental Botany. 2018;**69**:2159-2170. doi: 10.1093/jxb/ery055.

[66] Li M, Li Y, Zhang W, Li S, Gao Y, Ai X, Zhang D, Liu B, Li Q. Metabolomics analysis reveals that elevated atmospheric CO₂ alleviates drought stress in cucumber seedling leaves. Analytical Biochemistry. 2018;**559**:71-85.

[67] Oliveira MFD, Marengo RA. Gas exchange, biomass allocation and water-use efficiency in response to elevated CO₂ and drought in andiroba (*Carapasurinaensis*, Meliaceae). iForest-Biogeosciences and Forestry. 2019;**12**:61-68

[68] Mphande W, Nicolas ME, Seneweera S, Bahrami H. Dynamics and contribution of stem water-soluble carbohydrates to grain yield in two wheat lines contrasted under drought and elevated CO₂ conditions. Journal of Plant Physiology. 2016;**214**:1037-1058

- [69] Vicente R, Bolger AM, Martinez-Carrasco R, Pérez P, Gutiérrez E, Usadel B, Morcuende R. De novo transcriptome analysis of durum wheat flag leaves provides new insights into the regulatory response to elevated CO₂ and high temperature. *Frontiers in Plant Science*. 2019;**10**:1605.
- [70] Sgherri C, Pérez-López U, Micaelli F, Miranda-Apodaca J, Mena-Petite A, Muñoz-Rueda A, Quartacci MF. Elevated CO₂ and salinity are responsible for phenolics-enrichment in two differently pigmented lettuces. *Plant Physiology and Biochemistry*. 2017;**115**:269-278
- [71] Agüera E, De la Haba P. Leaf senescence in response to elevated atmospheric CO₂ concentration and low nitrogen supply. *Biologia Plantarum*. 2018;**62**:401-408
- [72] Collins M, Knutti R, Arblaster J, Dufresne JL, Fichefet T, Friedlingstein P,Krinner G. Long-term climate change: projections, commitments and irreversibility. In *Climate Change 2013. The Physical Science Basis: Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press. 2013. pp. 1029-1136
- [73] Rosenzweig C, Elliott J, Deryng D, Ruane AC, Müller C, Arneth A,Khabarov N. Assessing agricultural risks of climate change in the 21st century in a global gridded crop model intercomparison. *Proceedings of the National Academy of Sciences*. 2014;**111**:3268-3273
- [74] Cai C, Yin X, He S, Jiang W, Si C, Struik PC, . . .Xiong Y. Responses of wheat and rice to factorial combinations of ambient and elevated CO₂ and temperature in FACE experiments. *Global Change Biology*. 2016;**22**:856-874
- [75] Tan K, Zhou G, Lv X, Guo J, Ren S. Combined effects of elevated temperature and CO₂ enhance threat from low temperature hazard to winter wheat growth in North China. *Scientific Reports*. 2018;**8**:1-9
- [76] EL Sabagh A, Hossain A, Barutçular C, Gormus O, Ahmad Z, Hussain S, Islam MS, Alharby H, Bamagoos A, Kumar N, Akdeniz A, Fahad S, Meena RS, Abdelhamid M, Wasaya A, Hasanuzzaman M, Sorour S, Saneoka H. Effects of drought stress on the quality of major oilseed crops: implications and possible mitigation strategies—a review. *Applied Ecology and Environmental Research*. 2019;**17**:4019-4043. https://doi.org/10.15666/aeer/1702_40194043
- [77] Mubeen M. et al. Evaluating the climate change impact on crop water requirement of cotton- wheat in semi-arid conditions using DSSAT model. *Journal of Water and Climate Change*. 2019; 201979. doi: 10.2166/wcc.2019.179.
- [78] Hatfield JL, Boote KJ, Kimball B, Ziska L, Izaurralde RC, Ort D, . . . Wolfe D. Climate impacts on agriculture: implications for crop production. *Agronomy Journal*. 2011;**103**:351-370
- [79] Walker BJ, VanLoocke A, Bernacchi CJ, Ort DR. The costs of photorespiration to food production now and in the future. *Annual Review of Plant Biology*. 2016;**67**:107-129.
- [80] Madan P, Jagadish S, Craufurd P, Fitzgerald M, Lafarge T, Wheeler T. Effect of elevated CO₂ and high temperature on seed-set and grain quality of rice. *Journal of Experimental Botany*. 2012;**63**:3843-3852
- [81] Reddy KR, Vara Prasad P, Kakani VG. Crop responses to elevated carbon dioxide and interactions with

temperature: cotton. Journal of Crop Improvement. 2005;**13**:157-191

[82] Ziska LH. et al. Food security and climate change: on the potential to adapt global crop production by active selection to rising atmospheric carbon dioxide. Proceedings of the Royal Society B: Biological Sciences. 2012;**279**:4097-4105 <http://doi.org/10.1098/rspb.2012.1005>

[83] Prasad PV, Boote KJ, Allen Jr LH, Thomas JM. Effects of elevated temperature and carbon dioxide on seed-set and yield of kidney bean (*Phaseolus vulgaris* L.). Global Change Biology. 2002;**8**:710-721

[84] Lobell DB, Gourdji SM. The influence of climate change on global crop productivity. Plant Physiology. 2012;**160**:1686-1697. Doi: <https://doi.org/10.1104/pp.112.208298>

[85] Farooq M, Bramley H, Palta JA, Siddique KHM. Heat stress in wheat during reproductive and grain-filling phases. Critical Reviews in Plant Sciences. 2011;**30**:491-507. DOI: 10.1080/07352689.2011.615687

[86] Prasad PVV, Djanaguiraman M. Response of floret fertility and individual grain weight of wheat to high temperature stress: sensitive stages and thresholds for temperature and duration. Functional Plant Biology. 2014;**41**:1261-1269. <http://dx.doi.org/10.1071/FP14061>

[87] Dreccer MF, Wockner KB, Palta JA, McIntyre CL, Borgognone MG, Bourgault M, Reynolds M, Miralles DJ. More fertile florets and grains per spike can be achieved at higher temperature in wheat lines with high spike biomass and sugar content at booting. Functional Plant Biology 2014;**41**:482-495. <http://dx.doi.org/10.1071/FP13232>.

[88] Wahid A, Gelani S, Ashraf M, Foolad MR. Heat tolerance in plants:

an overview. Environmental and Experimental Botany. 2007;**61**:199-223

[89] Shanmugam S, Kjaer KH, Ottosen CO, Rosenqvist E, Kumari Sharma D, Wollenweber B. The alleviating effect of elevated CO₂ on heat stress susceptibility of two wheat (*Triticum aestivum* L.) cultivars. Journal of Agronomy and Crop Science. 2013;**199**:340-350

[90] Chavan SG, Duursma RA, Tausz M, Ghannoum O. ECO₂ alleviates the negative impact of heat stress on wheat physiology but not on grain yield. Journal of Experimental Botany. 2019;**70**:6447-6459. doi:10.1093/jxb/erz386