We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Alcohol Consumption Practice and Associated Risk Factors among University of Limpopo Students

*Mbelege Rosina Nkwana, Gift Makaleng,
Mafoloa Suzan Monyeki, Hlengani James Siweya
and Kotsedi Daniel Monyeki*

Abstract

In South Africa, substance rehabilitation centers found alcohol as the primary substance abuse in eight out of the nine provinces. This study aimed to investigate the prevalence of alcohol use and associated risk factors among University of Limpopo students with mean age of 21.5 years. This was a cross-sectional study and constituted a total of 500 female students and 417 male students from the University of Limpopo. Logistic regression was used to calculate the association between alcohol consumption and its predictors. Information about their alcohol consumption, the type of alcohol and the practice related to alcohol consumption was collected using a validated questionnaire. Tobacco and marijuana product use were significantly ($P < 0.05$) associated with alcohol use among University of Limpopo students for unadjusted (OR ranges 4.31 95%CI 2.71 6.87 to 4.92 95%CI 3.16 7.70) and after adjusting for age gender and amount of money deposited into the student account by the bursary scheme (OR ranges from 4.14 95%CI 2.59 6.62 to 4.50 95%CI 2.87 7.06). Tobacco product use, marijuana use and enhancing interpersonal relationships are major risk factors associated to current alcohol use. Future studies are required to investigate the development of lifestyle and cardiovascular disease over time among University of Limpopo students.

Keywords: alcohol, tobacco, marijuana, students, university of Limpopo

1. Introduction

A university living setting provides a unique opportunity for students who enter this environment for the first time. Yet, living in a new environment is associated with stress and changes of lifestyle [1]. At the university level, students adjust to a certain level of being an adult and experiencing freedom. These students create a social and cultural environment by drinking alcohol to obtain a positive outcome (e.g. enhancement, coping) or to avoid a negative outcome (like peer rejection) [2]. This environment could further create unhealthy lifestyles such that students are more involved in risky behaviour and serious drinking problems than their peers

who are not enrolled in the University [2–4]. Unhealthy behaviour such as tobacco product use, marijuana use, injuries and interpersonal problems are the vital risk factors associated with excessive alcohol consumption. Besides, alcohol use among university students has been associated with poor academic performance, personal injuries, loss of memory, temporary illness (such as nausea, vomiting, and hang-over), unprotected sexual activity, acquaintance rape, sexual violence and other problems like suicide, impaired driving skills, vandalism or fights [5].

The high percentage of alcohol use and abuse in the university environment by students is viewed as representing the prime drinking years for individuals [6]. For example, Young and De Klerk. [7] reported that more than 50% of Rhodes University student respondents who participated in their study were dependent on alcohol. The prevalence of alcohol use in a university in the South-East of Nigeria was reported to be 78.4% [8].

The prevalence of alcohol use by university students in South Africa has been found to range from 20% to 80% in 2009 by Peltzer & Ramlagan [9]. The prevalence of alcohol drinking at the University of Venda, South Africa was reported to be 65% of the current drinkers [10]. In South Africa, substance rehabilitation centers found alcohol as the primary substance abuse in eight out of the nine provinces [9]. Excessive alcohol consumption during University year's impacts negatively on student lifestyles even on post-University level, leading to accidents, health problems, and personal problems [11].

Alcohol drinking may be responsive to internal rewards such as the manipulation of management of one's own internal emotional state and external reward such as social acceptance [2]. Various factors affect the alcohol drinking patterns of students in the university environment. Some university students prefer beer and other drugs for economic reasons. Regardless of the various alcoholic beverages like cider, wine, spirits, champagne, whisky, gin or brandy [12]. Gender differences in alcohol use was reported among university students in the USA with a significantly high number of males compared to females drinking alcohol [13]. Furthermore, staying on-campus has been linked with excessive alcohol drinking than staying off-campus [14]. Little is known about the prevalence of alcohol use and associated risk factors among the University of Limpopo. This study aimed to investigate the prevalence of alcohol use and associated risk factors among University of Limpopo students aged 17 to 43 years.

2. Method

2.1 Geographical area

The study was conducted at the University of Limpopo, that was previously known as the University of The North (Sovenga). The University of Limpopo is situated at the foothill of Hwiti (Wolkberg range) in Mankweng, midway between Polokwane and Makgoebaskloof in Limpopo province, South Africa. The research was approved by the University of Limpopo research ethics committee (TREC/61/2019: IR).

2.2 Sampling and sample size

All the participants of the study were full-time students enrolled for the academic year 2019 in University of Limpopo. The participants represented the four available faculties (Science and Agriculture, Humanities, Management and Law, and Health Sciences). The total number of participants was $n = 917$ of which 45.5%

(n = 417) were males and 54.5% (n = 500) were females (mean age of 21.5 years). A convenient sampling method was used to have easy access to the participants.

2.3 Instruments

The questionnaire used in this study was based on questions, which have been used in the Ellisras Longitudinal Study and other studies [15, 16]. The questionnaire was shared with experts to ensure content, face and construct validity. The questionnaire was then revised and piloted with a sample of students to make sure it was valid, reliable, acceptable and accurately understood.

The questionnaire comprised of three sections. The first section of the questionnaire included of participants's age, gender, marital status and place of residence, field of study, year of entering the university and the level of study. The academic level was categorised as lower level (first and second year), moderate high (third-year level) and high level (postgraduate level).

Current alcohol drinkers were defined as anyone who drinks alcohol regularly for the past 30 days. The following question was asked: "During the past 30 days, have you had at least one drink of any alcoholic beverage such as beer, wine, a malt beverage, or liquor regularly?" The number of days in the past month where you had a drink between 07:30 and 16:00 and the frequency of drinking alcohol regularly the past 30 days were classified into: Less than four times in the past 30 days, 2 to 3 days a week in the past 30 days and more than four days a week the past 30 days. This was classified into the less than four times in 30 days. The onset (initiation age) age for alcohol drinker use was determined by the question "If yes, indicate how old you were when you first tried this _____. How old were you when you first drank alcohol regularly _____." Onset age was grouped as less than 15 years, between 15 and 19 years and over 20 years.

2.4 Other associated factors

All studied characteristics together with peer relation, personal enjoyment and tension reduction were chosen based upon literature that had identified associations between these variables and binge drinking [17, 18]. The selection questions included: I drink alcohol because it will boost my creativity (Yes/NO)? I drink alcohol because I want to be popular (Yes/NO)? I drink alcohol because it helps me to face difficulties with confidence (Yes/NO)? I drink alcohol because it would comfort me when I get blamed (Yes/NO)?

Tobacco and marijuana products use was assessed using the following questions: Does anyone you live with use tobacco products (Yes/No)? Does anyone you live with smoke hubbly bubbly (Yes/No)? Does anyone you live with smoke weed/marijuana (Yes/No)? Does anyone you live with drink alcohol (Yes/No)?

2.5 Statistical analysis

Descriptive statistics including frequency distribution and percentage frequencies were used to determine the prevalence of current alcohol use among University of Limpopo students. A chi-squared test was used to compare sets of nominal data that had larger frequency counts while the Fisher's exact test was used when frequency cells were small (less than five or ten) between genders [19, 20]. Logistic regression was used to determine the associated risk factors for current alcohol use among the University of Limpopo students. All statistical analyses were performed using SPSS version 25. The statistical significance was set at $P < 0.05$.

3. Results

Table 1 shows the descriptive statistics for positive response on alcohol consumption of the University of Limpopo students aged 17 to 43 years. The prevalence of current regular alcohol use was significantly ($p < 0.021$) higher for males (61.9%) compared to females (47.8%). Students who first consumed alcohol at age of less than 15 years were 24.7% males and 8.8% females, the difference was

	Males		Females	
	%	(n)	%	(n)
Faculties of the University of Limpopo				
Science and Agriculture	33.8	(141)	19.6	(98)
Humanities	26.9	(112)	23.8	(119)
Management and Law	17.0	(71)	13.0	(65)
Health Sciences	22.3	(93)	43.6	(218)
Bursaries Received				
Receiving funding from other bursaries	11.0	(46)	9.2	(46)
Receiving funding from NSFAS	70.3	(293)	80.4	(402)
Students Study level				
Lower level (First and Second year)	56.8	(237)	53.8	(269)
Moderate high level (third-year level)	23.0	(96)	25.6	(128)
High level (post graduate level)	20.1	(84)	20.6	(103)
Alcohol consumption				
Have you ever drunk alcohol?	74.3	(310)	62.8	(314)
Age when you had first drink of alcohol				
Less than 15 years	31.4	(131)	14.2	(71)
15 to 19 years	35.7	(149)	40.0	(200)
20 years and above	7.2	(30)	8.6	(43)
Are you currently drinking alcohol regularly during the past 30 days	61.9 [*]	(258)	47.8 [*]	(239)
Age when you first started drinking alcohol regularly during the past 30 days				
Less than 15 years	24.7 [*]	(103)	8.8 [*]	(44)
15 to 19 years	30.0	(125)	30.8	(154)
20 years and above	7.2	(30)	8.2	(41)
Types of alcoholic drinks you drink regularly in the past 30 days				
Beer	44.4 [*]	(185)	9.6 [*]	(48)
Cider	35	(146)	30.2	(151)
Wine	24.9	(104)	29.2	(146)
Spirit	29.3 [*]	(122)	12.8 [*]	(64)
Home made	8.4 [*]	(35)	3.0 [*]	(15)
Champagne	15.8	(66)	12.4	(62)

^{*}≥0.05; ^{**}≥0.001.

Table 1.
Descriptive statistics for positive response on alcohol consumption of the University of Limpopo students aged 17 to 43 years.

significant ($P < 0.05$). Many males (44.4% and 29.3%) in the current study significantly ($P < 0.05$) prefer beer and spirits compared to females (9.6% and 12.8%), respectively. Females preferred wine than males (29.2% and 24,9%) respectively. Though, the difference was not significant. The use of homemade alcohol was significantly ($p < 0.05$) higher for males (8.4%) compared to females (3.0%).

Table 2 presents the prevalence of positive response on the frequency of drinking and reasons for drinking alcohol among University of Limpopo students aged 17 to 43 years. A significantly high number of females (27.6%) drink alcohol fewer than 4 times in the past 30 days than males (21.6%). There was a significant ($P < 0.05$) number of males (13.4%) who drink alcohol regularly between the period of 07:30 and 16:00 2 to 3 days a week in the past month compared to females (7.4%). The prevalence of students who drink alcohol for popularity and bosting confidence is also higher in males (34.8% and 23.5%) than in females (1.8%, and 7.8%), respectively. The prevalence of tobacco and marijuana products use among relatives/friends who live with males students were significantly higher (ranges from 16.8 to 42.2%) compared to those who live with females students (prevalence ranges from 10.6 to 27.4%).

Table 3 shows the logistic regression (Odds ratio and 95% confidence interval and p-value) for the association of current regular alcohol drinking use and the associated risk factors for University of Limpopo students aged 17 to 43 years. Tobacco and marijuana product use of relatives/friends living with students were

	Males		Females	
	%	(n)	%	(n)
Rate of drinking alcohol regularly the past 30 days				
< than 4 times in 30 days	21.6*	(90)	27.6*	(138)
2 to 3 days a week in the past 30 days	11.8	(49)	6.8	(34)
> than 4 days a week in the past 30 days	28.5*	(119)	13.4*	(67)
Number of days in the past month where you had a drink between 07:30 and 16:00				
< than 4 times in the past 30 days	29.0	(121)	22.4	(112)
2 to 3 days a week in the past 30 days	13.4*	(56)	7.4*	(37)
> than 4 days a week the past 30 days	7.2	(30)	4.0	(20)
Reasons for drinking alcohol				
I drink alcohol because it will enhance my creative ability	6.7	(28)	6.6	(33)
I drink alcohol because I want to be popular	34.8*	(145)	1.8*	(9)
I drink alcohol because it helps me to face difficulties with confidence	23.5*	(99)	7.8*	(39)
I drink alcohol because it would ease me when I get blamed	18*	(75)	4.6*	(23)
Tobacco and marijuana product use				
Does anyone you live with use tobacco product?	25.2*	(105)	14.2*	(71)
Does anyone you live with smoke hubbly bubbly?	16.8*	(70)	11.4*	(57)
Does anyone you live with smoke weed/marijuana?	22.8*	(95)	10.6*	(53)
Does anyone you live with drink alcohol?	42.2*	(178)	27.4*	(137)

* ≥ 0.05 ; ** ≥ 0.001 .

Table 2.
Positive response on frequency of drinking and reasons for drinking alcohol among University of Limpopo students aged 17 to 43 years.

	Unadjusted		Adjusted for age, gender and receiving money into own Bank account					
	OR	P-value	(95%CI)		OR	P-value	95 %CI	
Educational level								
Lower level	0.96	0.765	(0.74	1.25)	0.98	0.900	(0.74	1.31)
Moderate high	0.84	0.254	(0.62	1.13)	0.84	0.264	(0.62	1.14)
High level	1.30	0.113	(0.94	1.80)	1.28	0.161	(0.91	1.82)
Tobacco and marijuana product use of relatives/friend living with participated students								
Smoking hubbly bubbly	4.31	0.000**	(2.71	(6.87)	4.14	0.000**	(2.59	6.62)
Smoke Marijuana	4.93	0.000**	(3.16	(7.70)	4.50	0.000**	(2.87	7.06)
Tobacco smoking	4.54	0.000**	(3.04	(6.78)	4.24	0.000**	(2.82	6.35)
Student residence								
On campus	0.86	0.322	(0.64	(1.16)	1.07	0.650	(0.79	1.45)
Off Campus	1.16	0.322	(0/86	(1.57)	0.93	0.650	(0.69	1.26)
Reasons for drinking alcohol								
I drink alcohol because it will enhance my creative ability	4.52	0.000*	(3.73	(8.21)	3.65	0.000**	(2.74	6.66)
I drink alcohol because I want to be popular	1.86	0.001**	(0.07	(3.71)	2.31	0.005*	(1.87	7.86)
I drink alcohol because it helps me to face difficulties with confidence	1.10	0.321	(0.31	(4.42)	0.24	0.371	(0.15	3.34)
I drink alcohol because it would ease me when I get blamed	3.41	0.561	(2.45	(4.73)	2.21	0.461	(1.57	4.63)
*≥0.05; **≥0.001.								

Table 3.
Odds ratio and 95% confidence interval and p-value for the association of current regular alcohol drinking use and the associated risk factors for University of Limpopo students aged 17 to 43 years.

significantly ($P < 0.05$) associated with the current regular alcohol use among University of Limpopo students for unadjusted (OR ranges 4.31 95%CI 2.71 6.87 to 4.92 95%CI 3.16 7.70) and after adjusted for age, gender, and amount of money deposited into the student account by the bursary scheme (OR ranges from 4.14 95%CI 2.59 6.62 to 4.50 95%CI 2.87 7.06). Place of residence and educational level of students were not significantly associated with current regular alcohol use among students. Current regular alcohol use among students was significantly ($P < 0.05$) associated with enhancing creativity (OR = 4.52 95%CI 3.73 8.210) and popularity (OR = 1.86 95%CI 0.07 3.71) motive among University of Limpopo students before adjusting the cofounders, even after adjusting cofounders, creativity (OR = 3.65 95%CI 2.74 6.66) popularity (OR = 2.31 95%CI 1.87 7.86)).

4. Discussions

This study aimed to investigate the prevalence of alcohol use and related risk factors among University of Limpopo students aged 17 to 43 years. The prevalence of current alcohol drinkers was significantly higher for males compared to females in a University of Limpopo. Similar patterns were reported earlier in South African Universities [7, 9, 10]. Among students in South Africa, Lategan *et al.*, [12] found significant differences in drinking behaviour between males and females, with male students drinking frequently more alcohol than female students. Kypri, *et al.*, [21] reported similar results among students in a university in New Zealand and further indicated that these students drank over the legal limit.

In the current study, a significant number of males compared to females ($P < 0.05$) preferred drinking beer and spirits. Weschler *et al.*, [22] reported that several students (both genders) chose spirits (43.8%), beer (19.2%) cider (17.4%) and very few chose wine (3.7%).

Students who preferred spirits and beer showed risky drinking behaviour such as driving while drunk and smoking [12]. Alcohol use at university tends to reach a peak among students aged between 18 and 25 years and are also at the risk of increasing their alcohol consumption later [23]. These university students may develop drinking patterns early in life and it will eventually affect their health and reduce their life expectancy later in their lifestyle.

In the current study alcohol use among students is associated with the use of tobacco and alcohol among friends and family they are staying with. Tareman *et al.*, [24] showed a relatively high incidence of alcohol (17%) abuse among University students in Tehran, however drugs abuse was also high with 34% water-pipe smoking, 24% cigarette, and 2.2% marijuana. Patterns of alcohol consumption and use of drugs vary depending on drinking motives. Cox and Klinger, [25] reported that alcohol drinking and drug use motives could be characterised into two underlying dimensions reflecting the valence (positive or negative) and source (internal or external) of the outcome an individual hopes to achieve. These drinking and drug problems of university students affect their academic achievement badly, emotional adaptableness, and the ability to get jobs after obtaining their career qualification [26].

In our current study alcohol use was not significantly associated with the students residents (off campus/on campus). Dawson *et al.*, [27] found the similar results were there was no differences in amount of alcohol drinking among students living on and off-campus. Similar results were found in the current study. Contrary, Simons-Morton *et al.*, [14] reported that independency and peer pressure increases the risk of alcohol consumptions among students living off-campus.

In the present study, most students drank alcohol to improve interpersonal relationships. The greatest motive for drinking was to be popular followed by difficulties with confidence among males in the current study. Yoo *et al.*, [28], reported a similar response among Korean Medical Students. The high prevalence of drinking among university students not only undermines academic performance, it also places student at risk of injuries, mortality, crime, and sexual assault [29]. It is necessary and essential to identify the significance of alcohol consumption complications in the institutional environment and system that can launch a healthy drinking culture. The problem of university student alcohol consumption can be solved by Policy-based tactics. A healthy drinking culture should be advocated by both the university authority, staff, students and parents such that skilled graduates of the University of Limpopo could pursue his/her career with dignity and vision.

This study has some limitations. We did not randomly select the students to participate in the study but have invited them to take part in the study. There is a possibility that our sample may not be generalizable to the University of Limpopo students. This study forms the base; ever since the higher education sector decided to deposit the bursary money into the student account every month to cover their essentials. Given the cross-sectional nature of the current study, a causal relationship could not be achieved. This was based on interviewing students by well-trained field workers hence recall bias should not be ruled out as an interview was conducted between lecture classes.

5. Conclusions

The prevalence of the current alcohol consumption is high among University of Limpopo male students compared to female students. Living with a person using Tobacco product, marijuana and alcohol are major associated risk factors for current alcohol use in the University of Limpopo students. Drinking problems should not be handled in isolation by the university authority in policy formulation for the culture of healthy drinking but also educate the students on the use of drugs and the motives for drinking. Future studies are required to investigate the development of lifestyle and cardiovascular disease over time among University of Limpopo students.

Acknowledgements

The authors are obligated to the participants (University of Limpopo students) for taking part in the study, resident manager for granting us permission to enter student resident to interview student. Ellirras longitudinal study team for helping with data collection.

Conflict of interest

Authors declare no conflict of interest.

IntechOpen

Author details

Mbelele Rosina Nkwana¹, Gift Makaleng¹, Mafoloa Suzan Monyeke¹,
Hlengani James Siweya² and Kotsedi Daniel Monyeke^{1*}

1 Department of Physiology and Environmental Health, University of Limpopo,
South Africa

2 Executive Dean Faculty of Science and Agriculture, University of Limpopo,
Sovenga, South Africa

*Address all correspondence to: kotsedi.monyeki@ul.ac.za

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Lapsley DK, Edgerton J. Separation-Individuation, Adult Attachment Style, and College Adjustment. *Counseling & Development J.* 2011; 80(4): 484-492.
- [2] Cooper ML. Motives for alcohol use among adolescent: Development and validation of a four factor model. *Psychological Assessment* 1994; 6 (2):117-128
- [3] Lorant V, Nicaise P, Soto VE, d'Hoore W. Alcohol drinking among college students: College responsibility for personal troubles. *BMC Public Health*, 2013; 13: 615
- [4] Lukács A, Simon N, Demeter JS, Kissné Dányi É, Kiss-Tóth E. Alcohol consumption among university students. *EGÉSZSÉGTUDOMÁNYI KÖZLEMÉNYEK: A MISKOLCI EGYETEM KÖZLEMÉNYE.* 2013;3(2):57-61.
- [5] Perkins HW. Surveying the damage: a review of research on consequences of alcohol misuse in college populations. *Journal of Studies on Alcohol, supplement.* 2002 Mar(14):91-100.
- [6] Schulenberg JE, Maggs JL. A developmental perspective on alcohol use and heavy drinking during adolescence and the transition to young adulthood. *Journal of Studies on Alcohol, Supplement.* 2002 Mar(14):54-70.
- [7] Young C, de Klerk V. Patterns of alcohol use on a South African university campus: The findings of two annual drinking surveys. *African Journal of Drug and Alcohol Studies.* 2008;7(2).
- [8] Chikere EI, Mayowa MO. Prevalence and perceived health effect of alcohol use among male undergraduate students in Owerri, South-East Nigeria: a descriptive cross-sectional study. *BMC Public health.* 2011 Dec 1;11(1):118.
- [9] Peltzer K, Ramlagan S. Alcohol use trends in South Africa. *Journal of Social Sciences.* 2009 Jan 1;18(1):1-2.
- [10] Kyei, K.A. and Ramagoma, M. 2013. Alcohol consumption in South African universities: Prevalence and factors at the University of Venda, Limpopo province. *Journal of Social Sciences*, 36: 77-86.
- [11] Arria AM, Caldeira KM, Allen HK, Vincent KB, Bugbee BA, O'Grady KE. Drinking like an adult? Trajectories of alcohol use patterns before and after college graduation. *Alcoholism: clinical and experimental research.* 2016 Mar;40(3):583-90.
- [12] Lategan BW, Du Preez R, Pentz CD. Socio-demographic insights into South African student drinking behaviour. *South African Journal of Higher Education.* 2017 Nov;31(5):90-115.
- [13] Harrell ZA, Karim NM. Is gender relevant only for problem alcohol behaviors? An examination of correlates of alcohol use among college students. *Addictive behaviors.* 2008 Feb 1;33(2):359-65.
- [14] Simons-Morton B, Haynie D, Liu D, Chaurasia A, Li K, Hingson R. The effect of residence, school status, work status, and social influence on the prevalence of alcohol use among emerging adults. *Journal of studies on alcohol and drugs.* 2016 Jan;77(1):121-32.
- [15] Matshipi M, Monyekei KD, Mafumo N, Monyekei SM, Siweya HJ, Kemper HC. The use of alcohol and knowledge of cardiovascular diseases among Ellirsas rural children aged 14-22 years: Ellirsas longitudinal study. *International journal of environmental*

research and public health. 2019 Jan;16(15):2650.

[16] Benz MB, DiBello AM, Balestrieri SG, Miller MB, Merrill JE, Lowery AD, Mastroleo NR, Carey KB. Off-campus residence as a risk factor for heavy drinking among college students. *Substance use & misuse*. 2017 Jul 29;52(9):1236-41.

[17] Bedendo A, Andrade AL, Opaleye ES, Noto AR. Binge drinking: a pattern associated with a risk of problems of alcohol use among university students. *Revista Latino-Americana de Enfermagem*. 2017;25.

[18] Suchy W, Gaczowska A, Pawelczyk A, Pukacki P, Chudzik R. Alcohol use among medicine and law students in Poland. *Journal of Medical Science*. 2015;84(3):183-8.

[19] Lauer RM, Lee JU, Clarke WR. Predicting adult cholesterol levels from measurements in childhood and adolescence: the Muscatine Study. *Bulletin of the New York Academy of Medicine*. 1989 Dec;65(10):1127.

[20] Altman RD. Criteria for classification of clinical osteoarthritis. *Journal of Rheumatology*. 1991 Jan 1;18(SUPPL. 27):10-2.

[21] Kypri K, Langley JD, McGee R, Saunders JB, Williams S. High prevalence, persistent hazardous drinking among New Zealand tertiary students. *Alcohol and Alcoholism*. 2002 Sep 1;37(5):457-64.

[22] Wechsler H, Nelson TF. What we have learned from the Harvard School of Public Health College Alcohol Study: Focusing attention on college student alcohol consumption and the environmental conditions that promote it. *Journal of Studies on alcohol and Drugs*. 2008 Jul;69(4):481-90.

[23] Nkambule NR, Bhayat A, Madiba TK. Knowledge, attitude and practices of alcohol and smoking among undergraduate oral health students at a South African University. *South African Dental Journal*. 2018 Feb;73(1):27-30.

[24] Taremian F, Bolhari J, Pairavi H, Ghazi Tabatabaei M. The prevalence of drug abuse among university students in Tehran. *Iranian J psychiatry and clinical psychology*. 2008; 13(4):335-42.

[25] Cox WM, Klinger E. Incentive motivation, affective change, and alcohol use: A model. Why people drink: Parameters of alcohol as a reinforcer. 1990:291-314.

[26] Zvaunya R, Oyeboode F, Day EJ, Thomas CP, Jones LA. A comparison of stress levels, coping styles and psychological morbidity between graduate-entry and traditional undergraduate medical students during the first 2 years at a UK medical school. *BMC research notes*. 2017 Dec 1;10(1):93.

[27] Dawson DA, Grant BF, Stinson FS, Chou PS. Another look at heavy episodic drinking and alcohol use disorders among college and noncollege youth. *Journal of studies on alcohol*. 2004 Jul;65(4):477-88.

[28] Yoo HH, Cha SW, Lee SY. Patterns of Alcohol Consumption and Drinking Motives Among Korean Medical Students. *Medical Science Monitor: International Medical Journal of Experimental and Clinical Research*. 2020;26:e921613-1.

[29] Hingson RW, Zha W, Weitzman ER. Magnitude of and trends in alcohol-related mortality and morbidity among US college students ages 18-24, 1998-2005. *Journal of Studies on Alcohol and Drugs, Supplement*. 2009 Jul(16):12-20.