We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Epidemiology, Genetics and Resistance of *Alternaria* Blight in Oilseed *Brassica*

Subroto Das Jyoti, Naima Sultana, Lutful Hassan
and Arif Hasan Khan Robin

Abstract

Alternaria blight is one of the most deadly diseases of oilseed *Brassica*. This recalcitrant disease causes up to 50% yield loss across the globe. The disease is mainly caused by *Alternaria brassicae* and *Alternaria brassicicola*. These pathogens lack sexual stages and survive as conidia or conidiospores on the debris of previous crops and susceptible weeds. Developing resistant oilseed *Brassica* cultivars to this disease has become a prime concern for researchers over the years. In absence of resistant oilseed *Brassica* cultivar, identification and introgression of resistance related genes can be a potential source for *Alternaria* blight resistance. As resistance toward *Alternaria* blight is governed by polygenes, intercrossing between the tolerant genotypes and subsequent selection will be the most appropriate way to transfer the quantitative resistance. For that reason, future breeding goal should focus on screening of germplasms for selecting genotypes containing resistance genes and structural features that favors resistance, like thick epicuticular wax, biochemical components such as phenols, phytoalexins and lower soluble sugars, reducing sugars and soluble nitrogen. Selected genotypes should be brought under appropriate breeding programs for attaining *Alternaria* blight resistance.

Keywords: *Alternaria* blight, oilseed *Brassica*, disease resistance, resistance mechanism

1. Introduction

Oilseed crops are one of the crucial pillars of world agriculture, occupying 22% of the world's arable land [1]. Rapeseed-mustard dominates the total oilseed production after soybean globally [1]. *Alternaria* blight is one of the major biotic threats that drastically reduces oilseed production all over the world including Australia, Europe, China and Canada [2]. *Alternaria* blight is a recalcitrant disease caused by the *Alternaria* species primarily *A. brassicae* and *A. brassicicola*, of which *A. brassicae* is the most deadly [3–4]. This disease decreases photosynthetic potential, leads to abnormal growth of the seeds and reduces seed oil content and quality [5]. Disease intensity varies across seasons and regions, and also between crops within an area [6]. Controlling the disease is one of the foremost concerns for researchers for reviving the yield potential of the rapeseed-mustard varieties. Chemical management of this disease is not proposed because maximum foliage

coverage by aerial application of fungicides is hard to achieve. Beside this, application of large amounts of chemicals raises environmental concerns. It is crucial to genetically monitor the disease by breeding for resistance [7]. Despite the immense efforts of breeders throughout the world, no resistant genotypes have been found till date. Combining various breeding tools may be fruitful in defining resistant genotypes in these scenarios. The genetic base of the cultivated oilseed *Brassica* is narrow and resistance governing genes are hard to find. *Alternaria* blight resistance is controlled by additives or polygenes and has been identified in some wild species of oilseed *Brassica* [8]. Easy availability of microarray data led researchers to the identification and understanding of the expression patterns of key genes involved in the *Alternaria* resistance. Another reliable form of plant immunity is Nonhost Resistance (NHR) that is successful against all genetic variants of a pathogen [8–11]. The infected plants also show hypersensitive response by producing reactive oxygen species [12]. Improvement of modern genetic transformation methods is helping scientists to incorporate resistant genes from non-host wild cultivars. Tissue culture method is one of the biotechnological tools that are being used to transfer resistance genes from resistant genotypes to the susceptible ones. Resistant genotypes showed higher phenolic content than the susceptible one, whereas the total soluble sugars, lower sugars and soluble nitrogen levels were lower [13–15]. Apart from all of these conventional methods, exploration and utilization of systematically acquired resistance and *de novo* resistance can be an efficient way to induce resistance in oilseed *Brassica* cultivar. Besides, molecular markers associated with resistance genes may contribute to the successful improvement of the resistance breeding process. This chapter discusses *Alternaria* blight disease with respect to its epidemiology, genetics and possible resistance mechanisms involved in *Alternaria* resistance and revisits earlier work done by oilseed *Brassica* breeders to elucidate future strategies for *Alternaria* resistance breeding.

2. Epidemiology

Disease epidemiology provides better understanding of the disease, host and favorable factors that facilitates disease progression. It also creates a better opportunity to control the disease by manipulating different epidemiological factors [16]. Majority of the *Alternaria* species produce asexual spores, as it lacks sexual stage (**Figure 1**; [17]). It survives as conidiospores or conidia under unfavorable conditions [18–19]. It also survives in the susceptible weed and in the infected seeds in temperate regions [20–23]. Although in tropical and subtropical India, the survival of *Alternaria* inoculum in seeds is discarded [24]. At first, symptoms start with black dots. Later, these spots extend and grow into prominent round spots with concentric circles displaying the spot's target board features (**Figure 2**). Many spots coalesce to form large patches which cause the leaves to blight and defoliate [4]. Initially the infection starts from the cotyledonary leaves and forms a basis for the secondary infection. Four hours of leaf wetness is necessary for leaf infection. An increase in leaf wetness duration at 25 °C increases infection and spread of the disease rapidly. Spores attack other parts of the plant upon getting favorable conditions. New lesions arise within four-five days. The pathogen infects the seed by penetrating the pod [25]. The critical factors for spore germination have been reported as darkness or low light intensity (<1000 lux), 25 °C temperature and more than 90% RH in some previous studies [26]. Some studies reported the increase of disease severity with the increase of inoculum concentration [27–29]. The optimal assay temperature of 25 °C and > 90% relative humidity resulted in the highest severity of the disease, regardless of the apparent susceptibility of the

Figure 1.
 Cultured spores (a) and conidia of *Alternaria brassicae* from the infected field samples (b).

Figure 2.
 Symptoms and different level of severity of *Alternaria* blight. Symptoms from 'a' to 'e' show gradually higher severity of infection.

cultigen [27, 30–36]. Previous studies reported that older leaves are more affected by *Alternaria* than the younger ones [27, 30, 37–40]. Weather characteristics such as maximum temperature 18–27 °C and minimum temperature 8–12 °C facilitates *Alternaria* infection on leaves with an average relative humidity more than 92% while on pods, the infection occurs at temperatures ranging from 20–30 °C [41]. Closer spacing (30 × 15 cm), high nitrogen doses (80 Kg Nha⁻¹) and frequent irrigation rapidly increase severity of disease in rapeseed–mustard [12]. Frequent rains are favorable for the initiation and spread of the disease on the leaves of oilseed *Brassica*. In addition, the rate of infection during the flowering and pod phases is the highest [42].

3. Genetics and genomics of *Alternaria* blight resistance

Identifying resistance mechanisms at the genetic and genomic level has been a prime concern for the researchers over the recent years. Various sources suggest that the resistance against *Alternaria* is polygenic [3, 43–45]. On the contrary, other studies reported that resistance to this disease is mainly controlled by only additive genes or dominant nuclear genes [3, 43–46]. However, Kumar et al. [47] proved that inheritance of *Alternaria* blight resistance is governed by more than one gene and fixable and non-fixable gene effects are vital in the genetic control of *Alternaria* blight resistance. In *Arabidopsis*, six QTLs governing *Alternaria* blight resistance were identified. Among these QTLs, five QTLs were population specific and one was common among all mapping populations. Presence of both common and population specific QTLs indicates that resistance against *Alternaria* blight is quantitative and more than one gene potentially governs the resistance [48].

With the modern development of biotechnology, the discovery of resistance (R) and defense-related genes has opened up new scopes for inducing genetic resistance against different biotic and abiotic stresses [49]. Advances in microarray data processing also ease the process of identifying candidate genes in certain physiological processes. In previous studies, *A. brassicicola* infection contributed to the upregulation of different genes such as *WRKY*, *peroxidase*, *p450 oxidases*, *Chitinase* that modulates defense response in oilseed *Brassica* and *Arabidopsis*. A recent computational study identified vital genes involved in *Alternaria* resistance in *Brassica* by analyzing microarray data of model plant *Arabidopsis thaliana* challenged with *Alternaria* infection [50]. *NHL10*, *HCHIB* and *XLG2* were identified as major genes and *CZF1*, *ARF6*, *WRKY*, *MP*, *IAA1*, *IAA19*, *AXR3* as candidate genes associated in defense response against *Alternaria* [50]. PR (pathogenesis-related) proteins are a distinct group of molecules which are induced by phytopathogens and signaling molecules linked to defense. They are the vital components of the plant's inherent immune system, particularly systemic acquired resistance (SAR) [51]. Two genes under these proteins namely *Chitinase* and *NPR1* have been characterized in oilseed *Brassica* species. Their high expression level in resistant genotypes compared to the susceptible genotypes suggested that these genes are related to resistance against *Alternaria* blight [52–53]. Another study reported the expression of *PR-3* and *PR-12* only in *Camelina sativa* and *Sinapis alba* compared with *B. juncea* [54]. This clarifies the involvement of PR proteins in the resistance mechanism of *Alternaria* resistant varieties.

4. Biochemical resistance against *Alternaria*

Biochemical defense is triggered by any stress condition in a plant and is the most important tool of plant defense mechanism. The hypersensitive response is one of the plant's most effective defensive responses against the pathogen [55]. Resistance to *Alternaria* blight in mustard was reported to be linked with the synthesis of phenolic pathway-associated leaf enzymes and higher leaf sugar content [56]. The concentration of phenolic compounds at all stages of plant growth was reported to be high in resistant genotypes compared to susceptible genotypes. Nevertheless, soluble sugars, sugar reduction and soluble nitrogen levels in resistant genotypes were lower [14–15]. Another study reported that, total phenol, total sugar, reducing sugar, o-dihydroxy phenol, chlorophyll content and flavonol contents were higher in resistant genotypes [57]. By activating several defense responses that dissuade the infection process, plants can respond to a pathogen. These include the production of reactive oxygen species (ROS), the accumulation of proteins related to pathogenesis (PR) and phytoalexins and the synthesis of compounds that strengthen the plant cell wall [58]. Moreover the contents of ascorbic acid, total phenol, enzymatic activities of superoxide dismutase and peroxidase, that of cell protecting enzymes such as phenylalanine ammonia lyase and polyphenol oxidases were increased in the resistant genotypes of mustard [59]. β -Aminobutyric acid (BABA), a non-protein amino acid has been known to stimulate resistance to a variety of pathogens in a number of plant species [60–61]. Pretreatment of oilseed *Brassica* plants with BABA-mediated resistance to the necrotrophic pathogen *A. brassicae* through enhanced expression of protein genes linked to pathogenesis [62]. The colonization of *A. brassicae* on *Brassica carinata* leaves was substantially inhibited by the foliar application of BABA [63]. A higher and early accumulation of H_2O_2 was observed in resistant *C. sativa* and *S. alba* compared to *B. juncea*. Catalase activity was enhanced in both *C. sativa* and *S. alba*, but the opposite phenomenon was observed in case of *B. juncea* [54].

5. Utilization of non-host resistance

Non-host resistance is one of the most useful approaches for attaining resistance against different plant pathogens. Till date, no resistant cultivar is available in oilseed *Brassica* species. Therefore, utilizing the non-host resistance from wild species can be an efficient breeding tool. Plant pathogens manage to affect different species, but they fail to overcome the non-host resistance [64]. Examples of some non-host plants of *A. brassicae* are chickpea, lentil, wheat, sugarcane, barley, tomato, potato [64]. NHR is multilayered and can be splitted into two main forms: the layer of pre-invasion and the phase of post-invasion [65–67]. Preformed defenses may include structural features like abundance of trichomes and spore germination inhibitory chemical compounds [68–70]. Previous studies reported that spore germination occurs at an equal rate in both host and non-host plants [71]. Despite an accurate germination, pathogens might fail to reach the stomata. Stomata in non-host plants may not be correctly recognized by the pathogen because the topography of the surface may vary significantly from that of the host leaf [64]. Another structural feature that can prevent the entry of *Alternaria* is the epicuticular wax [72–74]. Non-host plants may have higher epicuticular wax than the susceptible host plants [64]. The non-host plant is capable of inducing stomatal closure, preventing pathogens from entering and constructing an inducible chemical barrier that suppresses hyphal production and differentiation by the rapid formation of phytoalexins, antimicrobial compounds [75–77]. In a non-host plant, the dietary deficiency and the presence of antimicrobial compounds in the apoplast can also prevent the production of hyphae into mycelium [71]. The pathogen also generates non-host specific or general toxins that might damage plant cells, leading ultimately to necrosis [78–80]. To avoid this, a non-host plant may recognize these toxins and employ defense mechanisms to detoxify these toxins [81]. In *Arabidopsis* and *S. alba* pathogenesis-related genes *PR-1*, *PR-2*, *PR-3* were highly expressed compared to *B. juncea* after *Alternaria* infection [82–86]. Furthermore, these two species showed non-host resistance toward *A. brassicicola* [81, 87]. Chitinase enzymes that hydrolyze the fungal cell wall and release fragments of chitin are actively secreted by these two species [82, 88]. The NHR action includes the stimulation by the plant cell of a signal transduction cascade following the detection of a pathogen, which triggers the activation of protein kinases and mitogen-activated protein kinase (MAPK) members and consequently lead to the activation of defensive genes in non-host plants [89]. The expression of *MAPK* was higher in *S. alba* and downregulated in *B. juncea* suggesting its possible role in *Alternaria* blight resistance.

6. Genetic transformation for *Alternaria* resistance

As the resistance of *Alternaria* has not yet been found, identification of resistance genes in non-host plants and transferring them into oilseed *Brassica* species could be a handy tool for resistance breeding. Introgression of genes under PR-proteins have been found effective in many cases. For instance, transgenic Indian mustard was developed with the *chitinase* gene in which the occurrence of disease symptoms was delayed by a duration of 10–15 days compared to control plants [90]. For enhancing resistance against *A. brassicae*, a PR protein-encoding glucanase was introduced from tomato into Indian mustard plants [91]. Glucanase hydrolyzes a main component of a fungal cell wall called glucan and destroys the invading fungal pathogens. In combating *Alternaria* blight disease, a barley anti-fungal class II *chitinase* gene and type I ribosome inactivating protein (*RIP*) gene were co-expressed in Indian mustard [92]. Transgenic mustard plants demonstrated

a 44% reduction in *A. brassicae* hyphal production relative to the control plants. When transgenic events were sprinkled with fungal spores through greenhouse screening, the late onset of the disease and a lower number of lesions with reduced size distribution were recorded. In addition, *Chitinase* gene was transferred from *Streptomyces griseus* HUT6037 to Indian mustard [93]. A previous study transformed *B. juncea* with the *osmotin* gene and documented resistance to the purified *A. brassicae* toxin in the transformed calli [94]. *B. juncea* was modified to add resistance to *Alternaria* blight and stem rot diseases with the *MSRA1* gene [95]. Bioassays after *Alternaria* infection *in vitro* showed that transgenic *B. juncea* lines inhibited the growth of *Alternaria* hyphae by 44–62% and reduced infection ranging from 69–85%. The *lectin* gene of chickpea was transferred to Indian mustard cv. Varuna to induce resistance against *A. brassicae* in transgenic lines [96]. Another study incorporated *B. juncea* with the gene *MPK3* and examined its role in providing tolerance against *A. brassicae* [97]. In transgenic plants, both ascorbate peroxidase (APX) and guaiacol peroxidase (GP) activity and proline content were higher, leading to the scavenging of ROS in transgenic plants developed as a result of infection with *Alternaria*.

When an *endochitinase* gene '*echh42*' from the *Trichoderma virens*, a fungal species used as a bio-control agent, was introduced to *B. juncea*— the transformed plants showed 7-fold higher endochitinase activity compared to the non-transformed plants based on fluorimetric analysis [98]. These results indicated that the *endochitinase* gene '*ech42*' could be a major gene that may provide resistance to oilseed *Brassica* plants against the *Alternaria* blight. In previous studies, the transgenic broccoli plants also showed expression of *chitinase* gene of *Trichoderma harzianum* [99–101]. Moreover, the synthetic *chitinase* gene (NIC) showed broad-spectrum resistance to the transgenic lines of *B. juncea* including *A. brassicicola* [102]. Further research utilizing RT-PCR validated that these *chitinase* genes were induced after wounding and exogenous treatments of jasmonic acid and salicylic acid similar to *Alternaria* infection [103]. A recent review summarized that the chitinases, glucanases or cry proteins provide broad-spectrum resistance against some major diseases including *Alternaria* blight and blackleg [104].

7. *De novo* resistance

It is assumed that the disease can be successfully managed by inducing protection inducers in plants. Some novel fungicides may mimic the action of different plant hormones that activate the plant's internal immune response. Jasmonic acid (JA) mediated defense response to *A. brassicae* fungus can prevent necrotrophic colonization mode. The JA receptor, coronatine insensitive 1 (COI1), is one of the possible targets to activate JA-mediated immunity via JA signal interaction [105]. It is understood that Jasmonates and its functional analogs play a crucial role in systemic defense, likely serving as the initiating signal of acquired systemic resistance [106]. It has been shown that necrotrophic fungal pathogens are the primary activators of JA-dependent defenses via COI₁ receptor activation [107]. A previous study identified some JA mimicking molecules that might be helpful in *de novo* resistance induction [108].

8. Tissue culture techniques in *Alternaria* resistance

Tissue culture is one of the most effective tools of modern biotechnology. Somaclonal variation provides an opportunity to extend the genetic variation of

crops, i.e. the variation caused by cell and tissue culture. By applying *in vitro* selection process, the efficiency of selection can be increased [109]. Somatic hybrids were produced through PEG-mediated symmetric and asymmetric protoplast fusion, in which *S. alba*, *B. nigra* and *B. juncea* were found to be the most effective resistance donor to *Alternaria* pathogen [110]. Through protoplast fusion, a previous study developed three hybrids between *B. juncea* and *S. alba* [111]. Among the hybrids, two of the hybrids were symmetric, while the third was asymmetric and had greater similarity to *B. juncea*. *Alternaria* resistant lines were developed through interspecific hybridization between *S. alba* and *B. juncea* [112]. *Alternaria* blight resistance was transferred from *B. tourneforti* to *B. juncea* cv. RH 30 through *in vitro* ovule culture [113]. Intergeneric hybrids of *B. campestris* and *B. spinenscens* were generated through sequential ovary, ovule and embryo culture [114]. The resistance trait was transferred to *B. napus* cv. Brutor from *S. alba* cv. Carine following *in vitro* fertilized ovary culture protocol [115]. *Erucastrum cardaminoides* and *B. oleracea* var. alboglabra were used to develop intergeneric hybrids with *Alternaria* blight resistance following sequential ovary and ovule culture procedures [116]. Previous studies reported transfer of *Alternaria* resistance through somatic hybridization such as, from *S. alba* to *B. napus* [117] and *Moricandida arvensis* to *B. oleracea* [118]. A research group in India transferred *Alternaria* resistance trait to *B. juncea* from *B. carinata* [119]. Disease resistant hybrid plants were produced from the hybridized leaf mesophyll protoplasts of *M. arvensis* and *B. napus* [120]. *B. carinata* was resynthesized by protoplast fusion between *B. nigra* and *B. oleracea* [121]. The hybrids thus obtained were fertile and grew into robust plants. Previous studies conducted hybridization between *S. alba* and *B. oleracea* and between *Camelina sativa* and *B. oleracea* for producing resistant hybrids [122–123]. Another study developed somatic hybrids between *S. alba* and *B. oleracea* by protoplast fusion followed by embryo rescue and managed to recover four highly resistant hybrid progenies after repeated backcrosses [124]. By inducing variations through gamma-irradiated mutagenesis the resistant varieties were obtained in *B. juncea* [125] while another study achieved the similar results by treating the embryos with chemical mutagens [126]. It is plausible to say that proper utilization of tissue culture techniques can be a successful means of incorporating *Alternaria* resistance into oilseed Brassica cultivars.

9. Molecular markers and *Alternaria* blight resistance

In any disease resistance breeding program, the primary approach is to quickly screen all the available germplasm including local races, improved variety and exotic genetic stocks. The traditional approach of screening of genotypes can be costly, time and space consuming, laborious, and involves large sample sizes [127]. The limitations of conventional approach can be solved through molecular markers. By utilizing molecular markers, economically important major genes and quantitative trait loci (QTLs) can be identified [128]. Pre-selection using molecular markers can minimize the size of a population and facilitate early detection of desirable genotypes [127]. Various molecular markers are being used nowadays for assessing genetic variability against *Alternaria* blight. For example, internal transcribed spacer regions (ITS), restriction fragment length polymorphism (RFLP), randomly amplified polymorphic DNA (RAPD), amplified fragment length polymorphism (AFLP), inter-simple sequence repeats (ISSRs), microsatellites (SSR), sequence tagged sites (STS), single nucleotide polymorphism (SNPs) etc. The ITS regions are the preserved areas in the fungal genome that are considered as the most common loci to study DNA based mycology at the species level. Berbee and co-workers

studied the ITS regions of rDNA to determine the pathogen's phylogeny [129]. RAPD technique was used successfully to examine the genetic differences in *Alternaria* infected species [130–132]. Later on, the assessment of genetic variability in *Alternaria* species has moved to more sensitive techniques such as AFLP [133] and microsatellite markers [134] due to the constraints of reproducibility of RAPD. Simple sequence repeats have been isolated and characterized from *B. napus*, *B. nigra*, and *B. rapa* [135, 136]. Moreover, SSR marker libraries have been developed for *B. rapa* those are being used to produce a genome map for *B. rapa* [137]. Recently, SNP markers have taken the supremacy over SSR as they are unique and plentiful in high and ultra-high-throughput and are able to find polymorphism within a single base pair [138].

10. Conclusions

Alternaria blight is one of the major diseases of oilseed *Brassica* causing enormous yield loss every year. In order to reduce the use of chemical fertilizers and to save the environment, breeding is important to attain resistance against *Alternaria* pathogens. Since the resistance against *Alternaria* blight is governed by additive or polygenes, molecular breeding for resistance could be more effective. All possible sources including wild relatives and non-host plants should be brought under the selection process for identifying ideal resistance donors. QTL mapping and continuous hybridization between resistant genotypes should be performed for better results. Emphasis should be given on functional analysis of PR proteins for engineering *Alternaria* resistance more effectively. In addition, accurate modeling of plant's internal defense responsive pathways can provide new insights on *de novo* and systematically acquired resistance.

Funding

This research is supported by the Ministry of Education, Bangladesh (Project title: Development of oilseed *Brassica* genotypes resistant to *Alternaria* blight through accelerated genetic gain; Project No. 2019/05/MoE).

Conflict of interest

The authors declare no conflict of interest.

IntechOpen

IntechOpen

Author details

Subroto Das Jyoti, Naima Sultana, Lutful Hassan and Arif Hasan Khan Robin*
Department of Genetics and Plant Breeding, Bangladesh Agricultural University,
Mymensingh, Bangladesh

*Address all correspondence to: gpb21bau@bau.edu.bd

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] FAO. World Food and Agriculture - Statistical Yearbook; 2020. Rome. DOI: 10.4060/cb1329en
- [2] Zheng X, Koopmann B, Ulber B, von Tiedemann A. A Global Survey on Diseases and Pests in Oilseed Rape—Current Challenges and Innovative Strategies of Control. *Frontiers in Agronomy*. 2020; 2:1-5. DOI: 10.3389/fagro.2020.590908
- [3] Meena PD, Awasthi RP, Chattopadhyay C, Kolte SJ, Kumar A. *Alternaria* blight: a chronic disease in rapeseed-mustard. *Journal of Oilseed Brassica*. 2016; 1:1-1.
- [4] Kumar D, Maurya N, Bharati YK, Kumar A, Kumar K, Srivastava K, Chand G, Kushwaha C, Singh SK, Mishra RK, Kumar A. *Alternaria* blight of oilseed *Brassicas*: A comprehensive review. *African Journal of Microbiology Research*. 2014; 8:2816-29. DOI: 10.5897/AJMR2013.6434
- [5] Saharan GS, Mehta DN, Meena PD, Dayal P. *Alternaria* diseases of crucifers: biology, ecology and disease management. Singapore: Springer; 2016.
- [6] Meena PD, Rani A, Meena R, Sharma P, Gupta R, Chowdappa P. Aggressiveness, diversity and distribution of *Alternaria brassicae* isolates infecting oilseed *Brassica* in India. *African Journal of Microbiology Research*. 2012; 6:5249-58. DOI: 10.5897/AJMR12.705
- [7] Sharma G, Kumar VD, Haque A, Bhat SR, Prakash S, Chopra VL. *Brassica* coenospecies: a rich reservoir for genetic resistance to leaf spot caused by *Alternaria brassicae*. *Euphytica*. 2002; 125:411-7. DOI: 10.1023/A:1016050631673
- [8] Heath MC. Nonhost resistance and nonspecific plant defenses. *Current Opinion in Plant Biology*. 2000; 3:315-9. DOI: 10.1016/S1369-5266(00)00087-X
- [9] Mysore KS, Ryu CM. Nonhost resistance: how much do we know? *Trends in Plant Science*. 2004; 9:97-104. DOI: 10.1016/j.tplants.2003.12.005
- [10] Senthil-Kumar M, Mysore KS. Nonhost resistance against bacterial pathogens: retrospectives and prospects. *Annual Review of Phytopathology*. 2013; 51:407-27. DOI: 10.1146/annurev-phyto-082712-102319.
- [11] Lee HA, Lee HY, Seo E, Lee J, Kim SB, Oh S, Choi E, Choi E, Lee SE, Choi D. Current understandings of plant nonhost resistance. *Molecular Plant-Microbe Interactions*. 2017; 30:515. DOI 10.1094/MPMI-10-16-0213-CR.
- [12] Lamb C, Dixon RA. The oxidative burst in plant disease resistance. *Annual Review of Plant Biology*. 1997; 48:251-75. DOI: 10.1146/annurev.arplant.48.1.251
- [13] Verma PR, Saharan GS. Monograph of *Alternaria* Diseases of Crucifers. Agriculture and Agri-Food Canada, Saskatoon Research Centre. Technical Bulletin E. 1994; 6:150.
- [14] Meena PD, Bala M, Sharma P, Singh D. Interaction and tolerance to *Alternaria brassicae* in Indian mustard (*Brassica juncea*) genotypes. *The Indian Society of Oilseeds Research*. 2014:136.
- [15] Meena PD, Kumar MS, Meena HS, Jambhulkar S, Pathak D, Srivastava S, Gupta R, Singh D, Gurung B, Rai PK. Confirmation of induced tolerance to *Alternaria* blight disease in Indian mustard (*Brassica juncea* L.). *Applied Biochemistry and Biotechnology*. 2020; 192:965-78. DOI: 10.1007/s12010-020-03362-2
- [16] Jones DG. An introduction to plant disease epidemiology. In *The*

- Epidemiology of Plant Diseases. Springer, Dordrecht. 1998; 3-13. DOI: 10.1007/978-94-017-3302-1_1
- [17] Mamgain A, Roychowdhury R, Tah J. *Alternaria* pathogenicity and its strategic controls. Research Journal of Biology. 2013; 1:1-9.
- [18] Humpherson-Jones FM, Maude RB. Studies on the epidemiology of *Alternaria brassicicola* in *Brassica oleracea* seed production crops. Annals of Applied Biology. 1982; 100:61-71. DOI: 10.1111/j.1744-7348.1982.tb07192.x
- [19] Humpherson-Jones FM. Survival of *Alternaria brassicae* and *Alternaria brassicicola* on crop debris of oilseed rape and cabbage. Annals of Applied Biology. 1989; 115:45-50. DOI: 10.1111/j.1744-7348.1989.tb06810.x
- [20] Sherf AF, MacNab AA. Vegetable diseases and their control. John Wiley & Sons. New York. 693: 267-269.
- [21] Maude RB, Humpherson-Jones FM. Studies on the seed-borne phases of dark leaf spot *Alternaria brassicicola* and grey leaf spot *Alternaria brassicae* of *Brassicas*. Annals of Applied Biology. 1980; 95:311-9. DOI: 10.1111/j.1744-7348.1980.tb04752.x
- [22] Shrestha SK, Munk L, Mathur SB. Survival of *Alternaria brassicae* in seeds and crop debris of rapeseed and mustard in Nepal. Seed Science and Technology. 2003; 31:103-9. DOI: 10.15258/sst.2003.31.1.11
- [23] Ahmad MS, Sinha KK. Mycotoxin contamination of mustard seeds during storage. Indian Phytopathology. 2002; 55:299-302.
- [24] Mehta N, Sangwan MS, Srivastava MP, Kumar R. Survival of *Alternaria brassicae* causing *Alternaria* blight rapeseed-mustard. Journal of Mycology and Plant Pathology. 2002; 32: 64-67.
- [25] Saharan GS, Mehta N, Sangwan MS, editors. Diseases of oilseed crops. Indus Publishing. 2005.
- [26] Mehta N. Epidemiology and forecasting for the management of rapeseed-mustard diseases. Journal of Mycology and Plant Pathology. 2014; 44:131.
- [27] Hong CX, Fitt BD. Effects of inoculum concentration, leaf age and wetness period on the development of dark leaf and pod spot (*Alternaria brassicae*) on oilseed rape (*Brassica napus*). Annals of Applied Biology. 1995; 127:283-95. DOI: 10.1111/j.1744-7348.1995.tb06673.x
- [28] Meena PD, Jambhulkar SJ, Gupta R, Meena HS, Singh D. Rapid screening technique for *Alternaria* blight resistance in Indian mustard (*Brassica juncea* L.) using cotyledonary leaf method. Journal of Plant Pathology. 2016: 463-9.
- [29] Nowakowska M, Wrzesińska M, Kamiński P, Szczechura W, Lichocka M, Tartanus M, Kozik EU, Nowicki M. *Alternaria brassicicola*–Brassicaceae pathosystem: insights into the infection process and resistance mechanisms under optimized artificial bio-assay. European Journal of Plant Pathology. 2019; 153:131-51. DOI: 10.1007/s10658-018-1548-y
- [30] Doullah MA, Meah MB, Okazaki K. Development of an effective screening method for partial resistance to *Alternaria brassicicola* (dark leaf spot) in *Brassica rapa*. European Journal of Plant Pathology. 2006; 116:33-43. DOI: 10.1007/s10658-006-9035-2
- [31] Gupta P, Ravi I, Sharma V. Induction of β -1, 3-glucanase and chitinase activity in the defense response of *Eruca sativa* plants against the fungal pathogen *Alternaria brassicicola*. Journal of Plant Interactions. 2013; 8:155-61. DOI: 10.1080/17429145.2012.679705

- [32] Mazumder M, Das S, Saha U, Chatterjee M, Bannerjee K, Basu D. Salicylic acid-mediated establishment of the compatibility between *Alternaria brassicicola* and *Brassica juncea* is mitigated by abscisic acid in *Sinapis alba*. Plant Physiology and Biochemistry. 2013; 70:43-51. DOI: 10.1016/j.plaphy.2013.04.025
- [33] Rashid MM, Hossain I, K. M. Khalequzzaman KM. Effect of weather factors on inoculum density and leaf spot development in radish seed crop infected with *Alternaria brassicae*. Bulletin of the Institute of Tropical Agriculture, Kyushu University. 2011; 34:43-7. DOI: 10.11189/bit.34.43
- [34] Sharma G, Kumar VD, Haque A, Bhat SR, Prakash S, Chopra VL. *Brassica* coenospecies: a rich reservoir for genetic resistance to leaf spot caused by *Alternaria brassicae*. Euphytica. 2002; 125:411-7.
- [35] Su'udi M, Kim MG, Park SR, Hwang DJ, Bae SC, Ahn IP. *Arabidopsis* cell death in compatible and incompatible interactions with *Alternaria brassicicola*. Molecules and Cells. 2011; 31:593-601.
- [36] Zală CR, Cristea ST, Gruia LI, Manole SA. Research on the biology of the *Alternaria brassicae* fungus isolated from mustard. In 3rd International Conference on Energy and Environment Technologies and Equipment. Brasov, Romania. p 2014 (pp. 161-164).
- [37] Allen SJ, Brown JF, Kochman JK. Effects of leaf age, host growth stage, leaf injury, and pollen on the infection of sunflower by *Alternaria helianthi*. Phytopathology. 1983; 73:896-8.
- [38] Deep SW, Sharma PR. Host age as predisposing factor for incidence of black leaf spot of cauliflower caused by *Alternaria brassicae* and *Alternaria brassicicola*. Indian Phytopathology. 2012; 65:71-5.
- [39] Rotem J. The biotic and physiological components of pathogenesis. The Genus *Alternaria*; Biology, Epidemiology, and Pathogenicity, The American Phytopathological Society, St. Paul, MN. 1998:95-118.
- [40] Saharan GS, Mehta N. Fungal diseases of rapeseed-mustard. Diseases of Field Crops. 2002:193-228.
- [41] Aneja JK, Agnihotri A. *Alternaria* blight of oilseed *Brassicas*: epidemiology and disease control strategies with special reference to use of biotechnological approaches for attaining host resistance. Journal of Oilseed Brassica. 2016; 1:1-10.
- [42] Prasad RA, Saxena DE, Chandra S. Yield losses by *Alternaria* blight in promising genotypes of Indian mustard. Indian Phytopathology. 2003; 56(2):205-6.
- [43] Tripathi NN, Kaushik CD, Yadava TP, Yadav AK. *Alternaria* leaf spot resistance in raya. Haryana Agricultural University Journal of Research. 1980; 10:166-8.
- [44] Zhang FL, Xu JB, Takahata Y. Inheritance of resistance to black leaf spot (*Alternaria brassicae*) in Chinese cabbage. Cruciferae Newsletter (United Kingdom). 1996.
- [45] Krishnia SK, Saharan GS, Singh D. Genetics of *Alternaria* blight resistance in inter and intraspecific crosses of *Brassica juncea* and *Brassica carinata*. Annals of Biology. 2000; 16:211-6.
- [46] Panja BN, De DK: Combining ability and heterosis for resistant to leaf blight disease in Indian mustard [*Brassica juncea* (L.) Czern & Coss]. Indian Journal of Genetics. 2005; 65: 55-6.
- [47] Kumar V, Kumari V, Meena PD, Kumar A. Role of inter genic

interactions in inheritance of *Alternaria* leaf blight resistance in inter and intraspecific crosses of *Brassica carinata* and *Brassica juncea*. *Journal of Oilseed Brassica*. 2020; 11:126-8.

[48] Rajarammohan S, Kumar A, Gupta V, Pental D, Pradhan AK, Kaur J. Genetic architecture of resistance to *Alternaria brassicae* in *Arabidopsis thaliana*: QTL mapping reveals two major resistance-conferring loci. *Frontiers in Plant Science*. 2017; 8:260. DOI: 10.3389/fpls.2017.00260

[49] Pandolfi V, Ribamar Costa Ferreira Neto J, Daniel da Silva M, Lindinalva Barbosa Amorim L, Carolina Wanderley-Nogueira A, Lane de Oliveira Silva R, Akio Kido E, Crovella S, Maria Benko Iseppon A. Resistance (R) genes: applications and prospects for plant biotechnology and breeding. *Current Protein and Peptide Science*. 2017; 18:323-34.

[50] Pathak RK, Baunthiyal M, Pandey D, Kumar A. Computational analysis of microarray data of *Arabidopsis thaliana* challenged with *Alternaria brassicicola* for identification of key genes in *Brassica*. *Journal of Genetic Engineering and Biotechnology*. 2020; 18:1-20. DOI: 10.1186/s43141-020-00032-y

[51] Ali S, Ganai BA, Kamili AN, Bhat AA, Mir ZA, Bhat JA, Tyagi A, Islam ST, Mushtaq M, Yadav P, Rawat S. Pathogenesis-related proteins and peptides as promising tools for engineering plants with multiple stress tolerance. *Microbiological Research*. 2018; 212:29-37. DOI: 10.1016/j.micres.2018.04.008

[52] Ali S, Mir ZA, Tyagi A, Mehari H, Meena RP, Bhat JA, Yadav P, Papalou P, Rawat S, Grover A. Overexpression of *NPR1* in *Brassica juncea* confers broad spectrum resistance to fungal pathogens. *Frontiers in Plant Science*. 2017; 8:1693. DOI: 10.3389/fpls.2017.01693

[53] Mir ZA, Ali S, Shivaraj SM, Bhat JA, Singh A, Yadav P, Rawat S, Paplao PK, Grover A. Genome-wide identification and characterization of *Chitinase* gene family in *Brassica juncea* and *Camelina sativa* in response to *Alternaria brassicae*. *Genomics*. 2020; 112:749-63. DOI: 10.1016/j.ygeno.2019.05.011

[54] Yadav P, Mir ZA, Ali S, Papolu PK, Grover A. A combined transcriptional, biochemical and histopathological study unravels the complexity of *Alternaria* resistance and susceptibility in *Brassica* coenospecies. *Fungal Biology*. 2020; 124:44-53.

[55] Goodman RN, Novacky AJ. The hypersensitive reaction in plants to pathogens: a resistance phenomenon. *American Phytopathological Society (APS)*; 1994.

[56] Singh DN, Singh NK, Srivastava S. Biochemical and morphological characters in relation to *Alternaria* blight resistance in rapeseed-mustard. *Annual Agriculture Research*. 1999; 20:472-7.

[57] Mathpal P, Punetha H, Tewari AK, Agrawal S. Biochemical defense mechanism in rapeseed-mustard genotypes against *Alternaria* blight disease. *Journal of Oilseed Brassica*. 2016; 1:87-94.

[58] Hammond-Kosack KE, Jones JD. Resistance gene-dependent plant defense responses. *The Plant Cell*. 1996; 8:1773. DOI: 10.1105/tpc.8.10.1773

[59] Mallick SA, Kumari P, Gupta M, Gupta S. Effect of *Alternaria* blight infection on biochemical parameters, quantity and quality of oil of mustard genotypes. *Indian Journal of Plant Physiology*. 2015; 20:310-316.

[60] Jakab G, Cottier V, Toquin V, Rigoli G, Zimmerli L, Métraux JP, Mauch-Mani B. β -Aminobutyric acid-induced resistance in plants. *European*

Journal of Plant Pathology. 2001; 107:29-37. DOI: 10.1023/A:1008730721037

[61] Cohen YR. β -aminobutyric acid-induced resistance against plant pathogens. Plant Disease. 2002; 86:448-57. DOI: 10.1094/PDIS.2002.86.5.448

[62] Kamble A, Bhargava S. β -Aminobutyric Acid-induced resistance in *Brassica juncea* against the necrotrophic pathogen *Alternaria brassicae*. Journal of Phytopathology. 2007; 155:152-8. DOI: 10.1111/j.1439-0434.2007.01209.x

[63] Chavan V, Bhargava S, Kamble A. Temporal modulation of oxidant and antioxidative responses in *Brassica carinata* during β -aminobutyric acid-induced resistance against *Alternaria brassicae*. Physiological and Molecular Plant Pathology. 2013; 83:35-9. DOI: 10.1016/j.pmpp.2013.03.002

[64] Fatima U, Bhorali P, Borah S, Senthil-Kumar M. Perspectives on the utilization of resistance mechanisms from host and nonhost plants for durable protection of *Brassica* crops against *Alternaria* blight. PeerJ. 2019; 7:e7486. DOI: 10.7717/peerj.7486

[65] Nurnberger T, Lipka V. Non-host resistance in plants: new insights in an old phenomenon. Molecular Plant Pathology. 2005; 6:335-345. DOI 10.1111/j.1364-3703.2005.00279.x.

[66] Lee HA, Lee HY, Seo E, Lee J, Kim SB, Oh S, Choi E, Choi E, Lee SE, Choi D. Current understandings of plant nonhost resistance. Molecular Plant-Microbe Interactions. 2017; 30:5-15. DOI 10.1094/MPMI-10-16-0213-CR.

[67] Fonseca JP, Mysore KS. Genes involved in nonhost disease resistance as a key to engineer durable resistance in crops. Plant Science. 2018; 279:108-116.

[68] Heath MC. Evolution of plant resistance and susceptibility to fungal

parasites. In: Carroll GC, Tudzynski P, editors. The Mycota V, part B, plant relationships. Berlin: Springer. 1997; 257-276.

[69] Heath MC. Signalling between pathogenic rust fungi and resistant or susceptible host plants. Annals of Botany. 1997; 80:713-720. DOI 10.1006/anbo.1997.0507.

[70] Morrissey JP, Osbourn AE. Fungal resistance to plant antibiotics as a mechanism of pathogenesis. Microbiology and Molecular Biology Reviews. 1999; 63:708-724. DOI:10.1128/MMBR.63.3.708-724.1999

[71] McRoberts N, Lennard JH. Pathogen behaviour and plant cell reactions in interactions between *Alternaria* species and leaves of host and nonhost plants. Plant Pathology. 1996; 45:742-752. DOI: 10.1046/j.1365-3059.1996.d01-4.x

[72] Tewari JP, Skoropad WP. Relationship between epicuticular wax and blackspot caused by *Alternaria brassicae* in three lines of rapeseed. Canadian Journal of Plant Science. 1976; 56:781-785. DOI 10.4141/cjps76-127.

[73] Skoropad WP, Tewari JP. Field evaluation of the epicuticular wax in rapeseed and mustard in resistance to *Alternaria brassicae*. Canadian Journal of Plant Science. 1997; 57:1001-1003. DOI 10.4141/cjps77-146.

[74] Conn KL. Leaf epicuticular wax of canola: ultrastructure, chemistry and interaction with *Alternaria brassicae*. M. S.C. thesis, University of Alberta, Edmonton.1986; 159.

[75] Grayer RJ, Harborne JJ. A survey of antifungal compounds from higher plants. Phytochemistry. 1994; 37:19-42. DOI 10.1016/0031-9422(94)85005-4.

[76] Mert-Turk F. Phytoalexins: defence or just a response to stress? Journal of Cell and Molecular Biology. 2002; 1:1-6.

- [77] Iriti M, Faoro F. Review on innate and specific immunity in plants and animals. *Mycopathology*. 2007; 164:57-64. DOI 10.1007/s11046-007-9026-7.
- [78] Buchwaldt L, Green H. Phytotoxicity of destruxin B and its possible role in the pathogenesis of *Alternaria brassicae*. *Plant Pathology*. 1992; 41:55-63. DOI 10.1111/j.1365-3059.1992.tb02316.x.
- [79] Bains PS, Tewari JP, Ayer WA. A note on phytotoxicity of homodestruxin B—a compound produced by *Alternaria brassicae*. *Phytoprotection*. 1993; 74:157-160.
- [80] Parada RY, Oka K, Yamagishi D, Kodama M, Otani H. Destruxin B produced by *Alternaria brassicae* does not induce accessibility of host plants to fungal invasion. *Physiology and Molecular Plant Pathology*. 2007; 71:48-54. DOI 10.1016/j.pmpp.2007.10.003.
- [81] Pedras MSC, Zaharia IL, Gai Y, Zhou Y, Ward DE. In planta sequential hydroxylation and glycosylation of a fungal phytotoxin: avoiding cell death and overcoming the fungal invader. *Proceedings of the National Academy of Sciences of the United States of America*. 2001; 98:747-752. DOI 10.1073/pnas.98.2.747.
- [82] Narusaka Y, Narusaka M, Seki M, Ishida J, Nakashima M, Kamiya A, Enju A, Sakurai T, Satoh M, Kobayashi M. The cDNA microarray analysis using an *Arabidopsis* pad3 mutant reveals the expression profiles and classification of genes induced by *Alternaria brassicicola* attack. *Plant Cell and Physiology*. 2003; 44:377-387. DOI 10.1093/pcp/pcg050.
- [83] Van Wees SC, Chang HS, Zhu T, Glazebrook J. Characterization of the early response of *Arabidopsis* to *Alternaria brassicicola* infection using expression profiling. *Plant Physiology*. 2003; 132:606-617. DOI 10.1104/pp.103.022186.
- [84] Ghose K, Dey S, Barton H, Loake GJ, Basu D. Differential profiling of selected defence-related genes induced on challenge with *Alternaria brassicicola* in resistant white mustard and their comparative expression pattern in susceptible India mustard. *Molecular Plant Pathology*. 2008; 9:763-775. DOI 10.1111/j.1364-3703.2008.00497.x.
- [85] Nayanakantha NMC, Rawat S, Ali S, Grover A. Differential expression of defense-related genes in *Sinapis alba* and *Brassica juncea* upon the infection of *Alternaria brassicae*. *Tropical and Agricultural Research*. 2016; 27:123-136.
- [86] Mandal S, Rajarammohan S, Kaur J. *Alternaria brassicae* interactions with the model Brassicaceae member *Arabidopsis thaliana* closely resembles those with Mustard (*Brassica juncea*). *Physiology and Molecular Biology of Plants*. 2018; 24: 51-59. DOI: 10.1007/s12298-017-0486-z
- [87] Thomma BP, Eggermont K, Tierens KF, Broekaert WF. Requirement of functional ethylene-insensitive 2 gene for efficient resistance of *Arabidopsis* to infection by *Botrytis cinerea*. *Plant Physiology*. 1999; 121:1093-1101. DOI 10.1104/pp.121.4.1093.
- [88] Chatterjee M, Mazumder M, Basu D. Functional analysis of the promoter of a glycosyl hydrolase gene induced in resistant *Sinapis alba* by *Alternaria brassicicola*. *Phytopathology*. 2013; 103:841-850. DOI 10.1094/PHTO-11-12-0303-R.
- [89] Lawrence CB, Mitchell TK, Craven KD, Cho YR, Cramer RA, Kim KH. At death's door: *Alternaria* pathogenicity mechanisms. *Plant Pathology Journal*. 2008; 24:101-111. DOI: 10.5423/PPJ.2008.24.2.101.

- [90] Mondal KK, Chatterjee SC, Viswakarma N, Bhattacharya RC, Grover A. Chitinase-mediated inhibitory activity of *Brassica* transgenic on growth of *Alternaria brassicae*. *Current Microbiology*. 2003; 47:0171-3. DOI: 10.1007/s00284-002-3980-6
- [91] Mondal KK, Bhattacharya RC, Koundal KR, Chatterjee SC. Transgenic Indian mustard (*Brassica juncea*) expressing tomato glucanase leads to arrested growth of *Alternaria brassicae*. *Plant Cell Reports*. 2007; 26:247-52. DOI: 10.1007/s00299-006-0241-3
- [92] Chhikara S, Chaudhury D, Dhankher OP, Jaiwal PK. Combined expression of a barley class II chitinase and type I ribosome inactivating protein in transgenic *Brassica juncea* provides protection against *Alternaria brassicae*. *Plant Cell, Tissue and Organ Culture (PCTOC)*. 2012; 108:83-9. DOI: 10.1007/s11240-011-0015-7
- [93] Ahmad B, Ambreen Ms, Khan I. *Agrobacterium*-mediated transformation of *Brassica juncea* (L.) Czern. with *chitinase* gene conferring resistance against fungal infections. *Pakistan Journal of Botany*. 2015; 47:211-6.
- [94] Taj G, Kumar A, Bansal KC, Garg GK. Introgression of *osmotin* gene for creation of resistance against *Alternaria* blight by perturbation of cell cycle machinery. *Indian Journal of Biotechnology*. 2004, 3: 291-298.
- [95] Rustagi A, Kumar D, Shekhar S, Yusuf MA, Misra S, Sarin NB. Transgenic *Brassica juncea* plants expressing *MsrA1*, a synthetic cationic antimicrobial peptide, exhibit resistance to fungal phytopathogens. *Molecular Biotechnology*. 2014; 56:535-45. DOI: 10.1007/s12033-013-9727-8
- [96] Kumar D, Shekhar S, Bisht S, Kumar V, Varma A, Kumar M. Ectopic overexpression of lectin in transgenic *Brassica juncea* plants exhibit resistance to fungal phytopathogen and showed alleviation to salt and drought stress. *Journal of Bioengineering and Biomedical Science*. 2015; 5:147. DOI: 10.4172/2155-9538-1000147
- [97] Tasleem M, Baunthiyal M, Kumar A, Taj G. Determination of antioxidant activity in overexpressed *MPK3* transgenic *Brassica juncea* for induction of defense against *Alternaria* blight disease. *Journal of Pharmacognosy and Phytochemistry*. 2017; 6:2579-82.
- [98] Kamble S, Mukherjee PK, Eapen S. Expression of an endochitinase gene from *Trichoderma virens* confers enhanced tolerance to *Alternaria* blight in transgenic *Brassica juncea* (L.) Czern and Coss lines. *Physiology and Molecular Biology of Plants*. 2016; 22: 69-76.
- [99] Mora A, Earle E. Combination of *Trichoderma harzianum* endochitinase and a membrane-affecting fungicide on control of *Alternaria* leaf spot in transgenic broccoli plants. *Applied Microbiology and Biotechnology*. 2001; 55: 306-310.
- [100] Mora AA, Earle ED. Resistance to *Alternaria brassicicola* in transgenic broccoli expressing a *Trichoderma harzianum* endochitinase gene. *Molecular Breeding*. 2001; 8:1-9.
- [101] Singh A, Shukla N, Kabadwal BC, Tewari AK, Kumar J. Review on plant-*Trichoderma*-pathogen interaction. *International Journal of Current Microbiology and Applied Sciences*. 2018; 7: 2382-2397.
- [102] Munir I, Hussan W, Kazi M, Farhatullah A, Iqbal A, Munir R. Production of transgenic *Brassica juncea* with the synthetic chitinase gene (NIC) Conferring resistance to *Alternaria brassicicola*. *Pakistan Journal of Botany*. 2016; 48: 2063-2070.
- [103] Rawat S, Ali S, Mittra B, Grover A. Expression analysis of chitinase upon

challenge inoculation to *Alternaria* wounding and defense inducers in *Brassica juncea*. *Biotechnology Reports*. 2017; 13: 72-79.

[104] Poveda J, Francisco M, Cartea ME, Velasco P. Development of Transgenic *Brassica* Crops against Biotic Stresses Caused by Pathogens and Arthropod Pests. *Plants*. 2020; 9: 1664.

[105] Pathak RK, Baunthiyal M, Pandey N, Pandey D, Kumar A. Modeling of the jasmonate signaling pathway in *Arabidopsis thaliana* with respect to pathophysiology of *Alternaria* blight in *Brassica*. *Scientific Reports*. 2017; 7:1-2.

[106] Truman W, Bennett MH, Kubigsteltig I, Turnbull C, Grant M. *Arabidopsis* systemic immunity uses conserved defense signaling pathways and is mediated by jasmonates. *Proceedings of the national academy of sciences*. 2007; 104:1075-80. DOI: 10.1073/pnas.0605423104

[107] Antico CJ, Colon C, Banks T, Ramonell KM. Insights into the role of jasmonic acid-mediated defenses against necrotrophic and biotrophic fungal pathogens. *Frontiers in Biology*. 2012; 7:48-56. DOI: 10.1007%2Fs11515-011-1171-1

[108] Pathak RK, Baunthiyal M, Shukla R, Pandey D, Taj G, Kumar A. *In Silico* Identification of mimicking molecules as defense inducers triggering jasmonic acid mediated immunity against *Alternaria* blight disease in *Brassica* species. *Frontiers in Plant Science*. 2017; 8:609. DOI: 10.3389/fpls.2017.00609

[109] Van den Bulk RW. Application of cell and tissue culture and in vitro selection for disease resistance breeding—a review. *Euphytica*. 1991; 56:269-85. DOI: 10.1007/BF00042373

[110] Scholze P, Krämer R, Ryschka U, Klocke E, Schumann G. Somatic hybrids

of vegetable *Brassicas* as source for new resistances to fungal and virus diseases. *Euphytica*. 2010; 176:1-4. DOI: 10.1007/s10681-010-0205-0

[111] Kumari P, Bisht DS, Bhat SR. Stable, fertile somatic hybrids between *Sinapis alba* and *Brassica juncea* show resistance to *Alternaria brassicae* and heat stress. *Plant Cell, Tissue and Organ Culture (PCTOC)*. 2018; 133:77-86. DOI: 10.1007/s11240-017-1362-9

[112] Yadav RC, Singh D, Rath AR, Arya R, Singh R, Yadav NR, Singh A. Development of *Alternaria* blight resistant lines through interspecific hybridization between Indian mustard (*Brassica juncea* L.) and white mustard (*Brassica alba*) through embryo rescue. *Journal of Oilseed Brassica*. 2018:104-13.

[113] Yadav RC, Sareen PK, Chowdhury JB. Interspecific hybridization in *Brassica juncea* x *Brassica tournefortii* using ovary culture. *Cruciferae Newsletter*. 1991;84:14-5.

[114] Agnihotri A, Lakshmikumaran MS, Jagannathan V, Shivanna KR. Wide hybridization for improvement in cultivated *Brassicas*. In: *International Symposium on Plant Biotechnology and its Contribution to Plant Development, Multiplication and Improvement* 289. 1991 (pp. 213-214).

[115] Chèvre AM, Eber F, Margale E, Kerlan MC, Primard C, Vedel F, Delseny M, Pelletier G. Comparison of somatic and sexual *Brassica napus*–*Sinapis alba* hybrids and their progeny by cytogenetic studies and molecular characterization. *Genome*. 1994;37: 367-74. DOI: 10.1139/g94-052

[116] Mohanty A, Chrungu B, Verma N, Shivanna KR. Broadening the genetic base of crop *Brassicas* by production of new intergeneric hybrid. *Czech Journal of Genetics and Plant Breeding*. 2009; 45: 117-22. DOI: 10.17221/35/2009-CJGPB

- [117] Primard C, Vedel F, Mathieu C, Pelletier G, Chèvre AM. Interspecific somatic hybridization between *Brassica napus* and *Brassica hirta* (*Sinapis alba* L.). Theoretical and Applied Genetics. 1988; 75:546-52. DOI: 10.1007%2FBBF00289119
- [118] Toriyama K, Hinata K, Kameya T. Production of somatic hybrid plants, 'Brassicomoricaandia', through protoplast fusion between *Moricandia arvensis* and *Brassica oleracea*. Plant Science. 1987; 48:123-8. DOI: 10.1016/0168-9452(87)90139-7
- [119] Sharma TR, Singh BM. Transfer of resistance to *Alternaria brassicae* in *Brassica juncea* through interspecific hybridization among *Brassicaceae*. Journal of Genetics & Breeding. 1992; 46:373-8.
- [120] O'Neill CM, Murata T, Morgan CL, Mathias RJ. Expression of the C 3-C 4 intermediate character in somatic hybrids between *Brassica napus* and the C 3-C 4 species *Moricandia arvensis*. Theoretical and Applied Genetics. 1996; 93:1234-41. DOI: 10.1007%2FBBF00223455
- [121] Jourdan P, Salazar E. *Brassica carinata* resynthesized by protoplast fusion. Theoretical and Applied Genetics. 1993; 86:567-72. DOI: 10.1007%2FBBF00838710
- [122] Hansen LN, Earle ED. Somatic hybrids between *Brassica oleracea* L. and *Sinapis alba* L. with resistance to *Alternaria brassicae* (Berk.) Sacc. Theoretical and Applied Genetics. 1997; 94:1078-85. DOI: 10.1007%2Fs001220050518
- [123] Hansen LN. Intertribal somatic hybridization between rapid cycling *Brassica oleracea* L. and *Camelina sativa* (L.) Crantz. Euphytica. 1998; 104:173-9. DOI: 10.1023/A: 1018604919488
- [124] Sigareva MA, Earle ED. Production and characterization of somatic hybrids between *Camelina sativa* and rapid-cycling *B. oleracea*. Theoretical and Applied Genetics. 1999; 98:164-70. DOI: 10.1007/s001220051053
- [125] Verma, VD, Rai, B. Note on induced mutagenesis for spotting out the sources of resistance to *Alternaria* leaf spot in Indian mustard. Indian Journal of Agricultural Sciences. 1980; 50: 278.
- [126] Agnihotri A, Gupta K, Prem D, Sarkar G, Mehra VS, Zargar SM. Genetic enhancement in rapeseed-mustard for quality and disease resistance through in vitro techniques. Proceedings of 16th Australian research assembly on *Brassicaceae*, Ballarat, Australia. 2009; 28.
- [127] Ghosh S, Mazumder M, Mondal B, Mukherjee A, De A, Bose R, Das S, Bhattacharyya S, Basu D. Morphological and SSR marker-based genetic diversity analysis of Indian mustard (*Brassica juncea* L.) differing in *Alternaria brassicicola* tolerance. Euphytica. 2019; 215: 206. DOI: 10.1007/s10681-019-2523-1.
- [128] Suwabe K, Iketani H, Nunome T, Kage T, Hirai M. Isolation and characterization of microsatellites in *Brassica rapa* L. Theoretical and Applied Genetics. 2002; 104:1092-8. DOI: 10.1007/s00122-002-0875-7.
- [129] Berbee ML, Payne BP, Zhang G, Roberts RG, Turgeon BG. Shared ITS DNA substitutions in isolates of opposite mating type reveal a recombining history for three presumed asexual species in the filamentous ascomycete genus *Alternaria*. Mycological research. 2003; 107: 169-82. DOI: 10.1017/S0953756203007263.
- [130] Cooke DE, Forster JW, Jenkins PD, Jones DG, Lewis DM. Analysis of intraspecific and interspecific variation in the genus *Alternaria* by the use of RAPD-PCR. Annals of Applied Biology.

1998; 132:197-209. DOI: 10.1111/j.1744-7348.1998.tb05197.x

[131] Sharma TR, Tewari JP. Detection of genetic variation in *Alternaria brassicae* by RAPD fingerprints. *Journal of Plant Biochemistry and Biotechnology*. 1995; 4: 105-7. DOI: 10.1007/BF03262963

[132] Gherbawy YA. Genetic variation among isolates of *Alternaria* spp. from select Egyptian crops. *Archives of Phytopathology and Plant Protection*. 2005; 38: 77-89. DOI: 10.1080/03235400400027390

[133] Bock CH, Thrall PH, Brubaker CL, Burdon JJ. Detection of genetic variation in *Alternaria brassicicola* using AFLP fingerprinting. *Mycological Research*. 2002; 106: 428-34. DOI: 10.1017/S0953756202005762

[134] Avenot H, Dongo A, Bataillé-Simoneau N, Iacomini-Vasilescu B, Hamon B, Peltier D, Simoneau P. Isolation of 12 polymorphic microsatellite loci in the phytopathogenic fungus *Alternaria brassicicola*. *Molecular Ecology Notes*. 2005; 5: 948-50. DOI: 10.1111/j.1471-8286.2005.01125.x

[135] Lowe AJ, Jones AE, Raybould AF, Trick M, Moule CL, Edwards KJ. Transferability and genome specificity of a new set of microsatellite primers among *Brassica* species of the U triangle. *Molecular Ecology Notes*. 2002; 2:7-11. DOI: 10.1046/j.1471-8286.2002.00126.x

[136] Piquemal J, Cinquin E, Couton F, Rondeau C, Seignoret E, Perret D, Villeger MJ, Vincourt P, Blanchard P. Construction of an oilseed rape (*Brassica napus* L.) genetic map with SSR markers. *Theoretical and Applied Genetics*. 2005;111: 1514-23. DOI: 10.1007/s00122-005-0080-6

[137] Ramchiary N, Nguyen VD, Li XI, Hong CP, Dhandapani VI, Choi SR, Yu GE, Piao ZY, Lim YP. Genic

microsatellite markers in *Brassica rapa*: development, characterization, mapping, and their utility in other cultivated and wild *Brassica* relatives. *DNA research*. 2011; 18: 305-20. DOI: 10.1093/dnares/dsr017

[138] Mammadov J, Aggarwal R, Buyyarapu R, Kumpatla S. SNP markers and their impact on plant breeding. *International Journal of Plant Genomics*. 2012; 2012.