We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Bioinoculants in Technological Alleviation of Climatic Stress in Plants

Rafia Younas, Shiza Gul, Rehan Ahmad, Ali Raza Khan, Mumtaz Khan, Tauseef Anwar and Huma Qureshi

Abstract

Global climate change is leading to a series of frequent onset of environmental stresses such as prolonged drought periods, dynamic precipitation patterns, heat stress, and cold stress on plants and commercial crops. The increasing severity of such stresses is not only making agriculture and related economic sector vulnerable but also negatively influences plant diversity patterns. The global temperature of planet Earth has risen to 1.1°C since the last 19th century. An increase in surface temperature leads to an increase in soil temperature which ultimately reduces water content in the soil, thereby, reducing crop growth and yield. Moreover, this situation is becoming more intense for agricultural practices in arid and semi-arid regions. To overcome climatically induced stresses, acclimatization of plant species via bioinoculation with Plant Growth Promoting Rhizobacteria (PGPR) is becoming an effective approach. The PGPR are capable of colonizing rhizosphere (exophytes) as well as plant organs (endophytes), where they trigger an accumulation of osmolytes for osmoregulation or improving gene expression of heat or cold stress proteins, or by signaling the synthesis of phytohormones, metabolites, proteins, and antioxidants to scavenge reactive oxygen species. Thus, PGPR exhibiting multiple plant growth-promoting traits can be employed via bioinoculants to improve the plant's tolerance against unfavorable stress conditions.

Keywords: Plant growth promoting rhizobacteria, plant growth-regulating hormones, ACC deaminase, osmoregulation, psychrotrophic microbes

1. Introduction

The agriculture sector has been considered as the main channel through which irreparable climatic change will influence the global economy [1]. The commercial plantation of crops has been severely influenced by various environmental factors mainly being drought, salinity, and intense temperature variation. These factors have become more adverse with continuing fluctuation in climate change. Globally, only 10% of arable land is reported to be free from environmental stress [2]. In arable land, more than 50% is vulnerable to drought and salinization which is expected to inflict an almost 10% rise in global water consumption by 2050. Moreover, the consistent rise in the global population will be succeeded by an increase in food demand double fold in 2050 [3].

Plants adapt and develop their own mechanism to cope the environmental stress. The most effective means of combating environmental stress is the release and accumulation of plant growth-regulating hormones such as abscisic acid (ABA) and salicylic acid (SA). ABA, commonly known as stress hormone is released at the root tip in response to lower water potential, followed by its translocation to leaves where it alters osmotic pressure of the stomatal guard cells, leading to the closure of stomata openings to conserve moisture, thereby, preventing water loss via transpiration. Moreover, accumulated carbohydrates and amino acids being low molecular weight compounds play an active role in osmoregulation. SA, on the other hand, protects the plant against pathogenic attack [4].

PGPR/Treated Plant	Climatic stress	Alleviation of climatic stress
<i>Bacillus cereu/Solanum lycopersicum</i>	Heat stress	Increased production of exo-polysaccharide, cleavage of ACC-deaminase [9]
<i>Bacillus amyloliquefaciens, Agrobacterium fabrum/Triticum aestivum</i>	Water stress	Increase in biomass and grain yield [10]
<i>Aneurinibacillus aneurinilyticus, Paenibacillus sp./Phaseolus vulgaris</i>	Salinity stress	Facilitating ACC-deaminase activity, increased production of IAA, hydrogen cyanide and siderophore [11]
<i>Stenotrophomonas maltophilia/Triticum aestivum</i>	Salinity stress	Improved growth and yield, elevated antioxidative enzymatic activity, increased K ⁺ uptake [12]
<i>Paenibacillus polymyxa/Brassica napus</i>	Soil temperature	Production of antimicrobial and plant growth promoting volatile organic compounds (VOCs) [13]
<i>Bacillus subtilis, Bacillus thuringiensis, Bacillus megaterium/Cicer arietinum</i>	Climatically induced salinity stress	Regularized photosynthesis via assimilation of soluble sugars, chlorophyll and proline [14]

Table 1.
Alleviation of climatic stress via bioinoculants.

Figure 1.
Plant’s responses to various stressses in presence and absence of bioinoculants.

Along with plant internal defense mechanism, the supportive role of PGPR has been widely acknowledged as an effective tool for sustaining plant growth and yield [5–7]. The plant root system serves as a habitat for numerous microorganisms that also interact with soil, thereby developing a complex ecosystem [8]. Certain PGPR inhabiting the rhizosphere penetrates roots and migrates through plant tissues; thereby impact the physiological and biochemical traits of plant cells (**Table 1**). This ecological aspect of the microbial community can be employed to strengthen the stress tolerance in plants, hence increasing their adaptation towards invincible climatic changes (**Figure 1**) [15].

The escalating demand and supply of chemical fertilizers and pesticides along with the persistence of their residual particles has posed an ultimate threat to humans as well as to the ecosystem. As an alternative, the acceptability of bio-fertilizers bearing bioinoculants has been increasing day by day. These bioinoculants act as a potential source of phytohormones which can rectify the climatic stress faced by plants. This chapter will explore various aspects of the utilization of bioinoculants for combating the climatic stress faced by plants.

2. Amelioration of heat stress via thermotolerant microbes

The major climatic stress faced by plants due to the rise in global temperature is heat stress which leads to water stress with counter effect on photosynthetic rate and flowering and fruiting in both tropical and sub-tropical crop systems [16]. The steady increase in surface temperatures is followed by an increase in soil temperature which can influence the root elongation process. Although, root elongation requires optimum soil temperature (specific to each plant), beyond which root elongation is stopped resulting in a stunted root system. Moreover, the activity of rhizobacteria is also relying on optimum soil temperature. In the forest ecosystem, N being a macronutrient acts as a growth-limiting factor. The flow of N cycling is deliberately subjected to temperature variation. The relatively increased temperature triggers N mineralization by microbes and thereby, prompting N uptake by plants as nitrates. However, in this process, the net nitrification is decreased [17]. The inoculation of PGPR is found to be more convenient than hsfs (heat stress transcription factors) genes, transgenic varieties, and breeding of heat-tolerant cultivators [18, 19].

Thermotolerance of certain PGPR has been evaluated to decrease the effect of heat stress on plants. The soybean plants inoculated with *Bacillus cereus* strain SA1 were exposed to heat stress for 5–10 days. The inoculated plants exhibited improved physiological (biomass) and biochemical (chlorophyll content and chlorophyll fluorescence) characteristics. Moreover, SA1 inoculation reduced the synthesis of abscisic acid and increased salicylic acid (a phenolic phytohormone involved in signaling and defense against pathogens) along with remarkable production of antioxidants (ascorbic acid peroxidase, superoxide dismutase, and glutathione) for sequestration of reactive oxygen species (ROS). A continuous increase in the synthesis and assimilation of Heat Shock Protein (HSP) was also claimed throughout the stress period of 5 to 10 days, mainly due to the perpetual onset of GmHSP gene expression. Also, over gene expression of GmLAX3 and GmAKT2 were associated with enhanced potassium gradient and altered auxin and ABA stimuli (Aaqil et al. 2020). The bacterial strain *Bacillus cereus* capable of producing ACC-deaminase (0.76–0.9 $\mu\text{M}/\text{mg}$ protein/h) can cleavage ACC to α -ketobutyrate and ammonia to increase heat tolerance in tomato (*Solanum lycopersicum* L.). Moreover, increase in synthesis of exopolysaccharide (0.66–0.91 mg/mL) showed promising plant growth in tomato [9] (**Table 2**). Moreover, the presence of enzymatic antioxidants

Treated Plant/PGPR	Stress Alleviation Effect
Potato/ <i>Paraburkholderia phytofirmans</i>	ACC deaminase production [20]
Tomato/Mycorrhizae	Reduction in lipid peroxidation and H ₂ O ₂ , higher ROS scavenging activity [21]
<i>Bacillus amyloliquefaciens</i> , <i>Azospirillum brasilense</i> <i>Pantoea dispersa</i> , <i>Serratia marcescens</i> , <i>Pseudomonas</i> spp.	ROS reduction, pre-activation of heat shock proteins [22]
	ACC deaminase production [23]
Apple and pear/ <i>Ps. fluorescens</i>	Competition with INA+ bacteria [24]
Grapevine/ <i>Paraburkholderia phytofirmans</i>	ACC deaminase production [25]
Soy/ <i>Bacillus aryabhatthai</i>	ABA production [26]

Table 2.
Alleviation of heat stress via bioinoculants.

such as superoxide dismutase (SOD), peroxidase (POD) and catalase (CAT), and accumulation of metabolites such as amino acids and proline had boosted the scavenging activity against ROS under heat stress.

Among different agro-ecological zones, the arid and semi-arid regions are highly vulnerable to climatic changes mainly due to dynamical precipitation patterns. Being a staple cereal crop, the wheat and its associated endophytic and epiphytic bacterial diversity was investigated for thermotolerance at a temperature range of 30–40°C, pH 3–11, and 3–20% NaCl concentration [27]. Among 32 bacterial strains of 15 genera, 10 strains were reported to exhibit six different plant growth-promoting traits under heat stress. *Arthrobacter* sp. ($66.0 \pm 0.7 \text{ mg L}^{-1}$) possessed high solubilization of phosphorus followed by *Pseudomonas japonica* ($64.6 \pm 0.9 \text{ mg L}^{-1}$). Whereas, the potassium solubilization was highest for *Methylobacterium mesophilicum*. *Pseudomonas putida* exhibited the highest IAA production ($70.8 \pm 1.5 \text{ } \mu\text{g mg}^{-1} \text{ protein day}^{-1}$) followed by *Rhodobacter capsulatus* ($69.1 \pm 0.5 \text{ } \mu\text{g mg}^{-1} \text{ protein day}^{-1}$). The highest production of siderophore had been shown by *Alcaligenes faecalis* ($4.9 \pm 0.1 \text{ mm}$). Being thermotolerant, *Alcaligenes faecalis* and *Pseudomonas poae* were also characterized as alkali tolerant with 5% NaCl and 10% NaCl tolerance respectively. Besides the production of IAA, siderophore, and ammonia, these bacterial strains were also involved in the solubilization of phosphorus and zinc under heat stress. The chief bacterial groups characterized for nitrogen fixation at high temperatures were *Acromobacter*, *Alcaligenes*, *Bacillus*, *Delftia*, *Providencia*, *Pseudomonas*, *Rhodobacter*, and *Salmonella*.

Similarly, Sorghum being native to arid habitat was inoculated with a thermo-tolerant bacterial strain of *Psuedomonas aeruginosa* AKM-P6 which was isolated from the rhizosphere of Pigeon Pea plant grown under semi-arid conditions [28]. The inoculated Sorghum seedlings survived the temperature range of 47–50°C up to 15 days. The thermotolerance was imparted by the biosynthesis of high molecular weight proteins in leaves which protected cellular membranes from injury and improved metabolite production mainly proline, chlorophyll, sugars, amino acids, and proteins along with plant biomass. Another strain of *Pseudomonas* sp. PsJN was used to evaluate the heat-stress tolerance level in 18 clones of potatoes via both *in vitro* and *ex-vitro* inoculation [29]. The inoculated potato nodal cuttings were exposed to the temperature range of 20/15°C or 33/15°C day and night for six weeks. The increase in temperature had drastically affected the root system and tuber number and tuber fresh weight. An average root to shoot ratio decreased from 3.7 at 20/15°C to 1.7 at 33/15°C in non-inoculated plantlets and respectively, from 4.3

to 1.5 for inoculated ones. ABA deficient plantlets lack the tuber formation whereas tuberization in inoculated plantlets was significant at 33/15°C. As compared to in vitro, the ex-vitro performance of potato clones (LT-7) exhibited the effectiveness of rhizobacteria in colonizing certain potato clones under heat stress. Hence, the thermotolerant bacterial strains may serve as inoculants and bio-control agents for improving crop productivity with accelerating climatic temperature stress.

3. Amelioration of water stress via 1-aminocyclopropane-1-carboxylate deaminase/exopolysaccharide production

Due to the changes in minimum and maximum temperature, high water requirement and reduction in yield was observed in plants. Although the production of 'Ethylene' as plant stress hormone occurs in response to climatic stress encountered by plants; its accumulation in high amounts could be deleterious. The accumulation occurs under the influence of ACC oxidase that prompts exudation of ACC, a precursor to ethylene synthesis. The PGPR capable of producing ACC deaminase degrades ACC into its intermediates i.e., ammonia and α -ketobutyrate, thereby, declining ethylene level and restoring plant development [30]. The inoculation of *Mucuna pruriens* L. (velvet bean), a well-adapted plant to arid and semi-arid regions with *Bacillus* spp. and *Enterobacter* spp. had shown significantly reduced ACC accumulation by 41% and 21% in leaves respectively, and in the roots by 46% and 15%, respectively. Thereby, the ethylene synthesis was reduced by 45% with *Bacillus* spp. (G9) and 65% with *Enterobacter* spp. (HS9) [29].

In addition to ACC deaminase activity, certain bacterial strains were found to be efficient in plant growth. For instance, *Achromobacter piechaudii* ARV8, a bacterial strain isolated from the arid region, experiencing frequent episodes of water stress was found to be better plant growth promoter as compared to *Pseudomonas putida* GR12-2, a bacterial strain inhabiting area with surplus water supply [31]. The bioinoculation of tomato (*Lycopersicum esculentum* Mill) and pepper (*Capsicum annum* L.) seedlings with *A. piechaudii* strain improved relative water content (RWC) which subsequently succored plants in maintaining their fresh weight in water stress conditions. The maintenance of the fresh weight in bioinoculated seedling can be justified by the declined production of ethylene to 6.1 nlh^{-1} as compared to stressed seedling with the rise in ethylene level up to 23 nlh^{-1} . Overall, the plant biomass was enhanced four times as compared to uninoculated controls. Similarly bacterial strains mainly *Variovorax paradoxus*, *Pseudomonas* spp., *Achromobacter* spp. and *Ochrobactrum anthropi* isolated from rain-fed agricultural soil were reported to possess growth-promoting traits such as N_2 fixation, siderophore, and phosphate solubilization along with sufficient production of ACC deaminase. These bacterial strains both as single inoculation and consortium imparted significant growth, foliar concentration, and antioxidant activity to the wheat (*Triticum aestivum* L.) production [32].

The exopolysaccharide (EPS) released by PGPR tends to form rhizosheath or biofilms around the surface of the roots containing sufficient moisture content, thereby, protect them from prolonged desiccation. Moreover, plants inoculated with EPS are capable of accumulating sugars, amino acids, and proline. The survival of PGPR in dried sandy soils with low moisture content is conceivable due to EPS, which may serve as a biological tool to combat climatic water stress and ultimately providing a path towards global food security [33]. The coalition of ACC deaminase and EPS has been shown to have an affirmative effect on carotenoid pigments in certain plants. The inoculation of *Capsicum annum* plants with *Bulkholderia cepacia* (ACC deaminase activity 12.8 ± 0.44 , $\text{mM } \alpha\text{KB mg}^{-1} \text{ min}^{-1}$ and

EPS 4.89 ± 0.06 mg/mg protein) resulted in increased chlorophyll a and b content (chlorophyll a 5.7 gm L^{-1} and chlorophyll b 3.4 gm L^{-1} , respectively) as compared to control (chlorophyll a 3.2 gm L^{-1} , chlorophyll b 1.9 gm L^{-1}). Moreover, the plant biomass has also increased (fresh weight of 9 g and dry weight of 3.6 g) as compared to control (fresh weight 6 g, and dry weight 1.6 g) [34].

The co-application of PGPR producing ACC deminase with biochar has been utilized as an effective mechanism against prevalent water stress conditions. Timber-waste biochar was co-applied with *Bacillus amyloliquefaciens* strain to increase the productivity of wheat under simulated water stress conditions. Along with high productivity (59% of 100-grain weight), the grain composition (58% N, 18% P, and 23% K), carotenoid (114% chlorophyll a and 123% chlorophyll b), photosynthetic rate (118%), and transpiration rate (73%) in wheat were also reinforced [10]. To understand the PGPR potential for genetic improvement in plants, the physiology and biochemical characteristics of PGPR isolated from chickpea rhizosphere were implored. The seeds of chickpea were grown in the cultures of *Bacillus subtilis*, *Bacillus thuringiensis*, and *Bacillus megaterium*. The 25-day old chickpea seedlings were sprayed with plant growth regulators/plant growth retardants (PGRs) i.e., salicylic acid (SA) and putrescine (Put) at the rate of 150 mgL^{-1} . The consortium of PGPR and PGRs significantly improved carotenoid, sugar, and protein contents as compared to irrigated and non-irrigated conditions [14].

The inhibitory effect of water stress in two maize species was mitigated by inoculation with two endophytic bacterial species i.e., *Burkholderia phytofirmans* PsJN and *Enterobacter* sp. FD17 [35]. Maize seedlings were exposed to water stress after the 45th day of vegetative growth. The inoculated bacterial seedlings exhibited 30% relative water content in leaves along with the significant increase in root and shoot biomass, leaf area, chlorophyll content, and photochemical efficiency of PSII (Photosystem II). However, the *Paraburkholderia phytofirmans* PsJN was found to be more effective as compared to *Enterobacter* sp. FD17. The remedial potential of PGPR against water stress conditions can be used as a tool for sustainable agriculture practices (Table 3).

Treated Plant/PGPR	Stress Alleviation Effect
Maize/ <i>Azospirillum lipoferum</i> , <i>Bacillus</i> Spp.	Increase accumulation of soluble sugar, free amino acids, proline and decrease electrolyte leakage; reduced activity of antioxidant enzyme [36, 37]
Soybean/ <i>Pseudomonas putida</i> H-2-3	Lower the level of abscisic acid and salicylic acid and a higher level of jasmonic acid content; declined superoxide dismutase, flavonoids and radical scavenging activity [38]
Wheat/ <i>Bacillus amyloliquefaciens</i> , <i>Azospirillum brasilense</i> , <i>Rhizobium leguminosarum</i> , <i>Mesorhizobium ciceri</i> , <i>Rhizobium phaseoli</i>	Improved homeostasis; catalase, exopolysaccharides and IAA production [39, 40]
Chickpea/ <i>Pseudomonas putida</i>	Osmolyte accumulation, ROS scavenging ability and stress-responsive gene expressions [41]
Lettuce/ <i>Azospirillum</i> sp.	Promote chlorophyll, ascorbic acid content and antioxidant capacity [42]
Rice/ <i>Trichoderma harzianum</i>	Promote root growth independent of water status and delay drought response [43]

Table 3.
Alleviation of water stress via bioinoculants.

4. Amelioration of salinity via bioinoculants

The soil salinity is naturally occurring phenomena in arid and semi-arid regions. Besides, extensive use of chemical fertilizers and irregular irrigation practices, the continuously changing precipitation pattern is contributing 1–2% salinity to arable lands every year [44]. The plants growing under salinity conditions tend to have high levels of ethylene, stress hormone released in response to both biotic and abiotic stress. As mentioned earlier, PGPR possessing ACC deaminase enzyme can reduce ethylene level by disintegrating ACC (precursor to ethylene) into α -ketobutyrate and ammonia, thereby suppressing the formation of ethylene and ultimately sustaining the plant growth. In doing so, the strains of *Aneurinibacillus aneurinilyticus* and *Paenibacillus sp.* isolated from Garlic (*Allium sativum*) rhizosphere were used for seed bacterization of French bean (*Phaseolus vulgaris*) and evaluated for their ACC deaminase activity [11]. These strains produced more than ~1500 nmol of α -ketobutyrate mg protein⁻¹ h⁻¹ and more than ~30 μ g/ml Indole Acetic Acid (IAA) under saline and nonsalinity stress conditions. Moreover, the seed bacterization in the form of consortia led to a ~ 60% decline in stress stimulated ethylene levels.

Under saline conditions, water uptake is restricted by ionic stress which develops between Na⁺ and K⁺ due to Na⁺ accumulation in aerial parts of plants. Singh and Jha [12] successfully employed inoculation of *Stenotrophomonas maltophilia* SBP-9 on wheat (*Triticum aestivum*) and increased K⁺ uptake by 20–28%. The plant-endophytic relationship has been explored by co-inoculation of *Cicer arietinum* (Chickpea) with *Bacillus subtilis* NUU4 and *Mesorhizobium ciceri* IC53 to alleviate salt stress. These bacterial strains increased proline accumulation which in turn sequestered hydrogen peroxide to strengthen the tolerance level against saline conditions. Besides, coinoculation of endophytes also reduced root rot infection caused by *F. Solani* [45]. The PGPR and their relationship with plants proved to be effective in dealing salt stress simultaneous to other biotic and abiotic stress in given global climatic conditions (Table 4).

5. Amelioration of oxidative stress via osmoregulation

Global climate change is amplifying the intensity of environmental factors, causing a profound disturbance in the plant's biological processes mainly via intractable generation of ROS. A condition called 'oxidative stress' develops when the available concentration of antioxidants within plant cells become insufficient to neutralize an unrestrained number of reactive species (OH⁻, O⁻ and H₂O₂⁻) which leads to disruption of cellular components (lipids, protein, nucleic acids, and metabolites). As a defense mechanism, plants and related rhizobacteria synthesize and release osmolytes when required. Osmolytes are low molecular organic compounds (proline: amino acids, sugars, polyols, methylamines, methylsulfonium compounds, and urea) mainly aiming at neutralization of the osmotic pressure of cells under stress conditions. These plants and PGPR synthesized osmolytes synergistically act to capture ROS. The prominent PGPR strains such as *Azospirillum brasilense* SP-7 and *Herbaspirillum seropedicae* Z-152 managed to retain relatively higher water content in vegetal tissues of inoculated/non-irrigated maize plants as compared to non-inoculated/irrigated maize plants. The plants inoculated with *H. seropedicae* had a higher proline accumulation rate (fourfold) as compared to those inoculated with *A. brasilense* (two-fold), thus indicating better osmoregulation under drought conditions. However, the amount of proline in inoculated plants

Treated Plant/PGPR	Stress Alleviation Effect
Groundnut/ <i>Brachybacterium saurashtrense</i> , <i>Brevibacterium casei</i> , <i>Haererohalobacter</i>	Higher K ⁺ /Na ⁺ ratio and higher Ca ²⁺ , phosphorus, and nitrogen content, higher Shoot and root concentration of auxin [43, 46]
Mung bean/ <i>Rhizobium</i> , <i>Pseudomonas</i>	ACC-deaminase activity [47]
Wheat/ <i>Azospirillum</i> Sp., <i>Pseudomonas</i> Sp. <i>Serratia</i> Sp.	Osmotic adjustments via proline and soluble sugars [48] ACC deaminase activity, reduced ethylene level [49]
Maize/ <i>Pseudomonas</i> , <i>Enterobacter</i>	More N, P, K uptake and high K ⁺ –Na ⁺ ratios [50]
Rice/ <i>Pseudomonas pseudoalcaligenes</i> , <i>Bacillus pumilus</i> , <i>Bacillus amyloliquefaciens</i>	Reduced reactive oxygen species, lipid peroxidation and superoxide dismutase activity, modulating differential transcription in at least 14 genes [51, 52]
Lettuce/ <i>Azospirillum</i>	Promoted ascorbic acid content antioxidant capacity [42]

Table 4.
Alleviation of salinity stress via bioinoculants.

was relatively lower than control plants in both water-stressed and well-watered conditions. This decreased concentration of proline in inoculated plants may be attributed to a balanced amount of osmolytes to the available ROS leading to better osmoregulation in water-stressed conditions. Also, inoculated plants have lower ethylene content than that of control plants. However, the gene expression of ZmVP14 responsible for the biosynthesis of abscisic acid was suppressed in inoculated plants [53], allowing stomata to remain open for better CO₂ assimilation even in drought conditions also supported by other researchers [35].

Inoculation of chickpea varieties with *Pseudomonas putida*, also indicated reduced proline concentration with a reduced level of 114% in BG-362 and 214% in BG-1003 on the seventh day of water stress [41, 53]. However, in *Capsicum annum* inoculated with *Burkholderia cepacia*, the accumulation of proline content was higher (0.143 mmoles gm⁻¹) as compared to the control plant (0.065 mmoles gm⁻¹). Co-inoculation of *Azotobacter chroococcum* and *Azospirillum brasilense* significantly improved accumulation of proline and other osmolytes along with physic-chemical characteristics of *Mentha pulegium* L. [54]. Other than proline, some soluble sugars play a vital role in maintaining photosynthesis in leaves under water stress, therefore, also termed ‘osmoprotectant’. However, the reduced level of soluble sugar can become deleterious to the cellular membranes of plants. The co-inoculation of PGPR (*Bacillus subtilis*, *Bacillus thuringiensis*, and *Bacillus megaterium*) with PGRs (SA & Put) in chickpea seedlings prompt the accumulation of soluble sugar along with chlorophyll and protein content (majorly proline), thus, regularizing photosynthesis under water stress [14]. The accumulated soluble sugar also acts as a signaling molecule, where it triggers the activation and gene expression of certain genes relevant to photosynthesis in plants. Hence, osmolytes being ‘abiotic stress busters’ are pivotal in maintaining homeostatic equilibrium at the molecular and cellular level to enhance stress tolerance in plants [55].

6. Amelioration of cold stress via psychrotrophic microbes

Due to global warming, extreme environmental conditions are occurring more frequently including cold stress. In northern parts of the world, the cold weather is unexpectedly intensifying specifically due to the Polar vortex bringing cold Arctic

air towards southern parts of the planet, bringing more intense winters in some parts of the world. The biomes of extreme environments serve as excellent habitats for archaea, bacteria, and fungi. Agro-ecosystems at high altitudes are vulnerable to low productivity due to low surface temperature resulting in low soil temperature and in turn low soil fertility [56]. High altitude soils comprised diverse microbes that are capable of surviving cold environments mainly due to synthesis and assimilation of cryoprotective compounds such as metabolites (proline), anthocyanin (secondary metabolite), and carbohydrates (trehalose). The induction of such stress-resistance against abiotic stress is also coined with a term called ‘Induced Systematic Resistance’ (ISR). An *in vitro* inoculation of *Vitis vinifera* L. (grapevine) explants with *Burkholderia phytofirmans* strain PsJN successfully developed chilling resistance mainly via accumulation of increased level of sugars, proline, and phenolics as compared to non-inoculated plantlets [57].

Several biotechnological and microbiological attempts have been ongoing for the last two decades to utilize the cold-tolerant (psychrotrophic) or cold-loving (psychrophilic) microbes to improve the agricultural practices in chilling mountainous environments. Psychrophilic microbes inhabiting cold environments majorly belong to phyla *Verrucomicrobia*, *Thaumarchaeota*, *Spirochaetes*, *Proteobacteria*, *Planctomycetes*, *Nitrospirae*, *Mucoromycota*, *Gemmatimonadetes*, *Firmicutes*, *Euryarchaeota*, *Cyanobacteria*, *Chloroflexi*, *Chlamydiae*, *Basidiomycota*, *Bacteroidetes*, *Ascomycota*, and *Actinobacteria*. Like thermotolerant PGPR, the psychrotrophic PGPR explicit various attributes like ACC deaminase activity, solubilization of micronutrients (phosphorus, potassium, and zinc), biological N₂ fixation, and production of ammonia, hydrogen cyanide, indole-3-acetic acid, and Fe-chelating compounds (**Figure 2**). The alleviation of cold stress on the production of cereal crops such as wheat was investigated by inoculation with *Bacillus* spp. CJCL2 and RJGP41 strains isolated from Qinghai-Tibetan plateau [58]. The psychrophilic activity of this strain was also compared to temperate *B. velezensis* FZB42 at a temperature range of 14°C, 10°C, and 4°C after 4 h post-inoculation. The cold-tolerant strain CJCL2 produced the finest biofilm structure as compared to RJGP41 which produced slight biofilms at 96 h post-inoculation at 4°C, whereas FZB42 failed to develop any biofilm at 4°C. The basic mechanism against cold stress was based on the regulation of abscisic acid, lipid peroxidation and proline accumulation, and a lower level of ROS. Moreover, genetic expression for ACC deaminase, glucose dehydrogenase encoding phosphate solubilization, and phytohormones was quite effective and led to the improvement in plant growth under cold stress. The cold-tolerant *Bacillus* spp. maintained osmotic balance in a plant cell by synthesizing glycine betaine, a major osmoprotectant and produced due to OpuAC gene expression. All in all linear four-fivefold increase in the gene expression of cold shock proteins (CspB, CspC, and CspD) observed in CJCL2 and RJGP41 inoculated wheat

Figure 2.
Key mechanisms targeted by plant growth promoting rhizobacteria.

plants may trigger the modifications in RNA. Also, these proteins may stabilize the secondary structure of nucleotides leading to improved cellular components.

6.1 Plant growth promoting rhizobacteria and root-nodule symbiosis

The PGPR possessing root-nodule symbiosis has also been reported to exhibit psychrotolerant traits at 5°C [59]. These PGPRs were isolated from the root nodule of *Pisum sativum* L. (Pea), cultivated in the Northern Indian plains. Out of nineteen tested strains, four exhibited cold tolerance mainly by producing phytohormone IAA in the range of 62.7–198.1 µg/ml. The inoculation of edible crops growing under adverse climatic conditions with PGPR producing phytohormones showed promising results [60]. The *Serratia nematodiphila* PEJ1011 strain was inoculated to *Capsicum annuum* L. (pepper) at a low temperature of 5°C. The inoculated plants exhibited higher endogenous GA₄ content grown both at normal and low temperatures. Moreover, PGPR increased abscisic acid (phytohormone) level and reduced jasmonic acid and salicylic acid (phytohormones) contents to regulate plant adaptation and growth at low temperature (Figure 3).

6.2 Plant growth promoting rhizobacteria and chlorophyll pigments

Chloroplasts containing chlorophyll pigments for photosynthesis are the main organelles affected by the cold. Low temperature also led to stomatal closure in many cold-tolerant plants like *Arabidopsis thaliana*. Although the stomatal closure prevents leaf dehydration, CO₂ uptake is inhibited which eventually reduces photosynthesis. The inoculation of *A. thaliana* with endophytic *Burkholderia phytofirmans* strain PsJN (Bp PsJN) prevented disruption of the plasma membrane at 0°C, –1°C, and –3°C and induced cell wall strengthening in leaf cells [62]. Moreover, after night stress, the bacteria led to better photosynthetic pigment content. Similarly, *Vitis vinifera* L. (grapevine) cultivated in temperate and cool climates contains CBF4 gene which is a homolog to *A. thaliana* CBF1 accumulates carbohydrates and proline on exposure to low temperature [63]. The *V. vinifera* plantlets were inoculated with endophytic *B. phytofirmans* strain PsJN at 4°C. As expected, the bacterial strain had induced cold resistance principally by enhancing gene expression of CBF (specifically CBF4) genes, along with an accumulation of anti-freeze proteins (PR proteins) and metabolite level. The bacterial strain was found to be highly effective in inhibiting hydrogen peroxide (H₂O₂). However, the inoculated plantlets indicated high H₂O₂ accumulation

Figure 3.
Impact of plant growth promoting rhizobacteria on nutrient acquisition [61].

in the first three days of cold stress followed by H₂O₂ elimination after a week, further evidenced by a reduction in the level of metabolites after 1 week indicating inhibition of ROS via scavenging process carried out by endophytic PsJN strain.

6.3 Plant growth promoting rhizobacteria and sugar deposition

Certain bacteria, fungi, vascular plants, and invertebrate animals synthesize trehalose or mycose for energy consumption and to tolerate cold and water stress. Trehalose is a disaccharide sugar consisting of two glucose molecules mainly synthesized via phosphorylated intermediate trehalose 6-phosphate (T6P). Being capable of inducing cryoprotective compounds, *B. phytofirmans* strain PsJN was inoculated in grapevine to observe its impact on the accumulation of T6P and trehalose at 4°C. After 120 h of cold exposure, the level of T6P was increased in roots (>0.4 nmol. g⁻¹ FW) stems (>0.7 nmol. g⁻¹ FW), and leaves (>1.7 nmol. g⁻¹ FW) of inoculated plantlets. T6P acts as a signal metabolite in plants responding in particular to changes in sucrose level, thereby a strong correlation had been observed between T6P and sucrose content in all plant organs in both chilled and non-chilled plants. The authors were uncertain about the role of trehalose against chilling because the overall accumulation of trehalose was only found in leaves in low amounts i.e., <15 nmol. g⁻¹. Localized trehalose accumulation might have protected cell membranes against cold-induced dehydration [64, 65]. However, the organ-specific accumulation may indicate trehalose acted as an osmolyte [66]. Previous studies related to drought stress indicated overexpression of genes for the synthesis of trehalose which under water-stress inhibited dehydration [67]. Although psychrotolerant PGPRs are proved to be effective in inducing tolerance against cold stress, the quest of mechanism underlying is still being investigated by the researchers.

7. Conclusions

Despite awareness and efforts being planned or carried out, global temperature, a dynamic entity itself is hard to reverse to its previous levels in 1990. Instead, plant species have to acclimatize themselves to prevailing temperature variations. A natural symbiotic relationship between PGPR and plant growth is known for decades, has been widely considered and accepted as the most convenient and reliable approach to deal with climate-based environmental stresses. This chapter, therefore, quoted and discussed several research studies utilizing PGPR to alter physiological and biochemical processes occurring at inter and intracellular levels to boost the plant's defense mechanism. However, the complexity of the mechanism underlying is still in focus to fully harness the potential of PGPR for sustainable agricultural practices and for sustaining natural plant diversity.

Conflict of interest

The authors declare no conflict of interest.

Abbreviations

PGPR	Plant Growth Promoting Rhizobacteria
1-Aminocyclopropane-1-carboxylate Deaminase	ACC deaminase

ABA	Absciscic Acid
SA	Salicylic Acid
IAA	Indole-3-Acetic Acid
VOCs	Volatile Organic Compounds
ABA	Absciscic Acid (ABA)
HSP	Heat Shock Protein
SOD	Superoxide Dismutase
POD	Peroxidase
CAT	Catalase
ROS	Reactive Oxygen Species
RWC	Relative Water Content
EPS	Exopolysaccharide
PGRs	Plant Growth Retardants (PGRs)
Put	Putrescine
ISR	Induced Systematic Resistance

Author details

Rafia Younas¹, Shiza Gul¹, Rehan Ahmad¹, Ali Raza Khan¹, Mumtaz Khan², Tauseef Anwar³ and Huma Qureshi^{4*}

1 Department of Environmental Science, Institute of Soil and Environmental Sciences, Gomal University, Dera Ismail Khan, Pakistan

2 Directorate of Regional Services, Allama Iqbal Open University, Islamabad, Pakistan

3 Department of Botany, The Islamia University of Bahawalpur, Bahawalnagar Campus (New Campus), Bahawalnagar, Pakistan

4 Institute of Biological Sciences, Gomal University, Dera Ismail Khan, Pakistan

*Address all correspondence to: drhuma@gu.edu.pk

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Estrada F, Botzen WJW, Calderon-Bustamante O. The assessment of impacts and risks of climate change on agriculture (AIRCCA) model: a tool for the rapid global risk assessment for crop yields at a spatially explicit scale. *Spatial Economic Analysis*. 2020; 15:262-279. DOI: 10.1080/17421772.2020.1754448
- [2] Dudal R. Inventory of the major soils of the world with special reference to mineral stress, In: *Plant Adaptation to Mineral Stress in Problem Soils*, Wright MJ, (Ed.). Cornell University, Agricultural Experiment Station, Ithaca, NY. 1976; p. 3-13.
- [3] Vinocur B, Altman B. Recent advances in engineering plant tolerance to abiotic stress: achievements and limitations. *Current Opinion in Biotechnology*. 2005; 16:1-10. DOI: 10.1016/j.copbio.2005.02.001
- [4] Bashir T, Naz S, Bano A. Plant Growth Promoting Rhizobacteria in combination with Plant Growth Regulators attenuates the effect drought stress. *Pakistan Journal of Botany*. 2020; 52:783-792. DOI: 10.30848/PJB2020-3(17)
- [5] Khan N, Bano A, Zandi P. Effects of exogenously applied plant growth regulators in combination with PGPR on the physiology and root growth of chickpea (*Cicer arietinum*) and their role in drought tolerance. *Journal of Plant Interactions*. 2018; 13:239-247. DOI: 10.1080/17429145.2018.1471527
- [6] Kumar A, Patel JS, Meena VS, Srivastava R. Recent advances of PGPR based approaches for stress tolerance in plants for sustainable agriculture. *Biocatalysis and Agricultural Biotechnology*. 2019; 20:101271. DOI: 10.1016/j.bcab.2019.101271
- [7] Mega R, Abe F, Kim JS, Tsuboi T, Tanaka K, Kobayashi H, Sakata Y, Hanada K, Tsujimoto H, Kikuchi J, Cutler SR, Okamoto M. Tuning water-use efficiency and drought tolerance in wheat using abscisic acid receptors. *Nature Plants*. 2019; 5:153-159. DOI: 10.1038/s41477-019-0361-8
- [8] Vurukonda SSKP, Vardharajula S, Shrivastava M, Ali SKZ. Enhancement of drought stress tolerance in crops by plant growth promoting rhizobacteria. *Microbiological Research*. 2015; 184:13-24. DOI: 10.1016/j.micres.2015.12.003
- [9] Mukhtar T, Shafiq ur Rehman, Smith D, Sultan T, Seleiman MF, Alsadon AA, Amna, Ali S, Chaudhary HJ, Solieman THI, Ibrahim AA, Saad MAO. Mitigation of heat stress in *Solanum lycopersicum* L. by ACC-deaminase and exopolysaccharide producing *Bacillus cereus*: Effects on biochemical profiling. *Sustainability*. 2020; 12:2159. DOI: 10.3390/su12062159
- [10] Danish S, Zafar-ul-Hye M. Co-application of ACC-deaminase producing PGPR and timber-waste biochar improves pigments formation, growth and yield of wheat under drought stress. *Scientific Reports*. 2019; 9:5999. DOI: 10.1038/s41598-019-42374-9
- [11] Gupta S, Pandey S. ACC Deaminase producing bacteria with multifarious plant growth promoting traits alleviates salinity stress in French bean (*Phaseolus vulgaris*) plants. *Frontiers in Microbiology*. 2019; 10:1506. DOI: 10.3389/fmicb.2019.01506
- [12] Singh RP, Jha PN. The PGPR *Stenotrophomonas maltophilia* SBP-9 augments resistance against biotic and abiotic stress in wheat plants. *Frontiers in Microbiology*. 2017; 8:1945. DOI: 10.3389/fmicb.2017.01945
- [13] Rybakova D, Rack-Wetzlinger U, Cernava T, Schaefer A, Schmuck M,

Berg G. Aerial warfare: A volatile dialogue between the plant pathogen *Verticillium longisporum* and its antagonist *Paenibacillus polymyxa*. *Frontiers in Plant Science*. 2017; 8:1294. DOI:10.3389/fpls.2017.01294

[14] Khan N, Bano A, Rahman MA, Guo J, Kang Z, Babar MA. Comparative physiological and metabolic analysis reveals a complex mechanism involved in drought tolerance in chickpea (*Cicer arietinum* L.) induced by PGPR and PGRs. *Scientific Reports*. 2019; 9: 2097. DOI: 10.1038/s41598-019-38702-8

[15] Schmidt R, Köberl M, Mostafa A, Ramadan EM, Monschein M, Jensen KB, Bauer R, Berg G. Effects of bacterial inoculants on the indigenous microbiome and secondary metabolites of chamomile plants. *Frontiers in Microbiology*. 2014; 5:64. Doi: 10.3389/fmicb.2014.00064

[16] Drigo B, Kowalchuk GA, Van Veen JA (2008) Climate change goes underground: effects of elevated atmospheric CO₂ on microbial community structure and activities in the rhizosphere. *Biology and Fertility of Soil*. 44:667-679. DOI: 10.1007/s00374-008-0277-3

[17] Rennenberg H, Loreto F, Polle A, Brilli F, Fares S, Beniwal RS, Gessler A. Physiological responses of forest trees to heat and drought. *Plant Biology*. 2006; 8:556-571. DOI: 10.1055/s-2006-924084

[18] Baniwal SK, Bharti K, Chan KY, Fauth M, Ganguli A, Kotak S, von Koskull-Doring P. Heat stress response in plants: a complex game with chaperones and more than twenty heat stress transcription factors. *Journal of Biosciences*. 2004; 29:471-487. DOI: 10.1007/BF02712120

[19] Vanaja M, Ramakrishna YS, Rao GGSN, Rao KV, Subbarao VM. Climate change and dryland agriculture. *Dryland ecosystems: an Indian*

perspective. Central Arid Zone Research Institute (CAZRI) and Arid Forest Research Institute (AFRI), Jodhpur. 2007; p. 23-24.

[20] Bensalim S, Nowak J, Asiedu SK. A plant growth promoting rhizobacterium and temperature effects on performance of 18 clones of potato. *American Journal of Potato Research*. 1998; 75:145-152. DOI: 10.1007/BF02895849

[21] Duc NH, Csintalan Z, Posta K. Arbuscular mycorrhizal fungi mitigate negative effects of combined drought and heat stress on tomato plants. *Plant Physiology and Biochemistry*. 2018; 132:297-307. DOI: 10.1016/j.plaphy.2018.09.011

[22] El-Daim IA, Bejai S, Meijer J. Improved heat stress tolerance of wheat seedlings by bacterial seed treatment. *Plant Soil*. 2014; 379:337-350. DOI: 10.1007/s11104-014-2063-3

[23] Mishra PK, Bisht SC, Bisht JK, Bhatt JC. Cold-tolerant PGPRs as bioinoculants for stress management. In *Bacteria in Agrobiolgy: Stress Management*; Maheshwari DK. Springer: Berlin/Heidelberg, Germany, 2012; p. 95-118.

[24] Lindow SE. The role of bacterial ice nucleation in frost injury to plants. *Annual Review of Phytopathology*. 1983; 21:363-384. DOI: 10.1146/annurev.py.21.090183.002051

[25] Theocharis A, Bordiec S, Fernandez O, Paquis S, Dhondt-Cordelier S, Baillieul F, Clément C, Barka EA. *Burkholderia phytofirmans* PsJN primes *Vitis vinifera* L. and confers a better tolerance to low non-freezing temperatures. *APS Publications*. 2012; 2:241-249. DOI: <https://doi.org/10.1094/MPMI-05-11-0124>

[26] Park YG, Mun BG, Kang SM, Hussain A, Shahzad R, Seo CW,

- Kim AY, Lee SU, Oh KY, Lee DY, Lee IJ, Yun BW. *Bacillus aryabhatai* SRB02 tolerates oxidative and nitrosative stress and promotes the growth of soybean by modulating the production of phytohormones. PLoS ONE, 2017; 12: e0173203. DOI: 10.1371/journal.pone.0173203
- [27] Verma P, Yadav AN, Khannam KS, Mishra S, Kumar S, Saxena AK Suman A Appraisal of diversity and functional attributes of thermotolerant wheat associated bacteria from the peninsular zone of India. Saudi Journal of Biological Science. 2019; 26:1882-1895. DOI: 10.1016/j.sjbs.2016.01.042
- [28] Ali SKZ, Sandhya V, Grover K, Rao LV, Venkateswarlu B. *Pseudomonas* sp. strain AKM-P6 enhances tolerance of sorghum seedlings to elevated temperatures. Biology and Fertility of Soils. 2009; 46:45-55. DOI: 10.1007/s00374-009-0404-9
- [29] Saleem AR, Brunetti C, Khalid A, Rocca GD, Raio A, Emiliani G, De Carlo A, Mahmood T, Centritto M. Drought response of *Mucuna pruriens* (L.) DC. inoculated with ACC deaminase and IAA producing rhizobacteria. PLoS ONE. 2018; 13:e0191218. DOI: 10.1371/journal.pone.0191218
- [30] Babalola OO. Beneficial bacteria of agricultural importance. Biotechnology Letters. 2010; 32:1559-1570. DOI: 10.1007/s10529-010-0347-0
- [31] Mayak S, Tirosh T, Glick BR. Plant growth-promoting bacteria that confer resistance to water stress in tomatoes and peppers. Plant Science. 2004; 166:525-530. DOI: 10.1016/j.plantsci.2003.10.025
- [32] Chandra D, Srivastava R, Gupta VV, Franco CM, Sharma AK. Evaluation of ACC deaminase producing rhizobacteria to alleviate water stress impacts in wheat (*Triticum aestivum* L.) plants. Canadian Journal of Microbiology. 2019; 65:387-403. DOI: 10.1139/cjm-2018-0636
- [33] Naseem H, Ahsan M, Shahid MA, Khan N Exopolysaccharides producing rhizobacteria and their role in plant growth and drought tolerance. Journal of Basic Microbiology (2018); 58:1009-1022. DOI: 10.1002/jobm.201800309
- [34] Maxton A, Singh P, Masih SA. ACC deaminase-producing bacteria mediated drought and salt tolerance in *Capsicum annuum*. Journal of Plant Nutrition. 2018; 41:574-583. DOI: 10.1080/01904167.2017.1392574
- [35] Naveed M, Mitter B, Reichenauer TG, Wieszorek K, Sessitsch A Increased drought stress resilience of maize through endophytic colonization by *Burkholderia phytofirmans* PsJN and *Enterobacter* sp. FD17. Journal of Environmental and Experimental Botany. 2014; 97:30-39. DOI: 10.1016/j.envexpbot.2013.09.014
- [36] Bano Q, Ilyas N, Bano A, Zafar N, Akram A, Hassan F. Effect of Azospirillum inoculation on maize (*Zea mays* L.) under drought stress. Pakistan Journal of Botany. 2013; 45: 13-20. DOI: 10.3390/microorganisms5030041
- [37] Vardharajula S, Ali SZ, Grover M, Reddy G, Bandi V. Drought-tolerant plant growth promoting *Bacillus* spp.: effect on growth, osmolytes, and antioxidant status of maize under drought stress. Journal of Plant Interactions. 2011; 6:1-14. DOI: 10.1080/17429145.2010.535178
- [38] Kang S, Radhakrishnan R, Khan AL, Kim M, Park J, Kim B, Shin D, Lee I. Gibberellin secreting rhizobacterium, *Pseudomonas putida* H-2-3 modulates the hormonal and stress physiology of soybean to improve the plant growth under saline and drought conditions. Plant Physiology and Biochemistry. 2014; 84:115-124. DOI: 10.1016/j.plaphy.2014.09.001

- [39] Kasim WA, Osman ME, Omar MN, El-Daim IA, Bejai S, Meijer J. Control of drought stress in wheat using plant-growth-promoting bacteria. *Journal of Plant Growth Regulation*. 2013; 32:122-130. DOI: 10.1007/s00344-012-9283-7
- [40] Hussain MB, Zahir ZA, Asghar HN, Asghar M. Can catalase and exopolysaccharides producing rhizobia ameliorate drought stress in wheat? *International Journal of Agriculture and Biology*. 2014; 16: 3-13.
- [41] Tiwari S, Lata C, Chauhan PS, Nautiyal CS. *Pseudomonas putida* attunes morphophysiological, biochemical and molecular responses in *Cicer arietinum* L. during drought stress and recovery. *Plant Physiology and Biochemistry*. 2016; 99:108-117. DOI: 10.1016/j.plaphy.2015.11.001
- [42] Gabriela F, Casanovas EM, Quillehauquy V, Yommi AK, Goñi MG, Roura SI, Barassi CA. *Azospirillum* inoculation effects on growth, product quality and storage life of lettuce plants grown under salt stress. *Scientia Horticulturae*. 2015; 195:154-162. DOI: 10.1016/j.scienta.2015.09.015
- [43] Shukla PS, Agarwal PK, Jha B. Improved salinity tolerance of *Arachis hypogaea* (L.) by the interaction of halotolerant plant-growth-promoting rhizobacteria. *Journal of Plant Growth Regulation*. 2012; 31, 195-206. DOI: 10.1007/s00344-011-9231-y
- [44] FAO. Salt-Affected soils from sea water intrusion: strategies for rehabilitation and management: Report of the regional workshop. Bangkok. 2005.
- [45] Egamberdieva D, Wirth SJ, Shurigin VV, Hashem A, Abd-Allah EF. Endophytic bacteria improve plant growth, symbiotic performance of chickpea (*Cicer arietinum* L.) and induce suppression of root rot caused by *Fusarium solani* under salt stress. *Frontiers in Microbiology*. 2017; 8:1887. DOI: 10.3389/fmicb.2017.01887
- [46] Shukla N, Awasthi RP, Rawat L, Kumar J. Biochemical and physiological responses of rice (*Oryza sativa* L.) as influenced by *Trichoderma harzianum* under drought stress. *Plant Physiology and Biochemistry*. 2012; 54, 78-88. DOI: 10.1016/j.plaphy.2012.02.001
- [47] Ahmad M, Zahir ZA, Asghar HN, Asghar M. Inducing salt tolerance in mung bean through coinoculation with rhizobia and plant-growth-promoting rhizobacteria containing 1-aminocyclopropane-1-carboxylate deaminase. *Canadian Journal of Microbiology*. 2011; 57:578-589. DOI: 10.1139/w11-044
- [48] Chang P, Gerhardt KE, Huang XD, Yu XM, Glick BR, Gerwing PD, Greenberg BM. Plant growth-promoting bacteria facilitate the growth of barley and oats in salt-impacted soil: implications for phytoremediation of saline soils. *International Journal of Phytoremediation*. 2014; 16:1133-1147. DOI: 10.1080/15226514.2013.821447
- [49] Zahir ZA, Ghani U, Naveed M et al. Comparative effectiveness of *Pseudomonas* and *Serratia* sp. containing ACC-deaminase for improving growth and yield of wheat (*Triticum aestivum* L.) under salt-stressed conditions. *Archives of Microbiology*. 2009;191: 415-424. DOI: 10.1007/s00203-009-0466-y.
- [50] Nadeem SM, Zahir ZA, Naveed M, Arshad M. Rhizobacteria containing ACC-deaminase confer salt tolerance in maize grown on salt-affected fields. *Canadian Journal of Microbiology*. 2009; 55:1302-1309. DOI: 10.1139/w09-092
- [51] Jha Y, Subramanian RB. PGPR regulate caspase-like activity, programmed cell death, and antioxidant enzyme activity in paddy under salinity.

Physiology and Molecular Biology of Plants. 2014; 20:201-207. DOI: 10.1007/s12298-014-0224-8

[52] Nautiyal CS, Srivastava S, Chauhan PS, Seem K, Mishra A, Sopory SK. Plant growth-promoting bacteria *Bacillus amyloliquefaciens* NBRISN13 modulates gene expression profile of leaf and rhizosphere community in rice during salt stress. Plant Physiology and Biochemistry. 2013; 66:1-9. DOI: 10.1016/j.plaphy.2013.01.020

[53] Curá JA, Franz DR, Filosofía JE, Balestrasse KB, Burgueño LE. Inoculation with *Azospirillum* sp. and *Herbaspirillum* sp. bacteria increases the tolerance of maize to drought stress. Microorganisms. 2017; 5:41. DOI: 10.3390/microorganisms5030041

[54] Asghari B, Khademain R, Sedaghati B. Plant growth promoting rhizobacteria (PGPR) confer drought resistance and stimulate biosynthesis of secondary metabolites in pennyroyal (*Mentha pulegium* L.) under water shortage condition. Scientia Horticulturae. 2020; 263:109132. DOI: 10.1016/j.scienta.2019.109132

[55] Iqbal N, Nazar R, Khan NA. Osmolytes and plants acclimation to changing environment: Emerging omics technologies. Springer, India. 2016.

[56] Bhandari G. Microbes adapted to cold and their use as biofertilizers for mountainous regions. Recent Advancements in Microbial Diversity. 2020; 197-215.

[57] Barka EA, Nowak J, Clement C. Enhancement of chilling resistance of inoculated grapevine plantlets with a plant growth promoting rhizobacterium *Burkholderia phytofirmans* strain PsJN. Applied and Environmental Microbiology. 2006; 72:7246-7252. DOI: 10.1128/AEM.01047-06

[58] Zubair M, Hanif A, Farzand A, Sheikh TMM, Khan AR, Suleman M, Ayaz M, Gao X. Genetic screening and expression analysis of Psychrophilic *Bacillus spp.* reveal their potential to alleviate cold stress and modulate phytohormones in wheat. Microorganisms. 2019; 7:337. DOI: 10.3390/microorganisms7090337

[59] Meena RK, Singh RK, Singh NP, Meena SK, Meena VS. Isolation of low temperature surviving plant growth-promoting rhizobacteria (PGPR) from pea (*Pisum sativum* L.) and documentation of their plant growth promoting traits. Biocatalysis and Agricultural Biotechnology. 2015; 4:806-811. DOI: 10.1016/j.bcab.2015.08.006

[60] Kang SM, Khan AL, Waqas M, You YH, Hamayun M, Joo GJ, Shahzad R, Choi KS, Lee IJ. Gibberellin-producing *Serratia nematodiphila* PEJ1011 ameliorates low temperature stress in *Capsicum annuum* L. European Journal of Soil Biology. 2015; 68:85-93. DOI: 10.1016/j.ejsobi.2015.02.005

[61] Vacheron J, Desbrosses G, Bouffaud M, Touraine B, Moenne-Loccoz Y, Muller D, Legendre L, Wisniewski-Dyé F, Prigent-Combaret C. Plant growth-promoting rhizobacteria and root system functioning. Frontiers in Plant Science. 2013; 4:1-19. DOI: 10.3389/fpls.2013.00356

[62] Su F, Jacquard C, Villaume S, Michel J, Rabenoelina F, Clément C, Barka EA, Dhondt-Cordelier S, Vaillant-Gaveau N. *Burkholderia phytofirmans* PsJN reduces impact of freezing temperatures on photosynthesis in *Arabidopsis thaliana*. Frontiers in Plant Science. 2015; 6:810. DOI: 10.3389/fpls.2015.00810

[63] Theocharis A, Bordiec S, Fernandez O, Paquis S, Dhondt-Cordelier S, Baillieul F,

Barka EA. *Burkholderia phytofirmans* PsJN primes *Vitis vinifera* L. and confers a better tolerance to low nonfreezing temperatures. *Molecular Plant-Microbe Interactions*. 2012; 25:241-249. DOI: 10.1094/MPMI-05-11-0124

[64] Crowe JH. Trehalose as a 'chemical chaperone' fact and fantasy. In: Back N, Cohen IR, Lajtha A, Lambris JD, Paoletti R (eds). *Molecular aspects of the stress response: chaperones, membranes and networks*. Springer, New York. 2007; 143-158.

[65] Ruelland E, Vaultier MN, Zachowski A, Hurry V, Kader JC, Delseny M. Cold signaling and cold acclimation in plants. *Advances in Botanical Research* *Advances in Botanical Research*. 2009; 49:35-150.

[66] Fernandez O, Béthencourt L, Quero A, Sangwan RS, Clément C. Trehalose and plant stress responses: friend or foe? *Trends in Plant Science*. 2010; 15:409-417.

[67] Karim S, Aronsson H, Ericson H, Pirhonen M, Leyman B, Welin B, Mantyla E, Palva E, van Dijck P, Holmstrom KO. Improved drought tolerance without undesired side effects in transgenic plants producing trehalose. *Plant Molecular Biology*. 2007; 64:371-386. DOI: 10.1007/s11103-007-9159-6