We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Avifauna in Relation to Habitat Disturbance in Wildlife Management Areas of the Ruvuma Miombo Ecosystem, Southern Tanzania

Ally K. Nkwabi, John K. Bukombe, Hamza K. Kija, Steven D. Liseki, Sood A. Ndimuligo and Pius Y. Kavana

Abstract

Understanding of relative distribution of avifauna provides insights for the conservation and management of wildlife in the community managed areas. This study examined relative diversity, abundance, and distribution of avifauna in selected habitat types across five Wildlife Management Areas of the Ruvuma landscape in miombo vegetation, southern Tanzania. Five habitat types were surveyed during the study: farmland, swamps, riverine forest, dense and open woodland. Transect lines, mist-netting, and point count methods were used to document 156 species of birds in the study sites. Descriptive statistics and Kruskal-Wallis tests were used to compare species richness and diversity across habitat types. We found differences in avifaunal species distribution in the study area whereby farmland had the highest abundance of avifauna species and lowest in the riverine forest. These results suggest that variations of avifauna species abundance, diversity, and distribution could be attributed by human activities across habitat types; due to the reason that habitats with less human encroachment had good species diversity and richness. Therefore, to improve avitourism and avoid local extinction of species, we urge for prompt action to mitigate species loss by creating awareness in the adjacent community through conservation education on the importance of protecting such biodiversity resources.

Keywords: Avifauna, diversity, conservation, habitat destruction, wildlife management areas, miombo

1. Introduction

The miombo ecosystems are known worldwide for their higher biodiversity [1, 2]. Woodlands in the miombo ecosystems are dominated by trees of the genera *Brachystegia*, *Julbernardia*, and *Isobertinia* Leguminosae, subfamily Caesalpinioideae [1, 3]. The woodlands cover between 2.7 and 3.6 million km² in 11 African countries [2, 4–6]. In Tanzania, this vegetation type covers more than 90% of forested land

[4, 7–10], and some of the miombo woodlands are found within several of the iconic protected areas including Selous Game Reserves and the Mikumi, Ruaha, Nyerere National Park as well as the Ruvuma Landscape in southern Tanzania. The ecological services it provides include: the provision of forage for wild and domestic animals, nesting sites for birds, water catchments, carbon sequestration, and biodiversity conservation in general and is archived due to the presence of habitat heterogeneity in particular flora diversity that exists in the miombo areas [3, 4].

Floral species compositions are a very important component to determine the distribution and diversity of avifauna communities [11]. Bird species diversity in savannah landscapes increases with an increase in vegetation/habitat heterogeneity in the miombo woodlands [5, 6]. In heterogeneous habitats, some avian species tend to show preference on certain habitat types, which also influence avifaunal diversity, abundance, and distribution across landscapes [7, 8, 12]. For example, miombo pied barbet (*Tricholaema frontata*), miombo rock thrush (*Monticola angolensis*), stierling's wren warbler (*Calamonastes stierlingi*), racket-tailed roller (*Coracias spatulatus*) and white-tailed blue-flycatcher (*Elminia albicauda*) prefer miombo woodland, only stierling's wren warbler and racket-tailed roller were observed during data collection other species listed here were not recorded during this study possibly due to habitat degradation.

The Ruvuma landscape in Tunduru District, in southern Tanzania encompasses five Wildlife Management Areas (WMAs) namely: Mbarang'andu, Kimbanda, and Kisungule in Namtumbo District, Nalika and Chingoli WMAs in Tunduru District (**Figure 1**). It borders the Selous Game Reserve and Nyerere National Park in the north and the Niassa National Reserve (Mozambique) to the south. The Ruvuma River forms an international boundary between Tanzania and Mozambique within Namtumbo and Tunduru districts [13]. The two protected areas rely on the presence of the five Wildlife Management Areas as they provide dispersal and movement area (corridor) to Niassa National Reserve in Mozambique and to Nyerere National Park. Habitat destruction by humans is a serious threat that alters the integrity of ecosystems [8], also affects vegetation cover. It is possible that human activities occurring in the miombo woodland resulted in land cover change [7, 9, 10, 14, 15]. Currently, the Wildlife Management Areas (WMAs) of the Ruvuma region in southern Tanzania undergo fragmentations caused by human activities which include uncontrolled wildfires, collection of fuel wood, charcoal, timber, illegal hunting, cattle grazing, and agriculture. In this area, communities have formulated the Wildlife Management Area (WMA), which is the form of community-based conservation which ensures villagers or communities rich in wildlife sustainably conserve, utilize and benefit from wildlife. Wildlife Management Areas are formed within village land from which villagers set aside a piece of land purposely for sustainable conservation and utilization of wildlife resources. The Tanzania government actualized WMAs for the local community to participate in wildlife management and conserve wildlife habitats in the communal land.

Apart from the study investigated on abundance, nesting and habitat of the white-browed sparrow-weaver (*Plocepasser mahali*) conducted by Ngongolo and Mtoka [16] no other study attempted to describe the diversity, abundance, and distribution of avifaunal species across the habitat gradient, and assess the implication of ongoing human activities to the conservation of avifauna species across the Ruvuma Landscape. This gives an opportunity to assess avifauna diversity and distribution in relation to habitat disturbance and how avifauna responded to this habitat destruction. Studying avifauna in Ruvuma landscape will open a room for avitourism activities and conserve from habitat degradation. Therefore, this chapter aimed at presenting the diversity of avifauna species in the Wildlife Management Areas in the Ruvuma Landscape in relation to human activities. It is predicted that

Figure 1.
Map of Ruvuma WMAs showing the location of the sampling sites.

avian species diversity and abundance would be higher in protected habitats inside WMA than in areas dominated by human activities namely farmlands.

In this study we treated the presence of farmlands in WMAs where they are not supposed to be as disturbance, because all WMAs in Tanzania have land use planning. The land use planning in all WMAs provides guidelines by zoning communal land where different activities can be conducted, such cattle grazing, settlements, farming and wildlife conservation area (tourist areas). All plots selected in this study were from wildlife conservation zones where also farms existed. Potential actions for intervention have been highlighted.

2. Methods and materials

2.1 Climate and vegetation types

The rainfall pattern is unimodal spanning from late November to May with a mean annual rainfall of 800–1200 mm in a north–south gradient. The mean annual temperature is 21°C, following the Köppen system [17]. The area consists of extensive miombo woodland, including *Brachystegia* sp., *Julbernardia* sp., *Isoberlinia* sp., *Azelia quanzensis*, *Pterocarpus angolensis*, and rare and threatened plant species such as *Dalbergia melanoxylon*, which forms dense miombo along the hills and rivers [18]. Also, there are seasonal and permanent wetlands (swamps), riverine forests along numerous perennial and seasonal streams. Due to the increasing

anthropogenic activities, the area currently has farmlands and patches of wooded with scattered trees and grazing land.

2.2 Sampling design

Five sites of 200 m x 200 m were established in each WMA, making a total of 25 sites. We selected different habitat types for each of the five sites, namely miombo woodland (open and dense), farmland, swamps, and riverine forest.

2.3 Avifauna survey

Each site was sampled using three complementary methods to maximize the sample size. First, in each habitat type, avifauna counts were carried out using the point transects technique [6, 19]. This method consists of standing at a particular point or walking slowly across the site back and forth several times, to detect cryptic and skulking species in the area. These counts were repeated for 3 days, based on results from our pilot study, and the numbers for each site were averaged. A 20-minute counting period was used at each site, and the starting time (between 6:30 and 10:30 h) was rotated among the sites to reduce bias. Avifauna was identified by both sight and call, and numbers were recorded [20].

Secondly, the transect method was used. Three transects 40 km in length each were established in every WMA using existing roads. The locations of all transects were based on accessibility and were sampled using a vehicle driven at a speed of 20 km/hr. or less that stopped for each individual or group of birds encountered [21]. Two observers sighted and recorded all avifauna on either side of the vehicle and notes on habitat type were also taken [21].

Thirdly, mist-netting was used to the targeted cryptic, understory, and lower canopy avian species. Nets were erected and checked every 15 min in the early morning (between 6:30–10:30 h) and late afternoon (between 16:00–18:00 h). The total number of each species caught, and the associated habitat type was recorded. Each bird was marked with a drop of red permanent spray paints at the base of its toes on the right tarsi for verification, if recaptured, to avoid double counting [22].

2.4 Statistical analysis

The biodiversity indices in different habitats or within these WMAs were obtained following Magurran [23]. This index uses three biodiversity indices including, diversity, richness, and abundance. A non-parametric Kruskal-Wallis test was used to assess whether there were significant differences in mean species abundance among five WMAs, and across each habitat type [24]. Differences in mean bird numbers between habitats in each WMA were tested using Mann-Whitney tests to assess whether the number of species was significantly lower in human-encroached habitat (farmland), i.e., farmland, compared to riverine forest, and dense and open miombo woodland habitats. Statistical tests were computed using the software package PAST [24]. For all these analyses, farmland habitat in this study represented human encroachment into protected areas and was used to compare with other habitat types found in the WMAs. We further calculated the Jaccard similarity index (J_i) between different habitat types to determine the level of similarities in species composition using the formulae [24]:

$$\text{Jaccard similarity coefficient (J); } J = A / (A + B + C) \quad (1)$$

Where A = number of species found in both communities, B = number of species only found in community 1 and C = number of species found in community 2. The equation returns a number between 0 and 1, where a number close to 1 indicates a higher similarity in species composition [23]. We then multiplied J by 100 to obtain a percent, to easily interpret the results.

3. Results

3.1 Avian species diversity, distribution, and richness

A total of 156 avian species representing 18 orders and 61 families were recorded in the five WMAs. The overall avian species Shannon diversity (H') for all the habitat types ranged from 2.28–4.08, except for dense miombo woodland which had $H' = 1.69$ (**Table 1**). Riverine forest habitat had higher species richness ($n = 101$ species), representing almost 45% of the total recorded individuals (**Table 1**). Avian species diversity was highest in riverine forest and lowest in dense miombo woodlands (**Table 1**; **Figure 2**). The Shannon Index of diversity revealed that species evenness for the five habitats surveyed was relatively low ranging from 0.29–0.59 (**Table 1**).

Values bearing different letters within column are significantly different ($p < 0.05$) and values with similar letters within column are not significantly

Habitat type	Number of avian species	Overall abundance	Mean abundance	Shannon diversity (H')	Shannon evenness (E_H)
Dense miombo	14	105	7.50 ± 3.91	1.69	0.39
Farmland	40	580	14.50 ± 5.82	2.46	0.29
Open miombo	98	1338	13.65 ± 2.08	3.9	0.51
Riverine forest	101	759	7.52 ± 0.97	4.08	0.59
Swamp areas	20	188	9.40 ± 3.26	2.28	0.49

Table 1.
Avian species diversity, abundance, and evenness in different habitats of WMAs in Ruvuma landscape ($\pm$ standard error).

Figure 2.
Avian species diversity in different habitats.

different ($p > 0.05$; **Table 2**). Dense miombo woodland, farm and swamp exhibited higher number of birds per point count than in open miombo woodland and riverine forest implying that the avian species were more scattered in open miombo woodlands and riverine forests.

The overall mean abundance of avifauna in the WMAs differed significantly (Kruskal-Wallis test, $\chi^2 = 50.13$, $df = 4$, $P = 0.03$). Kimbanda had the highest mean abundance of species followed by Kisungule (**Figure 3**). There was a significant difference in the mean abundance of avifauna across the five habitats (Kruskal-Wallis test, $\chi^2 = 13.18$, $df = 4$, $P = 0.010$). Mean abundance of species was significantly higher in farmland than in dense miombo (Mann-Whitney tests, $U = 19$, $P < 0.0001$), open miombo woodland ($U = 66.5$, $P < 0.0003$), riverine forest ($U = 157$, $P < 0.019$) and swamps ($U = 93.5$, $P < 0.004$) (**Figure 3**).

The distribution of the 2970 avifauna species recorded in the five habitat types is given in (**Table 1** above; **Figure 4**). Some species were found in more than one habitat type, a total of six species with bronze mannikin (*Lonchura cucullata*) the most abundant (**Figure 5**). Tawny-flanked prinia (*Prinia subflava*), blue-spotted wood dove (*Turtur afer*), common bulbul (*Pycnonotus barbatus*), violet-backed starling (*Cinnyricinclus leucogaster*), and Jameson’s firefinch (*Lagonosticta rhodopareia*) were observed in four habitat types, except swamp habitat (see **Figure 5**; Appendix **Table A1**). Southern cordon-bleu (*Uraeginthus bengalus*) was observed in three habitat types and was the second most abundant species recorded

Habitats	Average bird count
Dense miombo woodland	6.18a
Farm	6.11a
Open miombo woodland	3.71b
Riverine forest	3.45b
swamp	6.48a

Table 2.
Average number of birds per point count in different habitats.

Figure 3.
Avian abundance in different habitats of wildlife management areas in southern Tanzania.

Figure 4. Avifauna species observed foraging in different habitats. Definition of abbreviation used (Demiwo = dense miombo woodland, riverfore = riverine forest, farmland = farmland habitat, opemiwo = open miombo woodland).

Figure 5. Distribution of avian species in different habitats within WMAs of the Ruvuma landscape in southern Tanzania.

during this study (**Figure 5**). Other species including pied crow (*Corvus albus*), brown-headed parrot (*Poicephalus cryptoxanthus*), and red-necked francolin (*Pternistis afer*) were observed in three habitat types (see Appendix **Table A1**) whereas black-faced waxbill (*Estrilda erythronotos*) and African pied wagtail (*Motacilla aguimp*), were observed only in farmland areas.

Cryptic species like African broadbill (*Smithornis capensis*) and red-capped robin-chat (*Cossypha natalensis*) and understory bird species including red-throated twinspot (*Hypargos niveoguttatus*) were observed only in the riverine forest using mist-nets and point count methods (Appendix Table A1). Palearctic migrants including European nightjar (*Caprimulgus europaeus*), European swift (*Apus apus*), and European bee-eater (*Merops apiaster*) were also recorded. Trumpeter hornbill (*Bycanistes bucinator*) is a bird of conservation status that was observed during the study in forest patches.

3.2 Species composition and similarities between different habitat types

We found strong contrast in species composition among habitat types (Table 3). The highest species similarities were between open woodland vs. Riverine forest (41%), Farmland vs. Open woodland (24%) and Farmland vs. Riverine forest (21%)

Habitat types	Dense woodland	Open woodland	Farmland	Riverine	Swamp area
	—	—	—	—	—
Open woodland	11	—	—	—	—
Farmland	15	24	—	—	—
Riverine forest	8	41	21	—	—
Swamp area	0	1	2	5	—

Table 3. Jaccard species composition similarity index (J) between habitat types of the WMAs in Ruvuma landscape, in southern Tanzania. In this table the similarity presented in percentage (%).

Figure 6. Plotted trend line to show species composition similarities between habitat types of the WMAs in Ruvuma landscape, in southern Tanzania. Definition of abbreviation used (dw vs. sw = dense woodland vs. swamp area; dw vs. fm = dense woodland vs. farmland; dw vs. ow = dense woodland vs. open woodland; dw vs. rf = dense woodland vs. riverine forest; fm vs.ow = farmland vs. open woodland; fm vs. rf = farmland vs. riverine forest; fm vs. sw = farmland vs. swamp area; ow vs. rf = open woodland vs. riverine forest; ow vs. sw = open woodland vs. swamp area; rf vs. sw = riverine forest vs. swamp area).

Figure 7. Cluster analysis of different habitat types based on bird species composition (presence/absence). Definition of abbreviation used (Demiwo = dense miombo woodland, riverfore = riverine forest, farmland = farmland habitat, opemiwo = open miombo woodland).

while dense woodland vs. Swamp areas had no similarity in composition (0%), Open woodland vs. Swamp area (1%) and Farmland vs. Swamp area (2%; **Table 3**). The Jaccard similarity indices among various pairs of habitat types compared (**Table 3; Figure 6**).

From the results, avian species adapted to open miombo woodlands and those adapted to riverine forest were very closely related and far from avian species adapted to swamps (**Figure 7**). Avian species adapted to swamps were separated from all other avian species adapted other habitats (**Figure 7**). Indeed, this entails a need for conservation of swamps to avoid local distinction of swamp adapted species.

4. Discussion and conclusion

4.1 Avian species diversity, distribution, and richness

Farmland habitats were observed in all WMAs except in Mbarang’andu where we did not encounter cultivated areas inside the core WMA. Possibly due to the presence of an anti-poaching office established inside WMA by Tanzania Wildlife Management Authority (TAWA, formerly Wildlife Division). In our study, we predicted that there would be higher avian diversity, richness, and abundance in WMAs than in human-modified areas named here as farmlands. We found strong support for this prediction for the species diversity and richness of avifauna but not for abundance. This suggested that the differing occurrence of avifauna species

across given habitats could be attributed to some reasons including food requirement as well as heat tolerance [25].

The richness and diversity imply a variety of taxa that exist in an area, many taxa should, therefore, survive in habitats that have a variety of favorable conditions and resources such as the presence of food, nesting areas, shade and water that might contribute to higher species richness and diversity. Therefore, low species diversity in the farmland might be contributed by the insufficient supply of food as well as insufficient cover for birds to hide against predators, lack of shade to hide from diurnal temperature [12, 26] low food supply compared to forests and woodlands. Suggesting that farmlands have reached maximum disturbance, as in lower farmlands heterogeneous vegetation offer foods and shelter for birds encouraging higher diversity and abundance [8]. Thus the granivores which are largely seed eaters such as the bronze mannikin, southern cordon-bleu, and red-billed quelea were dominant in farmlands than in other habitats because farmlands were rich in seed types vegetation, in line with the findings of others [12, 26]. Furthermore, for similar reasons, the abundance of the granivores species was also higher in open miombo where grassland patches are dominant than in forest areas. Birds that preferred mixed habitat of tree-covered vegetation and open areas chose forest and woodlands but are not water-bound and avoided farmlands such as red-throated twinspot, pygmy kingfisher and red-capped robin-chat, they co-existed in riverine forest and woodland, together with birds that prefer evergreen or lowland forest, dense deciduous thickets, or other dense woodlands such as black-throated wattle-eye and the African broadbill.

4.2 Species composition and similarities between different habitat types

The presence of higher species composition and similarities among habitat types suggests that miombo woodlands harbor unique avifauna species. Some avian species are observed to occur in more than one habitat type indicating that avian species are not habitat specialists. In this study, such patterns were observed; some species existed in more than 4 habitat types suggesting areas visited they provide similar resource abundance, types, and habitat heterogeneity.

Therefore, under no intervention strategies, the Ruvuma Landscape will result in a marked loss of avian richness and diversity. This suggests that measures that will reduce land clearance for agriculture need to be promptly implemented to reduce the ecological impacts on avifauna. Wildlife management areas should involve adjacent communities that are the key stakeholders of the habitats and species biodiversity conservation. Such measures can enhance the resilience of wildlife management areas and complement the goals of community-based conservation measures [27, 28]. Unfortunately, any proposed measures may be challenged by increasing human pressure due to agricultural intensification needs as well as a rapidly changing climate that may be beyond the WMA's management control. Examining the links of these threats to avian biodiversity and addressing such in an urgent manner is likely to abate current human disturbance in the WMAs of Ruvuma region.

Acknowledgements

We thank the District Game Officers of Namtumbo and Tunduru as well as Community Based Conservation Training Centre (CBCTC) staff for their assistance and positive cooperation they have rendered for the success of this project. We thank the Village Game Scouts (VGSs) and all WMA leaders for their guidance during data collection. Furthermore, we recognize the materials and technical support offered by

the Tanzania Wildlife Research Institute (TAWIRI). This study was funded by the WWF Tanzania grant to Geo Network Ltd. based at Dar es Salaam.

Conflict of interest

The authors have not declared any conflict of interests.

Appendix

No.	English name	Species name	Habitat type						Ratio
			Dense miombo woodland	Farmland	Open miombo woodland	Riverine forest	Swamp areas	Grand Total	
1	Bronze mannikin	<i>Spermestes culciata</i>	56	222	56	72	0	406	0.137
2	Southern (Blue- breasted) cordon-bleu	<i>Uraeginthus angolensis</i>	3	85	123	0	0	211	0.071
3	Red-billed quelea	<i>Quelea quelea</i>	0	15	115	0	0	130	0.044
4	Tawny- flanked prinia	<i>Prinia subflava</i>	18	27	38	34	0	117	0.039
5	Common waxbill	<i>Estrilda astrild</i>	0	9	63	20	0	92	0.031
6	Common bulbul	<i>Pycnonotus goiavier</i>	5	1	37	33	0	76	0.026
7	Ring-necked dove	<i>Streptopelia capicola</i>	0	21	51	0	0	72	0.024
8	European bee-eater	<i>Merops apiaster</i>	0	3	51	8	6	68	0.023
9	Violet-backed starling	<i>Cinnyricinclus leucogaster</i>	2	4	45	17	0	68	0.023
10	White-faced whistling- duck	<i>Dendrocygna viduata</i>	0	0	0	0	62	62	0.021
11	Helmeted guineafowl	<i>Numida meleagris</i>	0	0	59	2	0	61	0.021
12	Blue-spotted wood-dove	<i>Turtur afer</i>	5	7	26	11	0	49	0.016
13	African green-pigeon	<i>Treron calvus</i>	0	2	42	4	0	48	0.016
14	Pied crow	<i>Corvus albus</i>	0	30	10	5	0	45	0.015
15	Fork-tailed drongo	<i>Dicrurus adsimilis</i>	0	2	35	5	0	42	0.014
16	Arrow- marked babbler	<i>Turdoides jardineii</i>	0	0	12	26	0	38	0.013
17	Gray-backed (bleating) camaroptera	<i>Camaroptera brevicaudata</i>	0	0	6	32	0	38	0.013

No.	English name	Species name	Habitat type						
			Dense miombo woodland	Farmland	Open miombo woodland	Riverine forest	Swamp areas	Grand Total	Ratio
18	Little greenbul	<i>Eurillas virens</i>	0	0	2	36	0	38	0.013
19	African jacana	<i>Actophilornis africanus</i>	0	0	0	2	33	35	0.012
20	Black-crowned tchagra	<i>Tchagra senegalus</i>	0	4	26	5	0	35	0.012
21	Lesser striped swallow	<i>Cecropis abyssinica</i>	0	13	0	21	0	34	0.011
22	Wire-tailed swallow	<i>Hirundo smithii</i>	0	34	0	0	0	34	0.011
23	Rufous-naped lark	<i>Mirafra africana</i>	0	0	32	0	0	32	0.011
24	Brown-headed parrot	<i>Poicephalus cryptoxanthus</i>	2	0	25	2	0	29	0.010
25	Lesser blue-eared starling	<i>Lamprotornis chloropterus</i>	0	0	25	3	0	28	0.009
26	Black-backed puffback	<i>Dryoscopus cubla</i>	0	0	10	16	0	26	0.009
27	Black-headed oriole	<i>Riolus larvatus</i>	0	1	18	6	0	25	0.008
28	Collared sunbird	<i>Hedydipna collaris</i>	1	0	9	15	0	25	0.008
29	Mosque swallow	<i>Cecropis senegalensis</i>	0	6	12	6	0	24	0.008
30	Pied kingfisher	<i>Ceryle rudis</i>	0	0	0	6	18	24	0.008
31	Mottled spinetail	<i>Telacanthura ussheri</i>	0	0	0	23	0	23	0.008
32	Purple-crested turaco	<i>Gallirex porphyreolophus</i>	0	0	15	8	0	23	0.008
33	Pennant-winged nightjar	<i>Caprimulgus vexillarius</i>	0	0	19	2	0	21	0.007
34	Rattling cisticola	<i>Cisticola chiniana</i>	4	0	0	17	0	21	0.007
35	Tropical boubou	<i>Laniarius aethiopicus</i>	0	0	7	14	0	21	0.007
36	White-headed black chat	<i>Myrmecocichla arnotti</i>	0	0	17	4	0	21	0.007
37	African paradise-flycatcher	<i>Terpsiphone viridis</i>	0	3	12	5	0	20	0.007
38	Gray-headed bush-shrike	<i>Malaconotus blanchoti</i>	0	0	6	13	0	19	0.006
39	African palm-swift	<i>Cypsiurus parvus</i>	0	19	0	0	0	19	0.006
40	Brown-crowned tchagra	<i>Tchagra australis</i>	0	4	7	7	0	18	0.006

No.	English name	Species name	Habitat type						
			Dense miombo woodland	Farmland	Open miombo woodland	Riverine forest	Swamp areas	Grand Total	Ratio
41	Flappet lark	<i>Mirafr rufocinnamomea</i>	0	2	10	6	0	18	0.006
42	Pale-billed hornbill	<i>Lophoceros pallidirostris</i>	0	0	2	16	0	18	0.006
43	Red-throated twinspot	<i>Hypargos niveoguttatus</i>	0	0	0	18	0	18	0.006
44	Gray-headed kingfisher	<i>Halcyon leucocephala</i>	0	0	15	2	0	17	0.006
45	Jameson's frefinch	<i>Lagonosticta rhodopareia</i>	2	2	5	8	0	17	0.006
46	Red-necked francolin	<i>Pternmistis afer</i>	0	6	3	8	0	17	0.006
47	Yellow bishop	<i>Euplectes capensis</i>	0	0	8	9	0	17	0.006
48	African golden oriole	<i>Oriolus auratus</i>	0	2	14	0	0	16	0.005
49	Black-faced waxbill	<i>Estrilda erythronotos</i>	0	15	0	0	0	15	0.005
50	White- rumped swift	<i>Apus caffer</i>	0	3	0	12	0	15	0.005
51	Yellow- breasted apalis	<i>Apalis flavida</i>	0	0	3	12	0	15	0.005
52	Black- throated wattle-eye	<i>Platysteira peltata</i>	0	0	0	14	0	14	0.005
53	African firefinch	<i>Lagonosticta rubricata</i>	0	6	4	3	0	13	0.004
54	Green woodhoopoe	<i>Phoeniculus purpureus</i>	0	0	13	0	0	13	0.004
55	Spotted flycatcher	<i>Muscicapa striata</i>	0	0	12	1	0	13	0.004
56	Orange- breasted bush-shrike	<i>Chlorophoneus sulfureopectus</i>	0	0	8	5	0	13	0.004
57	White-backed duck	<i>Thalassornis leuconotus</i>	0	0	0	0	12	12	0.004
58	White- browed sparrow- weaver	<i>Plocepasser mahali</i>	0	0	12	0	0	12	0.004
59	Yellow- fronted canary	<i>Crithagra mozambica</i>	0	0	12	0	0	12	0.004
60	African darter	<i>Anhinga rufa</i>	0	0	0	0	11	11	0.004
61	Kurrichane thrush	<i>Turdus libonyana</i>	0	0	9	2	0	11	0.004
62	African gray hornbill	<i>Lophoceros nasutus</i>	0	2	3	5	0	10	0.003
63	Böhm's spinetail	<i>Neafrapus boehmi</i>	0	0	0	10	0	10	0.003

No.	English name	Species name	Habitat type						
			Dense miombo woodland	Farmland	Open miombo woodland	Riverine forest	Swamp areas	Grand Total	Ratio
64	Common squacco heron	<i>Ardeola ralloides</i>	0	0	0	0	10	10	0.003
65	Coqui francolin	<i>Peliperdix coqui</i>	0	0	10	0	0	10	0.003
66	Shelley's sunbird	<i>Cinnyris shelleyi</i>	0	0	3	7	0	10	0.003
67	Reichenow's woodpecker	<i>Campethera scriptoricauda</i>	1	0	9	0	0	10	0.003
68	African broadbill	<i>Smithornis capensis</i>	0	0	0	9	0	9	0.003
69	Black crake	<i>Zapornia flavirostra</i>	0	0	0	2	7	9	0.003
70	Green-capped eremomela	<i>Eremomela scotops</i>	0	0	6	3	0	9	0.003
71	Striped kingfisher	<i>Halcyon chelicuti</i>	0	0	7	2	0	9	0.003
72	Little bee-eater	<i>Merops pusillus</i>	0	0	6	2	0	8	0.003
73	Little swift	<i>Apus affinis</i>	0	8	0	0	0	8	0.003
74	Pied wagtail	<i>Motacilla aguimp</i>	0	8	0	0	0	8	0.003
75	Senegal lapwing	<i>Vanellus lugubris</i>	0	0	8	0	0	8	0.003
76	Amethyst sunbird	<i>Chalcomitra amethystina</i>	0	0	0	7	0	7	0.002
77	Greater honeyguide	<i>Indicator indicator</i>	0	0	7	0	0	7	0.002
78	Racket-tailed roller	<i>Coracias spatulatus</i>	0	1	6	0	0	7	0.002
79	Red-faced cisticola	<i>Cisticola erythrops</i>	0	0	0	4	3	7	0.002
80	Rufous-bellied tit	<i>Melaniparus rufiventris</i>	0	0	5	2	0	7	0.002
81	Broad-billed roller	<i>Eurystomus glaucurus</i>	0	0	5	1	0	6	0.002
82	Brown-hooded kingfisher	<i>Halcyon albiventris</i>	0	0	6	0	0	6	0.002
83	Dark chanting-goshawk	<i>Melierax metabates</i>	0	1	4	1	0	6	0.002
84	Eastern bearded scrub-robin	<i>Tychaemon quadrivirgata</i>	0	0	2	4	0	6	0.002
85	Great white egret	<i>Ardea alba</i>	0	0	0	0	6	6	0.002
86	Southern ground-hornbill	<i>Bucorvus leadbeateri</i>	0	0	6	0	0	6	0.002
87	Livingstone's turaco	<i>Tauraco livingstonii</i>	0	0	6	0	0	6	0.002

No.	English name	Species name	Habitat type						
			Dense miombo woodland	Farmland	Open miombo woodland	Riverine forest	Swamp areas	Grand Total	Ratio
88	Red-cheeked cordon-bleu	<i>Uraeginthus bengalus</i>	0	0	0	6	0	6	0.002
89	Southern gray-headed sparrow	<i>Passer diffusus</i>	0	0	6	0	0	6	0.002
90	Swallow- tailed bee- eater	<i>Merops hirundineus</i>	0	0	6	0	0	6	0.002
91	Trumpeter hornbill	<i>Bycanistes bucinator</i>	0	0	0	6	0	6	0.002
92	White-crested helmetshrike	<i>Prionops plumatus</i>	0	0	6	0	0	6	0.002
93	Willow warbler	<i>Phylloscopus trochilus</i>	0	0	6	0	0	6	0.002
94	Common hoopoe	<i>Upupa epops</i>	0	0	0	5	0	5	0.002
95	Black cuckoo	<i>Cuculus clamosus</i>	0	0	0	5	0	5	0.002
96	Black kite	<i>Milvus migrans</i>	0	2	3	0	0	5	0.002
97	Common sandpiper	<i>Actitis hypoleucos</i>	0	0	0	3	2	5	0.002
98	Golden-tailed woodpecker	<i>Campethera abingoni</i>	0	0	4	1	0	5	0.002
99	Little sparrowhawk	<i>Accipiter minullus</i>	0	0	4	1	0	5	0.002
100	Pale (East coast) batis	<i>Batis soror</i>	0	0	2	3	0	5	0.002
101	Pygmy kingfisher	<i>Ispidina picta</i>	0	0	0	3	2	5	0.002
102	Red-chested cuckoo	<i>Cuculus solitarius</i>	0	0	0	5	0	5	0.002
103	Miombo wren warbler	<i>Calamonastes stierlingi</i>	0	0	5	0	0	5	0.002
104	Wattled lapwing	<i>Vanellus senegallus</i>	0	0	0	3	2	5	0.002
105	White-bellied sunbird	<i>Cinnyris talatala</i>	0	0	3	2	0	5	0.002
106	White- breasted cuckoo-shrike	<i>Cebilepyris pectoralis</i>	0	0	5	0	0	5	0.002
107	Yellow-bellied greenbul	<i>Chlorocichla flaviventris</i>	0	0	2	3	0	5	0.002
108	Cardinal woodpecker	<i>Dendropicos fuscescens</i>	1	0	3	0	0	4	0.001
109	African pipit	<i>Anthus richardi</i>	0	0	3	0	2	5	0.002
110	Hamerkop	<i>Scopus umbretta</i>	0	0	0	4	0	4	0.001
111	Lilac-breasted roller	<i>Coracias caudatus</i>	0	0	2	2	0	4	0.001
112	Pearl-spotted owlet	<i>Glaucidium perlatum</i>	0	0	4	0	0	4	0.001

No.	English name	Species name	Habitat type						
			Dense miombo woodland	Farmland	Open miombo woodland	Riverine forest	Swamp areas	Grand Total	Ratio
113	Red-capped robin-chat	<i>Cossypha natalensis</i>	0	0	0	4	0	4	0.001
114	White-browed coucal	<i>Centropus superciliosus</i>	0	0	2	2	0	4	0.001
115	White-browed robin-chat	<i>Cossypha heuglini</i>	0	0	0	4	0	4	0.001
116	Black cuckoo-shrike	<i>Campephaga flava</i>	0	0	1	2	0	3	0.001
117	Böhm's bee-eater	<i>Merops boehmi</i>	0	0	0	3	0	3	0.001
118	Brubru	<i>Nilaus afer</i>	0	0	3	0	0	3	0.001
119	Cabanis's bunting	<i>Emberiza cabanisi</i>	0	2	1	0	0	3	0.001
120	Crested barbet	<i>Trachyphonus vaillantii</i>	0	0	1	2	0	3	0.001
121	Crowned hornbill	<i>Lophoceros alboterminatus</i>	0	3	0	0	0	3	0.001
122	European swift	<i>Apus apus</i>	0	0	0	3	0	3	0.001
123	African fish eagle	<i>Haliaeetus vocifer</i>	0	0	0	1	2	3	0.001
124	Hadada ibis	<i>Bostrychia hagedash</i>	0	0	0	0	3	3	0.001
125	Harlequin quail	<i>Coturnix delegorguei</i>	0	0	0	3	0	3	0.001
126	Namaqua dove	<i>Oena capensis</i>	3	0	0	0	0	3	0.001
127	Speckle-throated woodpecker	<i>Campethera scriptoricauda</i>	0	0	3	0	0	3	0.001
128	Parasitic weaver	<i>Anomalospiza imberbis</i>	0	0	3	0	0	3	0.001
129	Red-fronted tinkerbird	<i>Pogoniulus pusillus</i>	0	0	0	3	0	3	0.001
130	Red-headed weaver	<i>Anaplectes rubriceps</i>	0	0	3	0	0	3	0.001
131	Speckled mousebird	<i>Colius striatus</i>	0	0	0	3	0	3	0.001
132	Stripe-breasted seedeater	<i>Crithagra striatipectus</i>	0	0	3	0	0	3	0.001
133	White-browed scrub-robin	<i>Cercotrichas leucophrys</i>	0	0	1	2	0	3	0.001
134	Wood sandpiper	<i>Tringa glareola</i>	0	0	0	0	3	3	0.001
135	Black-headed heron	<i>Ardea melanocephala</i>	0	1	0	1	0	2	0.001

No.	English name	Species name	Habitat type						
			Dense miombo woodland	Farmland	Open miombo woodland	Riverine forest	Swamp areas	Grand Total	Ratio
136	Black-winged stilt	<i>Himantopus himantopus</i>	0	0	0	2	0	2	0.001
137	Brimstone canary	<i>Crithagra sulphurata</i>	0	0	2	0	0	2	0.001
138	Egyptian goose	<i>Alopochen aegyptiaca</i>	0	0	0	0	2	2	0.001
139	Fiscal shrike	<i>Lanius collaris</i>	0	2	0	0	0	2	0.001
140	Golden- breasted bunting	<i>Emberiza flaviventris</i>	2	0	0	0	0	2	0.001
141	Retz's helmet shrike	<i>Prionops retzii</i>	0	0	0	2	0	2	0.001
142	Scarlet- chested sunbird	<i>Chalcomitra senegalensis</i>	0	0	0	2	0	2	0.001
143	Tambourine dove	<i>Turtur tympanistria</i>	0	0	0	2	0	2	0.001
144	African barred owl	<i>Glaucidium capense</i>	0	0	1	0	0	1	0.000
145	Piping cisticola	<i>Cisticola fulvicapilla</i>	0	0	3	0	0	3	0.001
146	Red-eyed dove	<i>Streptopelia semitorquata</i>	0	0	0	3	0	3	0.001
147	Beautiful sunbird	<i>Cinnyris pulchellus</i>	0	1	0	0	0	1	0.000
148	Black coucal	<i>Centropus grillii</i>	0	0	0	1	0	1	0.000
149	Brown snake- eagle	<i>Circaetus cinereus</i>	0	0	0	1	0	1	0.000
150	European nightjar	<i>Caprimulgus europaeus</i>	0	0	1	0	0	1	0.000
151	Gray heron	<i>Ardea cinerea</i>	0	0	0	0	1	1	0.000
152	Olive sunbird	<i>Cyanomitra olivacea</i>	0	0	0	1	0	1	0.000
153	Saddlebill	<i>Ephippiorhynchus senegalensis</i>	0	0	0	0	1	1	0.000
154	Spectacled weaver	<i>Ploceus ocularis</i>	0	0	1	0	0	1	0.000
155	Spotted creeper	<i>Salpornis salvadori</i>	0	0	0	1	0	1	0.000
156	Woodland kingfisher	<i>Halcyon senegalensis</i>	0	0	0	1	0	1	0.000
Grand Total			105	580	1338	759	188	2970	

Table A1.
List of avifauna species observed in different habitats of WMAs in Ruvuma.

IntechOpen

Author details

Ally K. Nkwabi^{1,2*}, John K. Bukombe², Hamza K. Kija², Steven D. Liseki²,
Sood A. Ndimuligo³ and Pius Y. Kavana^{1,2}

1 Mahale-Gombe Wildlife Research Centre, Kigoma, Tanzania

2 Tanzania Wildlife Research Institute, Arusha, Tanzania

3 Department of Biosciences, Centre for Ecological and Evolutionary Synthesis
(CEES), University of Oslo, Norway

*Address all correspondence to: ally.nkwabi@tawiri.or.tz

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Frost, P., *The ecology of miombo woodlands*, in *The miombo in transition: Woodlands and welfare in Africa*, C. Campbell, Editor. 1996, Centre for International Forestry Research: Bogor, Indonesia. p. 11-57.
- [2] Mittermeier, R.A., et al., *Global biodiversity conservation: the critical role of hotspots*, in *Biodiversity hotspots*, F.E. Zachos and J.C. Habel, Editors. 2011, Springer: Verlag Berlin Heidelberg. p. 3–22.
- [3] Munishi, P., et al., *The role of the Miombo Woodlands of the Southern Highlands of Tanzania as carbon sinks*. *Journal of Ecology and The Natural Environment*, 2010. **2**(12): p. 261–269.
- [4] Sawe, T.C., P.K. Munishi, and S.M. Maliondo, *Woodlands degradation in the Southern Highlands, Miombo of Tanzania: Implications on conservation and carbon stocks*. *International Journal of Biodiversity and Conservation*, 2014. **6**(3): p. 230–237.
- [5] Daniels, R.J., N.V. Joshi, and M. Gadgil, *On the relationship between bird and woody plant species diversity in the Uttara Kannada district of south India*. *Proceedings of the National Academy of Sciences*, 1992. **89**(12): p. 5311–5315.
- [6] Pomeroy, D. and C. Dranzoa, *Methods of studying the distribution, diversity and abundance of birds in East Africa-some quantitative approaches*. *African Journal of Ecology*, 1997. **35**(2): p. 110–123.
- [7] Oindo, B.O., A.K. Skidmore, and P. De Salvo, *Mapping habitat and biological diversity in the Maasai Mara ecosystem*. *International journal of remote sensing*, 2003. **24**(5): p. 1053–1069.
- [8] Nyirenda, V.R., et al., *Avian occupancy along multiple land use type gradients in Miombo dominated landscapes of the Copper Belt Province, Zambia*. *Journal of Biodiversity Management and Forestry*, 2017. **6**: p. 1.
- [9] Lupala, Z., et al., *The land use and cover change in miombo woodlands under community based forest management and its implication to climate change mitigation: a case of southern highlands of Tanzania*. *International Journal of Forestry Research*, 2015. **2015**: p. Article ID 459102.
- [10] Nduwamungu, J., et al., *Recent land cover and use changes in Miombo woodlands of eastern Tanzania*. *Tanzania Journal of Forestry and Nature Conservation*, 2008. **78**: p. No. 1.
- [11] Oindo, B.O., R.A. De By, and A.K. Skidmore, *Environmental factors influencing bird species diversity in Kenya*. *African Journal of Ecology*, 2001. **39**(3): p. 295–302.
- [12] Nyirenda, V.R., et al., *Nest sites selection by sympatric cavity-nesting birds in miombo woodlands*. *Koedoe*, 2016. **58**(1): p. 1–10.
- [13] MNRT, *Vegetation study on the Biodiversity conservation values and Management strategies of the Selous-Niassa Wildlife Corridor*. 2006.
- [14] Luoga, E., E. Witkowski, and K. Balkwill, *Land cover and use changes in relation to the institutional framework and tenure of land and resources in eastern Tanzania miombo woodlands*. *Environment, Development and sustainability*, 2005. **7**(1): p. 71–93.
- [15] Handavu, F., P.W. Chirwa, and S. Syampungani, *Socio-economic factors influencing land-use and land-cover changes in the miombo woodlands of the Copperbelt province in Zambia*. *Forest policy and economics*, 2019. **100**: p. 75–94.pt
- [16] Ngongolo, K. and S. Mtoka, *Abundance, nesting and habitat for white-browed sparrow-weaver (Plocepasser*

mahali) in northeastern Selous Game Reserve, Tanzania. International Journal of Fauna and Biological Studies, 2013. 1 (1): p. 63–67.

[17] Köppen, W., *Outline of climate science*. 1931, Berlin: Walter de Gruyter.

[18] Baldus, R.D. and R. Hahn, *The Selous-Niassa Wildlife Corridor in Tanzania: Biodiversity Conservation from the Grassroots. Practical Experiences and Lessons from Integrating Local Communities into Transboundary Natural Resources Management*, Joint publication of FAO and CIC. 2009, Hungary: Budapest.

[19] Pomeroy, D.E., *Counting birds*. 1992, Nairobi: African Wildlife Foundation.

[20] Jones, M.J., M.D. Linsley, and S.J. Marsden, *Population sizes, status and habitat associations of the restricted-range bird species of Sumba, Indonesia*. Bird Conservation International, 1995. 5(1): p. 21–52.

[21] Mmassy, E.C., et al., *Kori bustard (Ardeotis kori struthiunculus) occurrence in the Serengeti grass plains, northern Tanzania*. African Journal of Ecology, 2017. 55(3): p. 298–304.

[22] Pierce, D.O., I. Flux, and M. Potter. *Kiwi (Apteryx spp.) Best Practice Manual: Abgerufen von http://www.kiwiforkiwi.org/wp-content/uploads/2012/09/Kiwi_Best_Practice_Manual_Sep03.pdf*. 2003 20 March 2018].

[23] Magurran, A.E., *Measuring Biological Diversity*. 2004, Oxford: Blackwell.

[24] Hammer, Ø., D.A.T. Harper, and P. D. Ryan, *PAST: Paleontological Statistics Software Package for Education and Data Analysis Palaeontol. Electronica*, 2001. 4: p. 1–9.

[25] Morimoto, T., et al., *Can surrounding land cover influence the avifauna in urban/suburban woodlands*

in Japan? Landscape and Urban Planning, 2006. 75(1–2): p. 143–154.

[26] Girma, Z., G. Mengesha, and T. Asfaw, *Diversity, relative abundance and distribution of avian fauna in and around wondo genet forest, South-central Ethiopia*. Research Journal of Forestry, 2017. 11(1): p. 1–12.

[27] Sinclair, A. and A.E. Byrom, *Understanding ecosystem dynamics for conservation of biota*. Journal of Animal Ecology, 2006. 75(1): p. 64–79.

[28] Songorwa, A.N., *Community-based wildlife management (CWM) in Tanzania: Are the communities interested?* World development, 1999. 27(12): p. 2061–2079.