We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Gender Disparities in Bladder Cancer

Yingsheng Zhang, Dan Theodorescu and Xue Li

Abstract

Biological sex is an independent risk factor of cancer. Men are three to five times more likely than women to develop bladder cancer even when known risk factors are taken into consideration. Development of sex in mammals is often viewed as a two-step process. The first step is sex determination, of which the XX and XY sex chromosome complements trigger gonad differentiation to form the ovary and testis, respectively. After that, sex hormones secreted by gonads initiate sexually dimorphic differentiation of nongonadal tissues. However, this model has been challenged by recent findings revealing an independent contribution of sex chromosomes to sexual dimorphism. In this chapter, we discuss how the sex chromosomes and sex hormones together cause gender disparities in bladder cancer. We propose a concept of epigenetic sex – epigenetic differences between males and females – and suggest that the sex epigenome is a previously unknown biasing factor contributing to gender disparities in bladder cancer.

Keywords: Bladder Cancer, Gender Disparities, Sex Hormones, Sex Chromosomes, Sex Epigenome, KDM6A, PRC2, COMPASS

1. Introduction

Bladder cancer (BC) originates primarily from the urothelium – the inner lining of bladder lumen. Men are disproportionately affected by the disease. Males are 3–5 times more likely than females to BC [1]. This difference persists even after adjusting other known risk factors [2–5], suggesting that male sex is an independent risk factor of BC.

Typical males have one copy each of the X and Y chromosomes (XY) while females have two copies of the X chromosome (XX). For XY individuals, sex-determining region Y (SRY) gene on the Y chromosome triggers gonadal differentiation to form testes, which secrete androgens and promote male primary and secondary sex characteristics. Females with XX chromosome complement have ovarian development and estrogen secretion leading to female primary and secondary sex characteristics [6].

Strong evidence exists that androgens acting through androgen receptor, promote bladder tumorigenesis [7]. Female gonadal hormones acting through estrogen receptors also influence BC risk albeit playing a minor role when compared to androgens [8]. While sex hormones clearly play important roles in gender disparities in BC, potential role of the sex chromosomes is not nearly as apparent. Because the sex chromosomes (*i.e.*, XX vs. XY) are coupled with the gonadal hormones (*e.g.*, estrogens vs. androgens), it is exceedingly difficult to ascertain independent effects of the sex chromosomes or the gonadal hormones [6]. Creative tools are needed

to overcome the challenges of studying sex differences in physiology and disease [9]. Here, we review how the sex hormones and chromosomes function together to cause gender disparities in BC and, furthermore, propose a novel concept of epigenome sex.

2. The impact of biological sex on bladder cancer development

2.1 The role of androgen

A retrospective study revealed that male patients who received α -reductase inhibitors before the diagnosis of BC had better survival and was positively correlated with duration of administration [10]. Similarly, prostate cancer patients who received androgen deprivation therapy (ADT) had a lower incidence of BC compared with patients that did not [11]. Paradoxically, a majority of expression analyses of androgen receptor (AR) in BC patients showed a negative correlation between AR expression and the aggressiveness of BC [12–17]. Offering a possible explanation underlying this observation is our finding that reduced AR expression may lead to upregulation of cancer stem cell related genes such as CD44 [18]. Clearly the role of androgens and AR in BC is complex and likely in molecular and cellular context-dependent manner.

Animal models of BC support the role of androgens/AR in modulating bladder tumorigenesis. Male mice were much more vulnerable than female mice to BC induced by N-butyl-N-(4-hydroxybutyl) nitrosamine (BBN), a bladder-specific carcinogen [19]. This sex difference was blunted by castration. Similarly, Okajima and colleagues have observed male-biased responses to BBN-induced BC in rats. Moreover, administration of Diethylstilbestrol (DES), a nonsteroidal estrogen, suppressed bladder carcinogenesis in male rats. In contrast, testosterone supplementation increased the incidence of BC in female rats [20]. Miyamoto *et al.* have also found that AR knockout (KO) mice, both males and females, were completely protected from BBN-induced BC [21]. Intriguingly, 25% AR KO male mice supplemented with dihydrotestosterone (DHT) had BC after BBN treatment, and 50% of castrated male mice still developed into BC [21]. These data suggest that androgens have AR-independent function in promoting BBN-induced BC, and AR may also function in an androgen-independent manner. In addition, they showed that androgen deprivation or androgen/AR disruption blunted growth of AR-positive human BC cells both *in vitro* and *in vivo* [21]. The genetic evidence of oncogenic role of androgen/AR signaling in bladder carcinogenesis was further confirmed in the subsequent studies. Hsu and colleagues revealed that urothelium-specific conditional KO of AR reduced the risk of BBN-induced BC in male mice. They further showed that AR expression was inversely associated with the level of p53-PCNA-DNA damage pathway, implicating a possible downstream event of AR oncogenic signaling [22]. Johnson *et al.* showed that conditional overexpression of human AR in mouse urothelium was sufficient to promote BBN-induced BC in both sexes and the phenotype in males could be alleviated by castration [23]. Surprisingly, the study also suggested that AR did not function through known molecular pathways which have been previously implicated, including p53 [22], Wnt/ β -catenin [24, 25], and CD24 [26–28]. Therefore, much more work is needed to define the precise role of AR in bladder tumorigenesis.

2.2 The role of estrogens and their receptors

Postmenopausal women had a higher risk of BC and early menopause enhanced this risk, suggesting that female sex hormones protect women from BC

development [29–31]. Classical estrogen receptors (ERs) include estrogen receptor α (ER α) and estrogen receptor β (ER β) [32]. Clinically, inconsistent results existed between the expression of ER α and ER β and the grades and stages of BC patients although more reports supported that expression of ER α favors a better while ER β is associated with a worse prognosis [14, 32–38]. Experimentally, both whole body ER α KO or urothelium-specific ER α KO increased the incidence of BBN-induced BC in female mice. Disruption of ER α decreased the expression of Inositol polyphosphate-4-phosphatase, type II (INPP4B), resulting in a higher activation of AKT [39]. On the contrary, whole body deletion of ER β impeded BBN-induced BC in female mice [40]. Moreover, knockdown (KD) of ER β suppressed transformation of normal bladder cells and growth of BC cells partly through reducing expression of mini-chromosome maintenance complex component 5 (MCM5) because reintroduction of MCM5 into BC cells blunted ER β KD phenotype [40]. Interestingly, tamoxifen treatment conferred a chemoprevention in female mice against BBN-induced BC [41]. Since tamoxifen is a selective estrogen-receptor modulator with mixed estrogenic and antiestrogenic activity depending on targeted tissues, it would be interesting to see which ERs, ER α or ER β , is activated or inhibited and whether any of these receptors plays a more dominant role in the BBN-induced bladder carcinogenesis.

2.3 The role of sex chromosomes in driving gender disparities

A potential role of the sex chromosomes in gender disparities in BC was implicated initially by cancer epidemiological findings of Turner and Klinefelter patients. Turner syndrome is a genetic disorder of female XO patients who lost one copy of the X chromosome. Conversely, Klinefelter syndrome has two or more copies of the X chromosome among the affected male patients. Turner patients displayed an increased risk of BC [42]; and Klinefelter patients had an overall reduced rate of solid tumors [43]. These observations suggest that an extra copy of X chromosome is tightly associated with low BC risk in both sexes. However, because the sex chromosomes are tightly coupled with their respective sex/gonadal hormones, the confounding effect of sex chromosomes cannot be excluded. As a result, the sex hormone-independent roles of the sex chromosomes have largely been overlooked.

To overcome the aforementioned challenges of studying independent roles of the sex-biasing factors, De Vries *et al.* developed “four-core genotype (FCG)” mouse model to uncouple the link between the sex chromosomes and their corresponding gonadal types [44]. In this model, *Sry* gene is “transferred” from the Y chromosome to an autosome. As a result, the FCG mouse model has four instead of two sex types: 1. XY male with testes (XYM); 2. XX male with testes (XXM); 3. XX female with ovaries (XXF); and 4. XY female with ovaries (XYF). Independent effects of the sex hormones can be evaluated by comparing XXF and XXM mice or XYM and XXM mice without the confounding issue of the sex/gonadal hormones [45]. Similarly, in the comparisons of XXF vs. XYF or XYM vs. XFM, one can evaluate independent effects of the sex chromosomes.

By taking advantage of the FCG mice, we showed that, independent of the sex hormones, the sex chromosomes had a sex-biasing effect on BC development [46]. We further showed that regardless of gonadal sex XY mice had 2.55 times of higher chance of developing BBN-induced BC than XX mice, demonstrating an independent role of the sex chromosomes. This study has also confirmed the sex-biasing role of androgens and further revealed that having testes was 4.71 times more likely than having ovaries to develop BC. More strikingly, wild type male mice with the XY chromosome complement and testes were 12.39 times more likely than wild type female mice with the XX chromosome complement and ovaries. This is unexpected

because it is close to the product of 2.55 and 4.71 instead of the sum. Such finding suggests that both the sex chromosomes and the sex hormones have independent and dependent sex-biasing effects on BC. Moreover, the sex chromosomes interact with the sex hormones to amplify the difference (**Figure 1**). The underlying mechanism of interaction between these sex-biasing factors is yet to be defined.

Because the Y chromosome is frequently lost in BC cells and its loss has been associated with a higher cancer risk [47–49], it is less likely that the Y chromosome explains the male dominance in BC. A more reasonable possibility is that copy number difference of the X chromosome may render females better protected than males. To understand the potential tumor suppressing role of the X chromosome, we have examined the X chromosome-linked genes that escape X chromosome inactivation (XCI) (**Figure 1a**) [50], hence, genes that are expressed in higher levels in XX than in XY urothelial cells. By comparing gene expression levels in the bladder urothelium of FCG mice, we have identified Lysine Demethylase 6A (KDM6A)

Figure 1. Major contributing factors of sex differences in bladder cancer. a) Schematic diagram of human sex chromosome complement of XX females and XY males. Genes that are reportedly escaped X-chromosome inactivation (XCI) in both mouse and human are indicated. PAR, Pseudo Autosomal Regions. b) Somatic DNA mutation rate of the candidate XCI escapees in bladder cancer. c) A proposed model that interactions among the sex chromosomes, sex hormones, and sex epigenome amplify male and female differences in bladder cancer.

as a top candidate of X-linked tumor suppressors [46]. *KDM6A* encodes a histone demethylase to remove methyl group from methylated histone H3 at lysine K27 to allow gene transcription. Somatic loss-of-function mutations of human *KDM6A* are tightly associated with BC, suggesting a tumor suppressive function in humans [51–54]. Interestingly, *KDM6A* mutations have been shown to be more common in female patients with non-muscle invasive BC (NMIBC) [54]. We showed that after conditional *Kdm6a* KO in the urothelium of female mice, susceptibility to BBN-induced BC was significantly increased. Similarly, down-regulation or mutation of *KDM6A* was tightly linked to advanced disease stages and poor outcomes among female but not male BC patients [46]. Urothelium-specific *Kdm6a* KO did not exhibit apparent phenotype in male mice. A lack of phenotype in males is potentially due to the compensatory effect of a paralog gene *UTY* on the Y chromosome, albeit that *UTY* lacks any detectable demethylase activity *in vivo* [55–58]. In addition to *KDM6A*, other X-linked genes have also been suggested to contribute to sex differences in cancer [59]. While these candidate genes were expressed in higher levels in females than in males, their expression was not associated with disease outcomes of BC (**Figure 1b**) [46]. Collectively, these findings suggest that *KDM6A* is a prototypical sex-biasing tumor suppressor in BC.

2.4 The sex epigenome

Through the intrinsic histone demethylase activity, *KDM6A* regulates downstream gene transcription by antagonizing Polycomb Repressive Complex 2 (PRC2)-dependent epigenetic gene silencing program [60–63]. Specifically, *KDM6A* catalyzes removal of the methyl groups from histone H3 lysine 27 trimethylation (H3K27me₃), making H3K27 available for acetylation (H3K27ac). H3K27me₃ and H3K27ac are closely associated with transcription repression and activation, respectively. In a demethylase-independent manner, *KDM6A* functions in the COMPASS-like protein complex that harbors MLL3/KMT2C and MLL4/KMT2D lysine methyltransferases [64]. KMT2C and KMT2D catalyze formation of H3K4 monomethylation (H3K4me₁), which is tightly linked to active transcription enhancers [65–67]. COMPASS and PRC2 display an antagonistic relationship in regulating downstream gene expression [68, 69]. Therefore, *KDM6A* plays a central role in shaping the epigenetic landscape by modulating the PRC2-dependent gene silencing and the COMPASS-dependent gene activation. Nearly 74% of NMIBC and 23% MIBC patients have mutations in *KDM6A* [51, 54]. About 18% and 28% of human BC patients have somatic mutations in *KMT2C* and *KMT2D*, respectively [51]. The sex-biased expression of *KDM6A* suggests that there is an epigenetic difference in bladder urothelium between sexes, hence the sex epigenome. It is conceivable that *KDM6A* plays a critical role in creating an epigenetic barrier to prevent malignant transformation. Females with more *KDM6A* may have a tougher barrier than males to overcome, thereby are less likely to develop BC.

Whole-genome transcriptome analysis of *Kdm6a* knockout urothelium indicates that the p53 tumor suppressor pathway is among the top affected pathways [46]. The canonical p53 downstream gene targets, *Cdkn1a* and *Perp*, which induce cell cycle arrest and apoptosis, respectively, are down-regulated significantly in *KDM6A* knockouts. UM-UC-13, a human BC cell line, lacks functional *KDM6A* due to genomic mutation [70]. Expression of wild type *KDM6A* induces both *CDKN1A* and *PERP* in UM-UC-13 cells while expression of the catalytically-dead form of *KDM6A* only induces *PERP* but not *CDKN1A*, suggesting that *KDM6A* regulates *PERP* gene expression in a demethylase-independent manner. While persistent expression of the enzymatically-dead form of *KDM6A* is required to suppress UM-UC-13 cell proliferation, a transient expression of *KDM6A* is sufficient to

achieve the same result, implying that the demethylase activity of KDM6A results in a lasting protective activity or an epigenetic memory of tumor suppression [46]. In supporting this notion, *Ler et al.* revealed that *KDM6A* mutant BC cells are more vulnerable to the pharmacological inhibition of PRC2 [52]. While the precise mechanism of *KDM6A*-dependent tumor suppression remains to be fully elucidated, we suspect that systematic profiling of the sex epigenome would shed new light on the gender disparities in BC (**Figure 1c**).

3. Conclusion

A new concept of sex epigenome begins to emerge. In addition to gonadal hormones, the copy number difference of the X chromosome between males and females contributes to sex differences in BC - an extra copy of the X chromosome confers a better protection of females. Moreover, there is a cooperative interaction between the sex hormones and chromosomes during BC development. The tumor suppressing effect of the X chromosome is largely mediated by *KDM6A*-dependent epigenetic program, or the sex epigenome. The sex chromosome, sex chromosome, and sex epigenome collectively contribute to sex differences in BC.

Acknowledgements

This work was supported by the NIH/National Cancer Institute (1R01DK110477, 1R01HL136921, and 5R21CA249701 to X.L.).

Conflict of interest

The authors declare no conflict of interest.

Notes/thanks/other declarations

We thank BioRender (<https://biorender.com/>) for providing the platform for plotting the **Figure 1c**.

IntechOpen

Author details

Yingsheng Zhang^{1,2}, Dan Theodorescu^{2,3} and Xue Li^{1,2*}

1 Department of Medicine, Cedars-Sinai Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, CA, USA

2 Department of Biological Sciences, Cedars-Sinai Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, CA, USA

3 Department of Surgery, Cedars-Sinai Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, CA, USA

*Address all correspondence to: sean.li@cshc.org

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Cumberbatch MGK, Jubber I, Black PC, Esperto F, Figueroa JD, Kamat AM, et al. Epidemiology of Bladder Cancer: A Systematic Review and Contemporary Update of Risk Factors in 2018. Vol. 74, European Urology. Elsevier B.V.; 2018. p. 784-95.
- [2] Hartge P, Harvey EB, Linehan WM, Silverman DT, Sullivan JW, Hoover RN, et al. Unexplained Excess Risk of Bladder Cancer in Men. JNCI J Natl Cancer Inst [Internet]. 1990 Oct 17 [cited 2021 Mar 9];82(20):1636-40. Available from: <https://academic.oup.com/jnci/article-lookup/doi/10.1093/jnci/82.20.1636>
- [3] Hemelt M, Yamamoto H, Cheng KK, Zeegers MPA. The effect of smoking on the male excess of bladder cancer: A meta-analysis and geographical analyses. Int J Cancer [Internet]. 2009 Jan 15 [cited 2021 Mar 9];124(2):412-9. Available from: <http://doi.wiley.com/10.1002/ijc.23856>
- [4] Krabbe LM, Svatek RS, Shariat SF, Messing E, Lotan Y. Bladder cancer risk: Use of the PLCO and NLST to identify a suitable screening cohort. Urol Oncol Semin Orig Investig. 2015 Feb 1;33(2):65.e19-65.e25.
- [5] Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. CA Cancer J Clin [Internet]. 2020 Jan 8 [cited 2021 Mar 9];70(1):7-30. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.3322/caac.21590>
- [6] Arnold AP. A general theory of sexual differentiation. J Neurosci Res [Internet]. 2017 Jan 2 [cited 2021 Mar 9];95(1-2):291-300. Available from: <http://doi.wiley.com/10.1002/jnr.23884>
- [7] Li P, Chen J, Miyamoto H. Androgen Receptor Signaling in Bladder Cancer. Cancers (Basel) [Internet]. 2017 Feb 22 [cited 2021 Mar 19];9(12):20. Available from: <http://www.mdpi.com/2072-6694/9/2/20>
- [8] Hsu NR| U. Role of oestrogen receptors in bladder cancer development. Nat Publ Gr [Internet]. 2013 [cited 2021 Mar 19];10:317-26. Available from: www.nature.com/nrurol
- [9] Arnold AP. Conceptual frameworks and mouse models for studying sex differences in physiology and disease: Why compensation changes the game [Internet]. Vol. 259, Experimental Neurology. Academic Press Inc.; 2014 [cited 2021 Mar 28]. p. 2-9. Available from: [/pmc/articles/PMC4125548/](http://pmc/articles/PMC4125548/)
- [10] Mäkelä VJ, Kotsar A, Tammela TLJ, Murtola TJ. Bladder Cancer Survival of Men Receiving 5 α -Reductase Inhibitors. J Urol. 2018 Oct 1;200(4):743-8.
- [11] Izumi K, Taguri M, Miyamoto H, Hara Y, Kishida T, Chiba K, et al. Androgen deprivation therapy prevents bladder cancer recurrence. Oncotarget [Internet]. 2014 [cited 2021 Mar 11];5(24):12665-74. Available from: [/pmc/articles/PMC4350350/](http://pmc/articles/PMC4350350/)
- [12] Laor E, Schiffman ZJ, Braunstein JD, Reid RE, Tolia BM, Koss LG, et al. Androgen receptors in bladder tumors. Urology. 1985 Feb 1;25(2):161-3.
- [13] Miyamoto H, Yao JL, Chaux A, Zheng Y, Hsu I, Izumi K, et al. Expression of androgen and oestrogen receptors and its prognostic significance in urothelial neoplasm of the urinary bladder. BJU Int [Internet]. 2012 Jun 1 [cited 2021 Mar 10];109(11):1716-26. Available from: <http://doi.wiley.com/10.1111/j.1464-410X.2011.10706.x>
- [14] Tuygun C, Kankaya D, Imamoglu A, Sertcelik A, Zengin K, Oktay M, et al. Sex-specific hormone receptors in urothelial carcinomas of the human urinary bladder: A comparative analysis

of clinicopathological features and survival outcomes according to receptor expression. *Urol Oncol Semin Orig Investig.* 2011 Jan 1;29(1):43-51.

[15] Nam JK, Park SW, Lee SD, Chung MK. Prognostic Value of Sex-Hormone Receptor Expression in Non-Muscle-Invasive Bladder Cancer. *Yonsei Med J* [Internet]. 2014 Sep 1 [cited 2021 Mar 10];55(5):1214. Available from: <https://eymj.org/DOIx.php?id=10.3349/ymj.2014.55.5.1214>

[16] Williams EM, Higgins JP, Ankur ., Sangoi R, Mckenney JK, Troxell ML. Androgen receptor immunohistochemistry in genitourinary neoplasms. *Int Urol Nephrol.* 2015;3:81-5.

[17] Boorjian S, Ugras S, Mongan NP, Gudas LJ, You X, Tickoo SK, et al. Androgen receptor expression is inversely correlated with pathologic tumor stage in bladder cancer. *Urology.* 2004 Aug 1;64(2):383-8.

[18] Sottnik JL, Vanderlinden LA, Joshi M, Chauca-Diaz A, Owens C, Hansel DE, et al. Androgen Receptor Regulates CD44 Expression in Bladder Cancer. *Cancer Res* [Internet]. 2021 Mar 9 [cited 2021 Mar 26];canres.3095.2020. Available from: <http://cancerres.aacrjournals.org/lookup/doi/10.1158/0008-5472.CAN-20-3095>

[19] Bertram JS, Craig AW. Specific induction of bladder cancer in mice by butyl-(4-hydroxybutyl)-nitrosamine and the effects of hormonal modifications on the sex difference in response. *Eur J Cancer.* 1972 Dec 1;8(6):587-94.

[20] Okajima E, Hiramatsu T, Iriya K, Ijuin M, Matsushima S, Yamada K. Effect of sex hormones on development of urinary bladder tumours in rats induced by N-butyl-N-(4-hydroxybutyl) nitrosamine. *Urol Res* [Internet]. 1975 Aug [cited 2021 Mar 10];3(2):73-9.

Available from: <https://link.springer.com/article/10.1007/BF00256185>

[21] Miyamoto H, Yang Z, Chen Y-T, Ishiguro H, Uemura H, Kubota Y, et al. Promotion of Bladder Cancer Development and Progression by Androgen Receptor Signals. *JNCI J Natl Cancer Inst* [Internet]. 2007 Apr 4 [cited 2021 Mar 10];99(7):558-68. Available from: <https://academic.oup.com/jnci/article-lookup/doi/10.1093/jnci/djk113>

[22] Hsu JW, Hsu I, Xu D, Miyamoto H, Liang L, Wu XR, et al. Decreased tumorigenesis and mortality from bladder cancer in mice lacking urothelial androgen receptor. *Am J Pathol.* 2013 May 1;182(5):1811-20.

[23] Johnson DT, Hooker E, Luong R, Yu E-J, He Y, Gonzalgo ML, et al. Conditional Expression of the Androgen Receptor Increases Susceptibility of Bladder Cancer in Mice. Agoulnik IU, editor. *PLoS One* [Internet]. 2016 Feb 10 [cited 2021 Mar 10];11(2):e0148851. Available from: <https://dx.plos.org/10.1371/journal.pone.0148851>

[24] Lin C, Yin Y, Stemler K, Humphrey P, Kibel AS, Mysorekar IU, et al. Constitutive β -catenin activation induces male-specific tumorigenesis in the bladder urothelium. *Cancer Res* [Internet]. 2013 Oct 1 [cited 2021 Mar 21];73(19):5914-25. Available from: <http://cancerres.aacrjournals.org/>

[25] Li Y, Zheng Y, Izumi K, Ishiguro H, Ye B, Li F, et al. Androgen activates β -catenin signaling in bladder cancer cells. *Endocr Relat Cancer* [Internet]. 2013 Jun 20 [cited 2021 Mar 21];20(3):293-304. Available from: <https://erc.bioscientifica.com/view/journals/erc/20/3/293.xml>

[26] Thomas S, Harding MA, Smith SC, Overdevest JB, Nitz MD, Frierson HF, et al. CD24 is an effector of HIF-1-driven primary tumor growth and metastasis.

- Cancer Res [Internet]. 2012 Nov 1 [cited 2021 Mar 21];72(21):5600-12. Available from: <http://cancerres.aacrjournals.org/>
- [27] Overdevest JB, Thomas S, Kristiansen G, Hansel DE, Smith SC, Theodorescu D. CD24 offers a therapeutic target for control of bladder cancer metastasis based on a requirement for lung colonization. Cancer Res [Internet]. 2011 Jun 1 [cited 2021 Mar 21];71(11):3802-11. Available from: <http://cancerres.aacrjournals.org/>
- [28] Overdevest JB, Knubel KH, Duex JE, Thomas S, Nitz MD, Harding MA, et al. CD24 expression is important in male urothelial tumorigenesis and metastasis in mice and is androgen regulated. Proc Natl Acad Sci U S A [Internet]. 2012 Dec 18 [cited 2021 Mar 21];109(51):E3588-96. Available from: www.pnas.org/cgi/doi/10.1073/pnas.1113960109
- [29] McGrath M, Michaud DS, De Vivo I. Hormonal and Reproductive Factors and the Risk of Bladder Cancer in Women. Am J Epidemiol [Internet]. 2006 Feb 1 [cited 2021 Mar 10];163(3):236-44. Available from: <http://academic.oup.com/aje/article/163/3/236/59618/Hormonal-and-Reproductive-Factors-and-the-Risk-of>
- [30] Dietrich K, Demidenko E, Schned A, Zens MS, Heaney J, Karagas MR. Parity, early menopause and the incidence of bladder cancer in women: A case-control study and meta-analysis. Eur J Cancer. 2011 Mar 1;47(4):592-9.
- [31] Wolpert BJ, Amr S, Ezzat S, Saleh D, Gouda I, Loay I, et al. Estrogen exposure and bladder cancer risk in Egyptian women. Maturitas. 2010 Dec 1;67(4):353-7.
- [32] Teng J, Wang ZY, Jarrard DF, Bjorling DE. Roles of estrogen receptor α and β in modulating urothelial cell proliferation. Endocr Relat Cancer [Internet]. 2008 Mar [cited 2021 Mar 11];15(1):351-64. Available from: pmc/articles/PMC3513362/
- [33] Basakci A, Kirkali Z, Tuzel E, Yorukoglu K, Mungan MU, Sade M. Prognostic significance of estrogen receptor expression in superficial transitional cell carcinoma of the urinary bladder. Eur Urol. 2002 Mar 1;41(3):342-5.
- [34] Shen SS, Smith CL, Hsieh J-T, Yu J, Kim IY, Jian W, et al. Expression of estrogen receptors- α and - β in bladder cancer cell lines and human bladder tumor tissue. Cancer [Internet]. 2006 Jun 15 [cited 2021 Mar 11];106(12):2610-6. Available from: <http://doi.wiley.com/10.1002/cncr.21945>
- [35] Miyamoto H, Yao JL, Chaux A, Zheng Y, Hsu I, Izumi K, et al. Expression of androgen and oestrogen receptors and its prognostic significance in urothelial neoplasm of the urinary bladder. BJU Int [Internet]. 2012 Jun 1 [cited 2021 Mar 11];109(11):1716-26. Available from: <http://doi.wiley.com/10.1111/j.1464-410X.2011.10706.x>
- [36] Bolenz C, Lotan Y, Ashfaq R, Shariat SF. Estrogen and Progesterone Hormonal Receptor Expression in Urothelial Carcinoma of the Bladder. Eur Urol [Internet]. 2009 Dec 7 [cited 2021 Mar 11];56(6):1093-5. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0302283809006769>
- [37] Croft PR, Lathrop SL, Feddersen RM, Joste NE. Estrogen receptor expression in papillary urothelial carcinoma of the bladder and ovarian transitional cell carcinoma. In: Archives of Pathology and Laboratory Medicine [Internet]. Arch Pathol Lab Med; 2005 [cited 2021 Mar 11]. p. 194-9. Available from: <https://pubmed.ncbi.nlm.nih.gov/15679420/>
- [38] Kontos S, Kominea A, Melachrinou M, Balampani E,

- Sotiropoulou-Bonikou G. Inverse expression of estrogen receptor- β and nuclear factor- κ B in urinary bladder carcinogenesis. *Int J Urol* [Internet]. 2010 Sep 1 [cited 2021 Mar 11];17(9):801-9. Available from: <http://doi.wiley.com/10.1111/j.1442-2042.2010.02603.x>
- [39] Hsu I, Yeh CR, Slavin S, Miyamoto H, Netto GJ, Tsai YC, et al. Estrogen receptor alpha prevents bladder cancer development via INPP4B inhibited Akt pathway in vitro and in vivo. *Oncotarget* [Internet]. 2014 [cited 2021 Mar 11];5(17):7917-35. Available from: [/pmc/articles/PMC4202170/](http://pmc/articles/PMC4202170/)
- [40] Hsu I, Chuang K-L, Slavin S, Da J, Lim W-X, Pang S-T, et al. Suppression of ER β signaling via ER β knockout or antagonist protects against bladder cancer development. *Carcinogenesis* [Internet]. 2014 Mar 1 [cited 2021 Mar 11];35(3):651-61. Available from: <https://academic.oup.com/carcin/article-lookup/doi/10.1093/carcin/bgt348>
- [41] George SK, Tovar-Sepulveda V, Shen SS, Jian W, Zhang Y, Hilsenbeck SG, et al. Chemoprevention of BBN-induced bladder carcinogenesis by the selective Estrogen receptor modulator tamoxifen. *Transl Oncol*. 2013 Jun 1;6(3):244-55.
- [42] Schoemaker MJ, Swerdlow AJ, Higgins CD, Wright AF, Jacobs PA. Cancer incidence in women with Turner syndrome in Great Britain: a national cohort study. *Lancet Oncol* [Internet]. 2008 Mar [cited 2021 Mar 11];9(3):239-46. Available from: <https://pubmed.ncbi.nlm.nih.gov/18282803/>
- [43] Ji J, Zöller B, Sundquist J, Sundquist K. Risk of solid tumors and hematological malignancy in persons with Turner and Klinefelter syndromes: A national cohort study. *Int J Cancer* [Internet]. 2016 Aug 15 [cited 2021 Mar 11];139(4):754-8. Available from: <http://doi.wiley.com/10.1002/ijc.30126>
- [44] De Vries GJ, Rissman EF, Simerly RB, Yang LY, Scordalakes EM, Auger CJ, et al. A model system for study of sex chromosome effects on sexually dimorphic neural and behavioral traits. *J Neurosci* [Internet]. 2002 Oct 15 [cited 2021 Mar 11];22(20):9005-14. Available from: <https://www.jneurosci.org/content/22/20/9005>
- [45] Burgoyne PS, Arnold AP. A primer on the use of mouse models for identifying direct sex chromosome effects that cause sex differences in non-gonadal tissues [Internet]. Vol. 7, *Biology of Sex Differences*. BioMed Central Ltd.; 2016 [cited 2021 Mar 22]. p. 1-21. Available from: [/pmc/articles/PMC5154145/](http://pmc/articles/PMC5154145/)
- [46] Kaneko S, Li X. X chromosome protects against bladder cancer in females via a KDM6A-dependent epigenetic mechanism. *Sci Adv* [Internet]. 2018 Jun 13 [cited 2021 Mar 10];4(6):eaar5598. Available from: <http://advances.sciencemag.org/>
- [47] Fadl-Elmula I, Gorunova L, Mandahl N, Elfving P, Lundgren R, Mitelman F, et al. Karyotypic characterization of urinary bladder transitional cell carcinomas. *Genes Chromosom Cancer* [Internet]. 2000 Nov 1 [cited 2021 Mar 11];29(3):256-65. Available from: <https://onlinelibrary.wiley.com/doi/full/10.1002/1098-2264%282000%2999999%3A9999%3C%3A%3AAID-GCC1034%3E3.0.CO%3B2-O>
- [48] Minner S, Kilgué A, Stahl P, Weikert S, Rink M, Dahlem R, et al. Y chromosome loss is a frequent early event in urothelial bladder cancer. *Pathology*. 2010;42(4):356-9.
- [49] Forsberg LA, Rasi C, Malmqvist N, Davies H, Pasupulati S, Pakalapati G, et al. Mosaic loss of chromosome y in peripheral blood is associated with shorter survival and higher risk of

cancer. *Nat Genet* [Internet]. 2014 Apr 28 [cited 2021 Mar 11];46(6):624-8. Available from: <https://www.nature.com/articles/ng.2966>

[50] Tukiainen T, Villani AC, Yen A, Rivas MA, Marshall JL, Satija R, et al. Landscape of X chromosome inactivation across human tissues. *Nature* [Internet]. 2017 Oct 11 [cited 2021 Mar 19];550(7675):244-8. Available from: <https://www.nature.com/articles/nature24265>

[51] Robertson AG, Kim J, Al-Ahmadie H, Bellmunt J, Guo G, Cherniack AD, et al. Comprehensive Molecular Characterization of Muscle-Invasive Bladder Cancer. *Cell*. 2017 Oct 19;171(3):540-556.e25.

[52] Ler LD, Ghosh S, Chai X, Thike AA, Heng HL, Siew EY, et al. Loss of tumor suppressor KDM6A amplifies PRC2-regulated transcriptional repression in bladder cancer and can be targeted through inhibition of EZH2. *Sci Transl Med* [Internet]. 2017 Feb 22 [cited 2021 Mar 11];9(378). Available from: <http://stm.sciencemag.org/>

[53] Nordentoft I, Lamy P, Birkenkamp-Demtröder K, Shumansky K, Vang S, Hornshøj H, et al. Mutational context and diverse clonal development in early and late bladder cancer. *Cell Rep*. 2014 Jun 12;7(5):1649-63.

[54] Hurst CD, Alder O, Platt FM, Droop A, Stead LF, Burns JE, et al. Genomic Subtypes of Non-invasive Bladder Cancer with Distinct Metabolic Profile and Female Gender Bias in KDM6A Mutation Frequency. *Cancer Cell*. 2017 Nov 13;32(5):701-715.e7.

[55] Shpargel KB, Sengoku T, Yokoyama S, Magnuson T. UTX and UTY Demonstrate Histone Demethylase-Independent Function in Mouse Embryonic Development. Wysocka J, editor. *PLoS Genet*

[Internet]. 2012 Sep 27 [cited 2021 Mar 11];8(9):e1002964. Available from: <https://dx.plos.org/10.1371/journal.pgen.1002964>

[56] Wang C, Lee JE, Cho YW, Xiao Y, Jin Q, Liu C, et al. UTX regulates mesoderm differentiation of embryonic stem cells independent of H3K27 demethylase activity. *Proc Natl Acad Sci U S A* [Internet]. 2012 Sep 18 [cited 2021 Mar 11];109(38):15324-9. Available from: www.pnas.org/cgi/doi/10.1073/pnas.1204166109

[57] Gozdecka M, Meduri E, Mazan M, Tzelepis K, Dudek M, Knights AJ, et al. UTX-mediated enhancer and chromatin remodeling suppresses myeloid leukemogenesis through noncatalytic inverse regulation of ETS and GATA programs. *Nat Genet* [Internet]. 2018 Jun 1 [cited 2021 Mar 11];50(6):883-94. Available from: <https://doi.org/10.1038/s41588-018-0114-z>

[58] Welstead GG, Creighton MP, Bilodeau S, Cheng AW, Markoulaki S, Young RA, et al. X-linked H3K27me3 demethylase Utx is required for embryonic development in a sex-specific manner. *Proc Natl Acad Sci U S A* [Internet]. 2012 Aug 7 [cited 2021 Mar 11];109(32):13004-9. Available from: www.pnas.org/cgi/doi/10.1073/pnas.1210787109

[59] Dunford A, Weinstock DM, Savova V, Schumacher SE, Cleary JP, Yoda A, et al. Tumor-suppressor genes that escape from X-inactivation contribute to cancer sex bias. *Nat Genet* [Internet]. 2017 Jan 1 [cited 2021 Mar 21];49(1):10-6. Available from: [/pmc/articles/PMC5206905/](https://pmc/articles/PMC5206905/)

[60] Agger K, Cloos PAC, Christensen J, Pasini D, Rose S, Rappsilber J, et al. UTX and JMJD3 are histone H3K27 demethylases involved in HOX gene regulation and development. *Nature* [Internet]. 2007 Oct 11 [cited 2021 Mar 19];449(7163):731-4. Available from:

<https://www.nature.com/articles/nature06145>

[61] Hong SH, Cho YW, Yu LR, Yu H, Veenstra TD, Ge K. Identification of JmjC domain-containing UTX and JMJD3 as histone H3 lysine 27 demethylases. *Proc Natl Acad Sci U S A* [Internet]. 2007 Nov 20 [cited 2021 Mar 22];104(47):18439-44. Available from: www.pnas.org/cgi/doi/10.1073/pnas.0707292104

[62] Lan F, Bayliss PE, Rinn JL, Whetstone JR, Wang JK, Chen S, et al. A histone H3 lysine 27 demethylase regulates animal posterior development. *Nature* [Internet]. 2007 Oct 11 [cited 2021 Mar 22];449(7163):689-94. Available from: <https://www.nature.com/articles/nature06192>

[63] Min GL, Villa R, Trojer P, Norman J, Yan KP, Reinberg D, et al. Demethylation of H3K27 regulates polycomb recruitment and H2A ubiquitination. *Science* (80-) [Internet]. 2007 Oct 19 [cited 2021 Mar 19];318(5849):447-50. Available from: <http://science.sciencemag.org/>

[64] Cho YW, Hong T, Hong SH, Guo H, Yu H, Kim D, et al. PTIP associates with MLL3- and MLL4-containing histone H3 lysine 4 methyltransferase complex. *J Biol Chem*. 2007 Jul 13;282(28):20395-406.

[65] Wang SP, Tang Z, Chen CW, Shimada M, Koche RP, Wang LH, et al. A UTX-MLL4-p300 Transcriptional Regulatory Network Coordinately Shapes Active Enhancer Landscapes for Eliciting Transcription. *Mol Cell*. 2017 Jul 20;67(2):308-321.e6.

[66] Dorigi KM, Swigut T, Henriques T, Bhanu N V., Scruggs BS, Nady N, et al. Mll3 and Mll4 Facilitate Enhancer RNA Synthesis and Transcription from Promoters Independently of H3K4 Monomethylation. *Mol Cell*. 2017 May 18;66(4):568-576.e4.

[67] Rickels R, Herz H-M, Sze CC, Cao K, Morgan MA, Collings CK, et al. Histone H3K4 monomethylation catalyzed by Trr and mammalian COMPASS-like proteins at enhancers is dispensable for development and viability. *Nat Genet* Vol. 2017;49(11).

[68] Cenik BK, Shilatifard A. COMPASS and SWI/SNF complexes in development and disease [Internet]. Vol. 22, *Nature Reviews Genetics*. Nature Research; 2021 [cited 2021 Mar 19]. p. 38-58. Available from: www.nature.com/nrg

[69] Piunti A, Shilatifard A. Epigenetic balance of gene expression by Polycomb and COMPASS families. *Science* (80-) [Internet]. 2016 Jun 3 [cited 2021 Mar 19];352(6290):aad9780. Available from: <https://www.sciencemag.org/lookup/doi/10.1126/science.aad9780>

[70] Nickerson ML, Witte N, Im KM, Turan S, Owens C, Misner K, et al. Molecular analysis of urothelial cancer cell lines for modeling tumor biology and drug response. *Oncogene* [Internet]. 2017 Jan 5 [cited 2021 Mar 22];36(1):35-46. Available from: [/pmc/articles/PMC5140783/](https://pubmed.ncbi.nlm.nih.gov/240783/)