We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Blunt Traumatic Aortic Injury

Domenico Calcaterra

Abstract

Traumatic aortic injuries represent a leading cause of death following motor-vehicular accidents. These injuries carry a very high mortality rate even though a significant number of patients reaches the hospital alive. These injuries are identified in the contest of a polytrauma work up and are almost always associated with multiple other severe traumatic injuries which makes the management of these patients very challenging. The technology advancements seen in recent years with radiologic imaging and the progress of the therapeutic options brought up by the uprise of endovascular therapy, along with the sophistication of the techniques of trauma resuscitation and intensive care management, have improved significantly the overall prognosis of these patients. Although traumatic aortic injuries need to be generally considered a life-threatening condition, their degree of severity may differ significantly from case to case requiring immediate repair in some patients, whereas their repair can be delayed in cases when the severity of the aortic injury does not represent an immediate threat to the patient life. Therefore, the challenge of treatment of the polytrauma patients with an aortic injury is to identify the best strategy of therapy able to prioritize the treatment of the injuries based on their lethal potential. In this contest, the ability of properly defining the severity of the aortic injury is the key-factor to allow the appropriate definition of a treatment strategy able to identify treatment priorities. In our experience, radiologic assessment of the aortic injury in correlation with the evaluation of clinical parameters and a comprehensive polytrauma assessment allows to optimize the ability of the trauma team to establish the most appropriate strategy for the care of this complex patients' group.

Keywords: polytrauma assessment, traumatic aortic injury, thoracic endovascular aortic repair, radiologic assessment

1. Introduction

Traumatic aortic injuries (TAI) represent the second leading cause of death from motor vehicle crashes, accounting for 15% of all motor vehicle accident associated deaths [1–3]. According to a historical case series, death occurs at the scene of the accident in 70 to 90% of these cases [1, 3–7], and of the patients (75%) who arrive to the hospital alive, although hemodynamically stable, only 10% survives more than 6 hours [1, 3]. Patients with TAI surviving at the scene who arrive to the hospital alive most frequently present with an injury at the aortic isthmus, since periadventitial tissue in this location seems to provide some degree of protection against free rupture, allowing the necessary time to transfer the patient from the trauma scene to the hospital [8–10]. The majority of patients with BAI have an

Figure 1. Theories of blunt aortic injury. Blunt aortic injuries involve a combination of forces, including stretching, shearing, torsion, a “water-hammer” effect (which involves simultaneous occlusion of the aorta and a sudden elevation in blood pressure), and the “osseous pinch” effect from entrapment of the aorta between the anterior chest wall and the vertebral column (modified with permission) [11].

associated closed head injury (51%), multiple rib fractures (46%), lung contusions (38%), or orthopedic injuries (20–35%) [1].

2. Pathophysiology

Blunt traumatic aortic injuries can involve any thoracic aortic segment, including occasionally even the abdominal aorta. The aortic isthmus is by far the most common location, followed by the ascending aorta (10–25%), the aortic arch (10–20%) and the abdominal aorta (5–10%). The theory is that a combination of sudden deceleration, associated to torsion, stretching and shearing forces, and thoracic compression, would cause the aortic injury (**Figure 1**) [11].

3. Clinical management

Diagnosis is made in the context of the trauma work up as defined by the Advanced Trauma Life Support (ATLS) guidelines. Computed tomography (CT)

scan is by far the test of choice to diagnose TAI with a sensitivity and negative predictive value close to 100%. In the very unusual circumstance that CT scan does not provide a definite answer and some doubt on the presence of a TAI remains, which would only occur in the case of a minimal injury, aortic angiogram, eventually with intravascular ultrasound (IVUS), represents the gold-standard to reach a definitive diagnosis. Once diagnosis is made, the severity of the injury assessed by imaging studies and the polytrauma assessment will dictate the treatment strategy [12].

Based on the application of ATLS protocols, treatment of the different trauma injuries will be prioritized based on their acute lethal potential [10]. Exsanguinating hemorrhages from any location and intracranial injury with mass effect take priority of treatment, unless hemodynamic instability obviously related to the imaging finding of an extremely unstable aortic injury would suggest to proceed with immediate endovascular aortic repair. In general, clinical management of the trauma patient arriving to the hospital requires the application of standard measures of trauma resuscitation aiming at establishing the best possible hemodynamic conditions. Once a TAI is diagnosed, anti-impulsive therapy with short-acting beta-blockers should be instituted, if allowed by hemodynamic conditions, to reduce aortic wall-stress. At that point the timing of the aortic repair should be decided based on the radiologic assessment of the aortic injury, the patient's general conditions, accounting in the decision-making process a polytrauma assessment which will allow to determine the sequence of therapeutic interventions offering the best chance of a positive clinical outcome [10].

In fact, although the vast majority of aortic injuries, based on a traditional 'old-school' approach, would represent an indication for therapeutic intervention, there has been more recently a strong school of thoughts proposing a conservative type of management for the type of injuries with a low lethal potential. In these cases, the therapeutic intervention can be delayed or completely aborted, selecting a strategy of radiologic monitoring which would allow to indicate a need for intervention only in cases showing evolution of the aortic injury in a growing pseudo-aneurysm [13–18].

The Society for Vascular Surgery (SVS) has proposed a grading system for TAI intended to rate the degree of severity of the injury (**Figure 2**) [19, 20]. Nevertheless, this grading system has failed to find reliable clinical correlation with risk of aortic rupture and death [15, 19–22], because this classification is qualitative but not quantitative, since can be useful to define the type of injury (intimal laceration versus intramural hematoma, versus pseudoaneurysm, versus free rupture), but does not include parameters to define size and extension of the injury. In our experience of blunt aortic injuries from 3 Level I Trauma Centers in the US from July 2008 to December 2016, we reviewed a total of 76 patients [12]. We analyzed overall mortality and TAI-related mortality (directly caused by the effects of the aortic injury) at 30 days in relation to factors such as: hemodynamic parameters on presentation (SBP, HR and need for vasopressor medications), timing of treatment, injury severity score (ISS) and aortic injury grade as defined by the Society for Vascular Surgery Clinical Practice Guidelines. Aortic injury (AI) grade was dichotomized as stable, grade I-II, and unstable, grade III-IV [12]. Using a new injury scale system, we classified the AI as "Severe" (Radiographic Severe Injury, RSI) when they included findings of [1] total/partial aortic transection (**Figure 3**), [2] active contrast extravasation (**Figure 4**), or [3] the association of 2 of more of the following: contained contrast extravasation >10 mm in bigger dimension, periaortic hematoma and/or mediastinal hematoma with >10 mm thickness, or left pleural effusion (**Figure 5**). We found that mortality caused by the aortic injury was associated with high ISS, SBP < 100, HR ≥ 100, and vasopressors requirement. Also, our

Figure 2.

The Society for Vascular Surgery classification of traumatic aortic injury. Grade I: Intimal tear; grade II: Intramural hematoma; grade III: Pseudoaneurysm; grade IV: Rupture. This grading system has failed to find reliable clinical correlation with risk of aortic rupture and death and therefore cannot find use to indicate the lethal potential of the aortic injury and support the necessary choices that need to be made to select the most appropriate therapeutic strategy to improve the prognosis of the polytrauma patient with TAI (modified with permission) [19].

new classification system of RSI, identifying patients with ‘unstable’ injuries, found statistically significant association with mortality (**Table 1**).

Therefore, our proposed system of grading of the aortic injury based on radiologic findings and the evaluation of clinical parameters, by the assessment of hemodynamic conditions (SBP, HR, and pressors requirement), is the most important elements to define the severity of the aortic injury and its lethal potential [12].

Besides the exceptional technical advancement of imaging studies that has allowed to increase tremendously the sensitivity to diagnose TAI, the most significant stride in the management of these injuries has been made by the rise of endovascular therapy, since treatment can be delivered with a faster approach using this much less invasive transcatheter technique and with substantial less operative and perioperative risk, compared to the ‘open’ surgical technique of aorta replacement used as the standard approach until a decade ago [15, 23–25].

Nonetheless, the choice of the most adequate timing for treatment of the aortic injury, particularly with respect to other major traumatic injuries, remains an area of active study. There are currently no clear guidelines for determining which patients may benefit from delayed aortic repair, nor there are validated methods of assessment of the severity of the aortic injury which would allow to choose when prioritize treatment of the aorta [15, 21]. A recent review of a small number of cases has suggested that some patients with small size pseudoaneurysms may be safely managed nonoperatively for the long-term [15, 25, 26]. Nevertheless, the ideal management for stable pseudoaneurysms after BAI remains a subject needing further study.

Figure 3.
CT scan of the chest with IV contrast demonstrating aortic transection near the isthmus (red arrow), also associated with intraluminal aortic thrombus as shown by the blue arrow. (transection is defined by total or partial interruption of the column of intra-venous contrast flowing within the aortic lumen).

In this contest, our system of classification created criteria for radiographic assessment of the degree of aortic injury used as a binary variable (severe versus non-severe), allowing to identify the patients in needs for immediate aortic repair.

The standard ‘open’ surgical technique of repair of TAI has been the replacement of the damaged aortic segment with a synthetic vascular graft. The most common

Figure 4.
Sagittal and transverse CT scan images of large intravenous contrast extravasation as shown by red arrows (irregular IV contrast contour outside the aortic wall boundaries).

location of the injury at the aortic isthmus requires a left thoracotomy approach, completing the aortic replacement with left atrial to femoral by-pass (**Figure 6**). This technique had replaced the previous approach of ‘clamp and saw’ (without use of partial cardiopulmonary bypass), which was associated with a very high incidence of complications, of which paraplegia secondary to spinal cord ischemia was the most common and devastating one. The strategy of replacement using left atrial to femoral bypass allows to maintain perfusion of the lower body after

Figure 5.
 Coronal and transverse CT scan images of periaortic hematoma (red arrows) and aortic pseudoaneurysm as shown by white arrows (pseudoaneurysm is defined by a regular intravenous contrast contour outside the aortic wall boundaries).

cross-clamping of the aorta above the injured segment, providing a better degree of protection of the spinal cord and other end organs from ischemia (**Figure 6**).

Nonetheless, the operation is still associated with significant morbidity and difficult postoperative recovery, considering that has to be accomplished under the very dangerous conditions of an extreme emergency in patients most likely affected by multiple other traumatic injuries.

The real revolution in the treatment of TAI has been accomplished with the uprise of endovascular aortic repair, which has impacted remarkably on the overall prognosis of these patients allowing to obtain an expedite aortic repair, without the need of an open surgical approach, which translates in much lesser procedural stress, much lesser operative risk, especially in the contest of the common poor general condition of the polytrauma patient, and much easier postprocedural recovery.

The other significant progress in the treatment of TAI has been the realization that under certain conditions the repair of TAI ought to be deferred, prioritizing the treatment of other major concomitant traumatic injuries which represented a more immediate danger for the patient. In a relatively recent prospective multicenter study sponsored by the American Association for the Surgery of Trauma (AAST), the effect of early versus delayed repair was observed in 178 patients admitted with BAI between 2005 and 2007. The study concluded that 'delayed repair of 'stable' blunt thoracic aortic injuries is associated with improved survival [16].

The decision establishing the timing of treatment of TAI should be exclusively based on the characteristics of the injury as seen on CT scan imaging and on the assessment of clinical factors in relation to other associated injuries [15, 21–24]. The Injury Severity Score (ISS) has been used to predict risk of morbidity and mortality associated with blunt trauma since the 1970s. It was demonstrated initially to correlate well with length of stay, need for major surgery, significant disability, and death [27, 28]. ISS does have known limitations, such as more limited applicability to penetrating trauma or other trauma patients in which injuries are localized only to one body area [28]. However, it continues to be a valuable tool used prominently in trauma databases to assign an objective value to traumatic injuries and predict risk for significant morbidity/mortality.

Unadjusted Mortality			Unadjusted BAI-related mortality	
	OR (CI)	P value ¹	OR (CI)	P value ¹
Age	1.01 (0.98, 1.04)	0.406	1.01 (0.98, 1.04)	0.665
ISS	1.07(1.02, 1.11)	0.003	1.07 (1.02, 1.12)	0.004
SBP < 100	10.54(2.61, 42.65)	<0.001	24.00 (2.84, 203.14)	0.004
HR ≥ 100	4.88 (1.37, 17.44)	0.015	7.48 (1.47, 38.17)	0.016
Pressors	7.56 (2.12, 29.12)	0.002	6.33 (1.52, 26.33)	0.011
AI grade(SVS)	2.65 (0.67, 10.45)	0.164	6.63 (0.79, 55.41)	0.081
RSI	3.02 (0.92, 9.90)	0.068	5.37 (1.28, 22.90)	0.023
Adjusted Mortality			Adjusted BAI-related mortality ²	
	OR (CI)	P value ²	OR (CI)	P value ²
ISS	1.04(1.00, 1.09)	0.074	1.05 (1.00, 1.11)	0.062
SBP < 100	5.54(0.71, 43.47)	0.103	13.16 (0.59, 195.18)	0.061
HR ≥ 100	0.99 (0.14, 6.88)	0.991	1.16 (0.12, 10.74)	0.898
Pressors	1.55 (0.28, 8.66)	0.616	0.77(0.12, 4.96)	0.786
RSI	—		2.51 (0.46, 13.75)	0.290

¹Univariate logistic regression.
²Multivariate logistic regression.
Abbreviations: AI, aortic injury grade group; BAI, blunt aortic injury CI, 95% confidence interval; HR, heart rate; ISS, injury severity score; OR, odds ratio; RSI, radio graphic severe injury; SBP, systolic blood pressure; SVS, society for vascular surgery.

Table 1.
Logistic regression of factors predicting risk of overall mortality and BAI-related mortality [12].

Thoracic endovascular aortic replacement (TEVAR) offers the advantages of a fast delivery of therapy, preventing a dangerous operation with partial cardiopulmonary bypass or hypothermic circulatory arrest, and less risk of postoperative paraplegia. Furthermore, endovascular therapy can be delivered in the operating room under portable fluoroscopy, offering the tremendous benefit of allowing simultaneous delivery of other therapies for associated life-threatening injuries, such as cranial decompression, transcatheter embolization or exploratory laparotomy, which would be significantly delayed by performing an open surgical aortic repair. The benefit of endovascular therapy is supported by the findings in the literature that have consistently shown substantial advantages of TEVAR over open repair in TAI [15, 29–36]. In our series, open repair was selected only when TEVAR was not feasible, such as in cases with no peripheral aortic access due to presence of intraluminal aortic thrombus, small size of femoral vessels, or presence of a total aortic transection which prevented delivery of endovascular therapy, as seen in the case of **Figure 3**.

As last consideration, in our experience we have observed a relatively small number of patients who died before any treatment was established. If historical series had reported that number to be significant, most recent reports have shown that of the patients surviving BAI at the scene, less than 5% would die of a direct aortic complication after arrival at the hospital [23]. The improvement of the techniques of resuscitation and trauma management, along with a consistent and early application of anti-impulsive therapy have positively impacted on the post-admission hospital mortality [23].

Figure 6.
Technique of proximal descending thoracic aortic replacement with partial cardiopulmonary bypass: Oxygenated blood is drained from the left inferior pulmonary vein and pumped in the left common femoral artery, allowing perfusion of end organs distal to the aortic cross-clamping.

4. Conclusion

The surge of TEVAR as the new standard for treatment for TAI has lowered the operative mortality for the treatment of this condition. However, the optimal timing for the delivery of therapy remains still unclear with respect to the identification of the patients who would require immediate intervention versus the ones for whom postponing treatment of the aortic injury would be preferable. The newly conceived radiologic classification system of TAI we use in our clinical experience is aimed at identifying the type of injuries associated with the highest mortality risk. Radiologic assessment of the severity of the aortic injuries with characterization of the presence of an ‘unstable’ and life-threatening condition should represent the primary factor to direct management strategy indicating the timing for the aortic repair and guiding treatment priorities.

Author details

Domenico Calcaterra
Florida Atlantic University, Boca Raton, FL, USA

*Address all correspondence to: domenicocalcaterra@hotmail.com

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Williams, JS, Graff JA, Uku JM, Steinig JP. Aortic injury in vehicular trauma. *Ann Thorac Surg.* 1994; 57:726.
- [2] Razzouk AJ, Gundry SR, Wang N, del Rio MJ, Varnell D, Bailey LL. Repair of traumatic aortic rupture: A 25-year experience. *Arch Surg.* 2000; 135:913; discussion 919.
- [3] Fabian TC, Richardson JD, Croce MA, Smith JS Jr, Rodman G Jr, Kearney PA, Flynn W, Ney AL, Cone JB, Luchette FA, et al. Prospective study of blunt aortic injury: Multicenter Trial of the American Association for the Surgery of Trauma. *J Trauma.* 1997; 42:374-80: discussion 380-3.
- [4] Parmley L, Mattingly T, Manion W. Nonpenetrating traumatic injury of the aorta. *Circulation.* 1958; 17:1086.
- [5] Greendyke RM. Traumatic rupture of aorta; special reference to automobile accidents. *JAMA.* 1966; 195:527.
- [6] Feczko JD, Lynch L, Pless JE, Clark MA, McClain J, Hawley DA. An autopsy case review of 142 nonpenetrating (blunt) injuries of the aorta. *J Trauma.* 1992; 33:846.
- [7] Smith RS, Chang FC. Traumatic rupture of the aorta: Still a lethal injury. *Am J Surg.* 1986; 152:660.
- [8] Arajärvi E, Santavirta S, Tolonen J. Aortic ruptures in seat belt wearers. *J Thorac Cardiovasc Surg.* 1989; 98:355.
- [9] Rabinsky I, Sidhu GS, Wagner RB. Mid-descending aortic traumatic aneurysms. *Ann Thorac Surg.* 1990; 50:155.
- [10] Gleason TG, Bavaria JE. Trauma to the Great Vessels. Cohn LH, ed. *Cardiac Surgery in the Adult.* New York: McGraw-Hill, 2008:1333-1354.
- [11] Neschis DG, Scalea TM, Flinn WR, Griffin BP. Blunt Aortic Injury. *N Engl J Med.* 2008; 359:1708-1716.
- [12] Payne RE, Nygaard RM, Fernandez JD, Sahgal P, Richardson CJ, Bashir M, Parekh K, Vardas PN, Suzuki Y, Corvera J, Krook JC, Calcaterra D. Blunt aortic injuries in the new era: radiologic findings and polytrauma risk assessment dictates management strategy. *Eur J Trauma Emerg Surg.* 2019 Dec;45(6):951-957.
- [13] Symbas PN, Sherman AJ, Silver JM, Symbas JD, Lackey JJ. Traumatic rupture of the aorta. Immediate or delayed repair? *Ann Surg.* 2002; 235(6):796-802.
- [14] Estrera AL, Gochnour DC, Azizzadeh A, Miller CC 3rd, Coogan S, Charlton-Ouw K, Holcomb JB, Safi HJ. Progress in the Treatment of Blunt Thoracic Aortic Injury: 12-Year Single-Institution Experience. *Ann Thorac Surg.* 2010; 90(1):64-71.
- [15] Heneghan R, Aarabi S, Quiroga E, Gunn ML, Singh N, Starnes BW. Call for a new classification system and treatment strategy in blunt aortic injury. *J Vasc Surg.* 2016 Jul;64(1):171-176.
- [16] Demetriades D, Velmahos GC, Scalea TM, Jurkovich GJ, Karmy-Jones R, Teixeira PG, Hemmila MR, O'Connor JV, McKenney MO, Moore FO, London J, Singh MJ, Spaniolas K, Keel M, Sugrue M, Wahl WL, Hill J, Wall MJ, Moore EE, Lineen E, Margulies D, Malka V, Chan LS. Blunt traumatic thoracic aortic injuries: early or delayed repair-results of an American Association for the Surgery of Trauma prospective study. *J Trauma.* 2009 Apr;66(4):967-973.
- [17] Nagy K, Fabian T, Rodman G, Fulda G, Rodriguez A, Mirvis S. Guidelines for the diagnosis and

management of blunt aortic injury: An EAST Practice Management Guidelines Work Group. *J Trauma*. 2000; 48:1128-1143.

[18] Williams MJ, Low CJ, Wilkins GT, Stewart RA. Randomised comparison of the effects of nicardipine and esmolol on coronary artery wall stress: Implications for the risk of plaque rupture. *Heart*. 2000; 84:377.

[19] Lee WA, Matsumura JS, Mitchell RS, Farber MA, Greenberg RK, Azizzadeh A, Murad MH, Fairman RM. Endovascular repair of traumatic thoracic aortic injury: clinical practice guidelines of the Society for Vascular Surgery. *J Vasc Surg*. 2011 Jan;53(1):187-192.

[20] Azizzadeh A, Keyhani K, Miller CC 3rd, Coogan SM, Safi HJ, Estrera AL. Blunt traumatic aortic injury: initial experience with endovascular repair. *J Vasc Surg*. 2009 Jun;49(6):1403-1408.

[21] Fortuna GR Jr, Perlick A, DuBose JJ, Leake SS, Charlton-Ouw KM, Miller CC 3rd, Estrera AL, Azizzadeh A. Injury grade is a predictor of aortic-related death among patients with blunt thoracic aortic injury. *J Vasc Surg*. 2016 May;63(5):1225-1231.

[22] Lamarche Y, Berger FH, Nicolaou S, Bilawich AM, Louis L, Inacio JR, Janusz MT, Evans D. Vancouver simplified grading system with computed tomographic angiography for blunt aortic injury. *J Thorac Cardiovasc Surg*. 2012 Aug;144(2):347-354.

[23] Pacini D, Angeli E, Fattori R, Lovato L, Rocchi G, Di Marco L, Bergonzini M, Grillone G, Di Bartolomeo R. Traumatic rupture of the thoracic aorta: ten years of delayed management. *J Thorac Cardiovasc Surg*. 2005 Apr;129(4):880-884.

[24] Patelis N, Katsargyris A, Klonaris C. Endovascular Repair of Traumatic

Isthmic Ruptures: Special Concerns. *Front Surg*. 2017 Jun 12;4:32.

[25] Tanizaki S, Maeda S, Matano H, Sera M, Nagai H, Nakanishi T, Ishida H. Blunt thoracic aortic injury with small pseudoaneurysm may be managed by nonoperative treatment. *J Vasc Surg*. 2016 Feb;63(2):341-344.

[26] Osgood M, Heck J, Rellinger E, Doran SL, Garrard CL 3rd, Guzman RJ, Naslund TC, Dattilo JB. Natural history of grade I-II blunt traumatic aortic injury. *J Vasc Surg*. 2014 Feb;59(2):334-341.

[27] Semmlow JL, Cone R. Utility of the Injury Severity Score: a confirmation. *Health Serv Res*. 1976;11:45-52.

[28] Smith B, Goldberg AJ, Gaughan JP, Seamon MJ. A comparison of Injury Severity Score and New Injury Severity Score after penetrating trauma: A prospective analysis. *Journal of Trauma and Acute Care Surgery*. 2015;79(2):269-274.26.

[29] Xenos ES, Abedi NN, Davenport DL, Minion DJ, Hamdallah O, Sorial EE, Endean Ed. Meta-analysis of endovascular vs open repair for traumatic descending thoracic aortic rupture. *J Vasc Surg*. 2008; 48:1343-1351.

[30] Kwolek CJ, Blazick E. Current management of traumatic aortic injuries. *Semin Vasc Surg*. 2010 Dec;23(4):215-220.

[31] Akowuah E, Baumbach A, Wilde P, Angelini G, Bryan AJ. Emergency repair of traumatic aortic rupture: endovascular versus conventional open repair. *J Thorac Cardiovasc Surg*. 2007; 134(4):897-901.

[32] Verdant A. Endovascular management of traumatic aortic injuries. *Can J Surg*. 2006; 49(3):217; author reply 217-218.

[33] Andrassy J, Weidenhagen R, Meimarakis G, Rentsch M, Jauch KW, Kopp R. Stent versus open surgery for acute and chronic traumatic injury of the thoracic aorta: a single-center experience. *J Trauma*. 2006;60(4):765-771; discussion 771-772.

[34] Kokotsakis J, Kaskarelis I, Misthos P, Athanasiou T, Kanakakis K, Athanasiu C, Romana C, Skouteli E, Lioulis A. Endovascular versus open repair for blunt thoracic aortic injury: short-term results. *Ann Thorac Surg*. 2007; 84(6):1965-1970.

[35] Akowuah E, Angelini G, Bryan AJ. Open versus endovascular repair of traumatic aortic rupture: a systematic review. *J Thorac Cardiovasc Surg*. 2009; 138(3):768-769.

[36] Canaud L, Alric P, Branchereau P, Marty-Ané C, Mercier G, Branchereau P. Open versus endovascular repair for patients with acute traumatic rupture of the thoracic aorta. *J Thorac Cardiovasc Surg*. 2011; 142(5):1032-1037.