We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Industrial Applications of Cyanobacteria

*Ayesha Algade Amadu, Kweku Amoako Atta deGraft-Johnson
and Gabriel Komla Ameka*

Abstract

Cyanobacteria also known as blue-green algae are oxygenic photoautotrophs, which evolved ca. 3.5 billion years ago. Because cyanobacteria are rich sources of bioactive compounds, they have diverse industrial applications such as algaecides, antibacterial, antiviral and antifungal agents, hence, their wide use in the agricultural and health sectors. Cyanobacterial secondary metabolites are also important sources of enzymes, toxins, vitamins, and other pharmaceuticals. Polyhydroxyalkanoates (PHA) which accumulate intracellularly in some cyanobacteria species can be used in the production of bioplastics that have properties comparable to polypropylene and polyethylene. Some cyanobacteria are also employed in bioremediation as they are capable of oxidizing oil components and other complex organic compounds. There are many more possible industrial applications of cyanobacteria such as biofuel, biofertilizer, food, nutraceuticals, and pharmaceuticals. Additionally, the metabolic pathways that lead to the production of important cyanobacterial bioactive compounds are outlined in the chapter along with commercial products currently available on the market.

Keywords: agricultural applications, bioactive compounds, biofuels, bioplastics, cosmetics, cyanobacteria, environmental remediation, pharmaceuticals

1. Introduction

Cyanobacteria are one of the oldest organisms on earth, based on the evidence of 3.5 billion year-old fossilized records [1]. These ancient organisms are ubiquitous photoautotrophic microorganisms found in fresh, brackish, marine or wastewater [2]. They are able to adapt to a wide range of conditions (salinities, temperatures, pH factors, light intensities, and so on). Oxygenic photosynthesis is believed to have started with these microbes, making them a big contributor to the oxygen-rich atmosphere we enjoy today. Cyanobacteria are by far, the only known prokaryotes that perform oxygen-evolving photosynthesis [3] and the concentration of CO₂ in the atmosphere would be twice as high had it not been for cyanobacteria [4]. Thus, cyanobacteria have helped to shape the evolutionary trajectory of the earth. It has been suggested that the oil-producing gene in plant chloroplasts originated from cyanobacteria. According to [5] a primordial plant cell “engulfed” a cyanobacterium about a billion years ago. The bacterium lived in the cell and supplied it with photosynthetic products. Thus, implying that the oil synthesis enzyme (acyltransferase) of

the chloroplasts originated from cyanobacteria. Similarly, according to [6] ethylene production existed before land plants colonized the earth as evidenced by unambiguously homologous ethylene-signaling pathways in *Spirogyra* and *Arabidopsis*; implying that cell elongation was possibly an ancestral ethylene production response.

Nearly 2,700 species of cyanobacteria have been described, however, prediction models suggest that between 2,000 and 8,000 cyanobacteria species exist in nature [7]. Some of the described species have been studied and known to have potential applications in agriculture, energy, food and pharmaceutical industries because of their ability to produce oil, fix atmospheric nitrogen, and also have high vitamin, mineral and protein contents among others [8].

Eutrophic water bodies facilitate the proliferation of cyanobacteria cells into blooms. Some of the blooms consist of cyanotoxin-producing species. The cyanotoxins are poisonous to humankind, animals and aquatic life. This has led to the closure of recreational centres and caused huge economic losses [9]. It is, therefore, imperative to understand the actions of these metabolites, their biosynthesis and possible applications in industry. This review will discuss the mechanisms involved in the cellular production of important metabolites and address the far-reaching industrial applications of these cyanobacteria compounds. Commercial products and companies producing cyanobacteria products are highlighted.

2. Biology of cyanobacteria

The composition of the earth's earlier environment may have influenced the evolution of the metal resistance features and the metal-utilizing proteins in cyanobacteria during a period when the atmosphere was limited in oxygen. Thus, some cyanobacteria still thrive in low oxygen conditions and some strains are highly tolerant to free sulfide [10]. Additionally, cyanobacteria have a high tolerance to ultraviolet-B and -C radiations along with high temperature tolerance, as high as 73°C [10]. Their tolerance to radiation is likely to have been crucial in the early evolution of the cyanobacteria.

Morphologically, cyanobacteria cells are coccoid, filamentous (filamentous non-heterocyst forming, and heterocyst forming) forms [10, 11]. Cyanobacterial cells have an outermost peptidoglycan layer composed mainly of proteins and lipopolysaccharides. Other minor components of the peptidoglycan cell wall include carotenoids and lipids that enhance permeability, mechanical stability and resistance toward chemical substances [12]. Despite their overall gram-negative structure, the peptidoglycan layer in cyanobacteria is significantly thicker than that of most gram-negative bacteria [12]. Outside the cell wall is a carbohydrate-rich glycocalyx with varying proportion of three distinct layers; a closely associated sheath, a well-defined capsule, and loosely attached slime [13]. These three layers protect cyanobacteria cells from desiccation and possibly from predators and phages, making their applications in biotechnology more feasible due to their robust nature.

Cyanobacteria cells divide by fission (binary or multiple), fragmentation or spore-formation processes. The doubling time of cells can occur between 2.1 hours and 72 hours under optimal conditions and as long as 10,000 years under stress conditions such as *Chroococcidiopsis* in dry deserts of Antarctica [10]. Their presence in such dry conditions confirms their ability to tolerate desiccation and water stress, allowing their use in biological processes under arid conditions. The fast doubling time of cyanobacteria cells leads to rapid growth rate which makes their application in industry economically sustainable.

In addition to their fast growth rate, cyanobacteria possess superior photosynthetic capabilities that enables them to convert about 10% of the solar energy received, into biomass which is relative to the 1% conversion done by traditional energy crops like sugarcane or corn, grown for biofuel production [14]. Some cyanobacteria cells possess gas-filled cavities that allow them to float on water surfaces, enhancing light capture for better photosynthetic efficiency. Internal thylakoid membranes are the site of photosynthetic reactions in cyanobacteria. The presence of chlorophyll and phycobilins increase the capture and conversion of light energy during photosynthesis. Other pigments including xanthophylls, carotenes, c-phycoerythrin and c-phycocyanin are present in cyanobacteria. C-phycoerythrin and c-phycocyanin are unique to blue-green algae.

Circadian rhythms are very important in cyanobacteria. These rhythms are fundamental adaptations to the earth's daily light and temperature fluctuations that lead to proper metabolic activity [15]. For example, nitrogen fixation is common in many species of cyanobacteria. However, oxygenic photosynthesis and nitrogen fixation are discordant processes because the nitrogenase enzyme is inactivated by oxygen. Two mechanisms are used to separate these activities. First, a biological circadian clock separates them temporally and cellular differentiation separates them spatially. For instance, a unicellular species such as *Cyanothece* strain ATCC 51142 stores glycogen at daytime and fixes nitrogen at night [15], whereas the filamentous strain, *Trichodesmium erythraeum* IMS101 fixes nitrogen during the day in groups of specialized cells known as heterocysts [16]. Based on this, heterocyst-forming cyanobacteria are able to differentiate highly specialized cells to provide fixed nitrogen to the vegetative cells in a filament. This property allows cyanobacteria to be used as bio-fertilizers in agriculture because of their ability to fix atmospheric nitrogen [14].

Fluctuation in environmental conditions causes oxidative stress to cells which inevitably lead to increased reactive oxygen species (ROS) production. As a form of defense, photosynthetic microorganisms including cyanobacteria have developed several mechanisms to evade the negative effects of ROS. Antioxidants and polyunsaturated fatty acids (PUFAs) are known to provide protection to the cell against oxidative stress by stabilizing free radicals [17]. Among the many fatty acids, PUFAs are of great interest due to their numerous health benefits and increasing global demand [18]. Additionally, they help regulate various cellular processes such as oxygen and electron transport, membrane fluidity, and heat adaptation [19]. Cyanobacteria lack organelles such as endoplasmic reticulum, mitochondria, chloroplasts and Golgi apparatus typically found in eukaryotic cells. However, ribonucleic acid (RNA)-containing organelles known as ribosomes are widespread in cyanobacterial cells. Ribosomes are responsible for protein synthesis that enables cells to perform efficient metabolic activities.

3. Metabolic activities and the production of bioactive compounds

Cellular metabolic processes lead to the production of valuable primary and secondary metabolites. Photosynthesis and carotenogenesis are examples of metabolic processes that lead to the production of primary metabolites like lipids (e.g., PUFAs), antioxidants (e.g., carotenoids) and some proteins (e.g., primary proteins). Primary metabolites are directly involved in normal developmental processes such as cell division, growth and reproduction [20]; and can also be transformed into products such as bio-fertilizers, bio-plastics, nutrient supplements, and dyes. These are beneficial in industrial applications (**Figure 1**). Secondary

Figure 1.

Industrial applications of cyanobacteria from laboratory to up-scaled production and the extraction of metabolites of commercial value for the production of pharmaceuticals, biofertilizers, biofuels, biopolymers and cosmetics.

metabolites on the other hand, are not utilized by the cells for their primary needs. These include hormonal compounds and antibiotics, or toxins [21].

Cyanobacteria possess a relatively simple genome [14], making it easy for gene modification and manipulations for the exploration of novel metabolites. From a nutritional perspective, microbes can alter their cellular metabolism naturally through stress response such as nutrient deficiency [22]. Growth conditions can be manipulated to promote the production of biomass rich in valuable secondary metabolites of economic value. This is usually achieved through a two-stage culture technique where in the first step the cells are grown under optimal conditions to maximize biomass production. The second step involves the introduction of stress factors, such as nutrient deprivation or high light intensity, to induce the production of valuable secondary metabolites. Although a deficiency in nitrogen is known to inhibit cell cycle processes and the production of several cellular components, surprisingly, lipid and biopolymer (PHB) syntheses rather increase under these conditions [23]. This leads to the accumulation of oil droplets and starch granules in starved cells [22, 24]. These adaptive responses help to ensure the cell's survival during stressful conditions. Both metabolites (lipids and starch granules) serve as energy stores coupled with the special role of the starch granules in osmotic balance, heat, freezing, and ultraviolet (UV) rays [22, 23].

The methylerythritol 4-phosphate (MEP) pathway, known to occur in algae, bacteria, and plants, is a classic example of a metabolic pathway that is exploited for drug discoveries. In cyanobacteria, the MEP pathway leads to chlorophyll and hormone production [17]. It is observed that secondary metabolite production is species-specific and environmental conditions can influence their production. For instance, microalgae growing under stress conditions are more likely to produce secondary metabolites with antibacterial activity. In cyanobacteria, fatty acid synthesis (FAS) is performed by a type II fatty acid synthase complex and these

fatty acids have anticarcinogenic, antibiotic, antifungal, and antiviral [25] properties that facilitates their use in pharmaceutical applications.

The shikimate pathway has been proposed to be responsible for mycosporine-like amino acids (MAAs) and scytonemin biosynthesis [26, 27]. These compounds have the potential to be used as natural UV blockers in product formulations such as cosmetic creams or paints, and varnishes (**Figure 1**).

4. Industrial applications of bioactive compounds

4.1 Bio-plastics

Cyanobacteria, have the potential to produce renewable biopolymers from natural resources such as, solar energy, water and CO₂, reducing the need for fertile soils, fertilizers, herbicides and potable water for crop production. Biodegradable polymers such as PHAs are produced as inclusion bodies within cells via the beta-oxidation pathway [28]. *Arthrospira* (*Spirulina*), *Synechococcus*, and *Synechocystis*, are examples of cyanobacteria species widely employed in biopolymer production [23, 29]. Currently majority of commercial bioplastics produced from biologically-derived polymers are sourced from first generation feedstock through fermentation processes. Companies like **Tianjin GreenBio** (China), **Metabolix** (USA), **Biocycle** (Brazil) and **Polyferm** (Canada) produce biopolymers from sugars and vegetable oils. Some biopolymers are also produced by microorganisms like methanotrophs as in the case of **Mango Materials** (USA). Although the potential for biopolymer production from cyanobacteria exists, commercialization on an industrial scale is yet to be achieved. Research is currently geared towards optimization of cultivation and genetic modification approaches for enhanced biopolymer production [23, 30, 31].

4.2 Agricultural applications and environmental remediation

Cyanobacteria are known to play a key role in maintaining the stability of the surface crusts of dry lands. Thus, to combat desertification, cyanobacteria can be used in conjunction with bacteria, algae, mosses, lichens, or fungi, which form the biological soil crusts in unique geographical regions [32]. These biological soil crusts help primary succession in arid regions by improving the nutrient and moisture contents [32]. Cyanobacteria help to form a complex of heavy metals and xenobiotics to limit their mobility and transport in plants. Additionally, they offer protection to plants from disease-carrying insects, act as bio-control agents as well as enhance the mineralization of simpler organic molecules for easy assimilation. Additionally, cyanobacteria have been extensively applied in the area of environmental bioremediation through wastewater treatment processes [33, 34].

Arthrospira maxima and *A. platensis*, generically referred to as *Spirulina* are widely used as dietary supplements in feed for the poultry and aquaculture industries for their nutritional benefits due to their protein, vitamin and fatty acid content. Species such as *Anabaena*, *Aulosira*, *Calothrix*, *Nostoc*, and *Plectonema* are also used for agricultural purposes because of their nitrogen-fixing capabilities. They are able to fix atmospheric nitrogen (N₂) by the conversion of nitrogen to ammonia (N₂ + 3H₂ → 2NH₃) also reducing soil salinity and controlling weed growth [35, 36]. Additionally, they increase soil phosphates [32] by converting insoluble phosphorus in the soil to phytoavailable forms [37], making them an excellent choice as bio-fertilizers. Studies have shown that endophytic cyanobacteria strains such as *Nostoc* strains have the ability to produce phytohormones,

indole-3-acetic acid and cytokinins in root cells of both rice and wheat for their growth and development [38]. In Asian regions for instance, *Azolla* is either mixed into soil prior to rice planting or grown as a mixed crop along with rice in rice fields due to the presence of a cyanobacterium endosymbiont, *Anabaena azollae*, which fixes nitrogen [39].

Cyanobacteria also have other agriculture potential. Some cyanotoxins demonstrate biocidal activity. These biocides inhibit the growth of microorganisms such as viruses, bacteria and fungi; they also affect invertebrates including crustaceans, bivalves and vertebrates such as fish, birds, and mammals [40–44]. Thus, cyanobacteria toxins could be developed into active biological compounds and applied in crop fields as algaecides, fungicides, herbicides and insecticides because of their allelopathic effects [17, 43, 44]. Biocides have low environmental risks and are thus preferable to synthetic pesticides negatively affect the environment [17].

4.3 Biofuels

Our reliance on petroleum products has resulted in polluting the environment. Aside the release of toxic fumes including green-house gasses, into the atmosphere, the process of obtaining the fuel in itself presents potential environmental hazards. Cyanobacteria hold great promise as sources of renewable by-products (biodiesel) especially for the energy sector [45]. Cyanobacteria cells can be engineered to convert CO₂ and water into biofuels through photosynthesis. In a study [46] four modules were optimized to achieve high titer values (4.8 g/L⁻¹) of a petroleum substitute, 1-butanol. Firstly, 1-butanol biosynthesis was introduced and re-cast by systematic screening of genes and pathways. Module 2 involved the optimization of the 5'-regions of expression units to tune protein expression levels. Module 3 rewired carbon flux by editing acetate metabolism. In module 4, photosynthetic central carbon metabolism was rewritten by installing a phosphoketolase (PK) pathway. Several other biofuels have been produced from engineered cyanobacteria e.g., acetone, 2,3-butanediol ethanol, ethylene, isobutanol, 2-methyl-1-butanol from *Synechocystis* sp. [47]. **Algenol**, and **JouleUnlimited** are USA-based companies that use genetically modified cyanobacteria capable of growing in brackish or saltwater to produce a range of biofuels such as ethanol, biodiesel, gasoline and jet fuel in addition to other valuable chemicals [48–50]. Their ability to grow in saltwater makes their industrial application more economically and environmentally sustainable as it reduces the burden of using limited freshwater resources for their cultivation.

4.4 Cosmetics

Numerous cyanobacteria species have been used in the cosmetics industry for many decades because of their anti-inflammatory, antioxidant, and detoxifying properties [15, 17]. The cosmetic industry has evolved from just topical skin products to a more invasive approach of beautifying from within. This booming industry seeks to address skin-related issues by resolving internal problems at the cellular level. Skin aging, wrinkling, drying and other skin conditions occur as a result of loss of elasticity to the skin. In the cosmetic industry, for instance, a novel extract, extracellular polysaccharide (EPS), extracted from *Pseudomonas fluorescens* PGM37 has been found to have higher moisturizing retention ability and has the potential to be used in cosmetics and medicinal products [46]. Similarly, sacran a known cyanobacteria gel extracted from *Aphanothece sacrum* can be used as a moisturizing and anti-inflammatory agent [51, 52].

Again, microalgae strains with high amino acid content are great for improving skin texture and elasticity. Other strains rich in lipids help soothe and moisturize skin tissues, while antioxidant-rich algae with chlorophyll are ideal for detoxifying. Many *Spirulina*-infused cosmetic products are already on the market in the form of tablets, lotions and facial masks. Skin perfection has garnered so much attention in recent years. For instance, skin whitening has become a common practice world-wide, with a booming market in Asia and Africa [47]. Tyrosinase inhibition is the most common approach to achieving skin hypo-pigmentation as this enzyme catalyzes the rate-limiting step of pigmentation. Tyrosinase inhibitors have been isolated from numerous marine macroalgae species such as *Ecklonia cava*, *Laminaria japonica* and *Sargassum silquastrum* [47]. Oscillapeptin G, is an example of a tyrosinase inhibitor isolated from toxic cyanobacteria, *Oscillatoria agardhii* [48]. There is a huge potential for commercialization of cyanobacteria since their cells grow faster than that of marine macroalgae and thus would be more economically efficient at industrial scale.

4.5 Food

Food supplements, animal feed, food additives, and colorants produced from cyanobacterial carotenoids such as canthaxanthin, beta-carotene, nostoxanthin, and zeaxanthin are on the rise. *Spirulina* is a cyanobacterial specie rich in riboflavin, thiamine, β -carotene, and vitamin B₁₂. These supplements are sold as capsules, granules and tablets on the market. A carotenoid such as the ketocarotenoid, astaxanthin is known to be a more powerful antioxidant than vitamin C and A or other carotenoids which play a vital role in preventing damage in human cells through photooxidation [17]. Astaxanthin obtained from *Haematococcus pluvialis* contains protease inhibitors that may be used to treat diseases, such as HIV [17]. These food supplements are usually made from cyanobacterial biomass and consumed whole unlike extracts used in the production of pharmaceuticals [53].

4.6 Pharmaceuticals

Natural products have become important contributing sources of semi-synthetic and synthetic drugs in all major disease fields; predominantly in antibiotic therapies, immunoregulation and oncology [54–56]. Most of these bio-medical natural products or metabolites have been derived from cyanobacteria. These cyanobacterial metabolites have exhibited both interesting and exciting biological activities including antibacterial, anticancer, antifungal, antimicrobial, and antiviral activities. Others are anticoagulant, anti-HIV, anti-inflammatory, anti-malarial, antiprotozoal, antituberculosis, antitumor, and immunosuppressant activities [54, 55]. Some of these bioactive compounds are Borophycin from *Nostoc* sp. against human carcinoma, Calothrix from *Calothrix* sp. against human HELa cancer cells and inhibition of growth of chloroquine-resistant strain of malaria parasite *Plasmodium falciparum* [8, 54]. Extracts from *Lyngbya lagerhaimanni* has anti-HIV activity, whilst *Lyngbyatoxin A* from toxic strains of *Lyngbya majuscula* is highly inflammatory [56, 57].

4.7 Bio-pigments

Phycobilisomes are phycobiliproteins accumulated by cyanobacteria and these include phycocyanin (blue), phycoerythrin (red), and allophycocyanin (blue-gray). They are major light-harvesting complexes in cyanobacteria [58, 59]. These compounds are used as bio-pigments in industrial applications. *Linablue*®

is an example of such bio-pigments approved by the USFDA [60] for use in food products. After extraction of phycocyanin the waste biomass is used as the starting material for the production of black ink, *Algae Black*. In the past, charcoal and carbon-derived soot were the ingredients required to make black ink. These petrochemically-derived materials generated a significant carbon footprint, and listed as a class 2b carcinogen [61]. Currently, however, a product extracted from *Spirulina* is used as the base material to create the renewable, bio-based carbon black ink known as *AlgaeBlack* by the **Living Ink** company whose aim is to create a sustainable alternative to traditional carbon-black ink pigment. A summary of major cyanobacteria product-related companies along with their founding years are provided in **Table 1**. Chlorophylls for instance have been used as a textile dye with antimicrobial properties and also as a biomordant to enhance dyeing processes in textile production [8].

4.8 Research and development

Other applications of cyanobacterial extracts include their use in scientific research experiments. For instance, phycobiliproteins have fluorescent properties that can be used in flow cytometry and in immunoassay techniques [61]. Among these numerous bioactive compounds are terpenoids such as terpenes, diterpenes

Company/Country (founding year)	Focus area	Industrial applications	Reference
LG Sonic/Netherlands (1999)	Algal blooms	Effective ultrasonic frequencies for algae control on large water surface areas	[62]
Cyano Biotech/Germany (2004)	Development of novel cyanobacteria products	Discovery and development of novel structures based on cyanobacterial natural products	[63]
Photanol/Netherlands (2008)	Genetic modification of cyanobacteria	Produces broad range of biochemicals from cyanobacteria	[64]
Algae Biotechnologia/Brazil (2009)	Technological development of microalgae and cyanobacteria cultivation systems	Treatment of liquid and gaseous agro-industrial effluents. Production of ingredients and additives for animal nutrition. Production of human food supplements and production of biofuels.	[65]
Algenuity/United Kingdom (2009)	Synthetic biotechnology	Produces lab scale photobioreactor for algae and cyanobacteria research and also harness specific microalgal strains for synthetic biology applications	[66]
Living Ink Technology/USA (2013)	Production of ecofriendly ink products	Develops a variety of ink products and colors, including digital ink	[67]
Spira Inc./USA (2016)	Incorporation of algae-based ingredients into everyday products	Extract high-value compounds from algae.	[68]

Table 1.
Major industrial companies involved in the production of cyanobacterial metabolites.

and sesquiterpenes. These organic compounds are widely found in cyanobacteria and are used as natural ingredients in flavors and perfumes. Such applications are recently gaining grounds in therapeutic and pesticide industries [17]. Terpenoid type compounds such as carotenoids and phytols are crucial for chlorophyll and hormone biosynthesis through the methylerythritol 4-phosphate (MEP) pathway in cyanobacteria that was discussed in Section 3.

The rapid development of molecular tools for whole genome sequences promotes the use of omics technology i.e. transcriptomics, proteomics, and system biology approaches to manipulate metabolic pathways for producing valuable products [8].

Experimental approaches that require fluorescent probes such as fluorescence microscopy for diagnostics and biomedical research utilizes the autofluorescent properties of cyanobacteria pigments such as phycobiliproteins. Among these, the most widely exploited fluorescent probe is phycoerythrin utilized in biomedical research [8].

5. Conclusions

Bioactive compounds synthesized by cyanobacteria are innumerable coupled with their vast industrial applications. More of these natural compounds are being discovered on a regular basis through research and development. There also exists an untapped pool of bioactive compounds that genetic engineering techniques can unfold. Current process engineering strategies in cyanobacteria research is centered on the regulation of metabolic pathways for the production of bioactive compounds. Such exploitations are feasible due to the small and simple genome of cyanobacteria species. Considering the fact that cyanobacteria are a promising feedstock for a circular economy, it is important to develop robust strains suitable for industrial applications.

The possibility of producing novel biopolymer blends, biofuel components, and pharmaceutical compounds that are capable of meeting the demands of a biotechnologically-driven society is vast. Thus, cyanobacteria will continue to play crucial roles in terms of health, energy, food, and many other aspects of our lives.

To achieve great strides in the cyanobacteria production sector, a synergistic approach should be adopted by cyanobacteria-related companies. Collaboratively, companies dealing in products such as biofuels, and bioplastics that involve extraction of compounds from cyanobacteria cell biomass, could supply their waste biomass to other companies that require the waste biomass as raw materials for production, such as algae ink or biochar for wastewater treatment. Thus, forming a biorefinery supply chain network that will essentially benefit both the producers and consumers from an economic and environmental point of view. Cyanobacteria industrial harmonization will ensure the valorization of production processes.

Conflict of interest

The authors declare no conflict of interest.

IntechOpen

Author details

Ayesha Algade Amadu¹, Kweku Amoako Atta deGraft-Johnson²
and Gabriel Komla Ameka^{3*}

1 Council for Scientific and Industrial Research – Water Research Institute, Accra, Ghana

2 CSIR-College of Science and Technology (CCST), Accra, Ghana

3 Department of Plant and Environmental Biology, University of Ghana, Legon, Accra, Ghana

*Address all correspondence to: gameka@ug.edu.gh

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Schopf J., Packer BM. Early Archean (3.3-Billion to 3.5-Billion-Year-Old) Microfossils from Warrawoona Group, Australia. *Science* (80-). 1987;237:70-3.
- [2] Costa JAV, Moreira JB, Lucas BF, Da Silva Braga V, Cassuriaga APA, De Moraes MG. Recent advances and future perspectives of PHB production by cyanobacteria. *Ind Biotechnol*. 2018; 14(5):249-56.
- [3] Rasmussen B, Fletcher IR, Brocks JJ, Kilburn MR. Reassessing the first appearance of eukaryotes and cyanobacteria. *Nature*. 2008;455(7216): 1101-4.
- [4] ScienceDaily. Unexpected discovery: Blue-green algae produce oil [Internet]. 2020. [cited 2021 Jan 15]. Available from: <https://www.sciencedaily.com/releases/2020/03/200305132125.htm#:~:text=Cyanobacteria--colloquially also called blue-green algae--can,by the University of Bonn>.
- [5] Aizouq M, Peisker H, Gutbrod K, Melzer M, Hölzl G, Dörmann P. Triacylglycerol and phytyl ester synthesis in *Synechocystis* sp. PCC6803. *Proc Natl Acad Sci U S A*. 2020;117(11):6216-22.
- [6] Ju C, Van De Poel B, Cooper ED, Thierer JH, Gibbons TR, Delwiche CF, et al. Conservation of ethylene as a plant hormone over 450 million years of evolution. *Nat Plants* [Internet]. 2015;1(1):1-7. Available from: <http://dx.doi.org/10.1038/nplants.2014.4>
- [7] Nabout JC, da Silva Rocha B, Carneiro FM, Sant'Anna CL. How many species of Cyanobacteria are there? Using a discovery curve to predict the species number. *Biodivers Conserv*. 2013;22(12):2907-18.
- [8] Singh PK, Kumar A, Kumar SV, Shrivastava AK, editors. *Advances in Cyanobacterial Biology*. Academic Press Inc.; 2020.
- [9] WHO. Algae and cyanobacteria in fresh water. In: *Guidelines for safe recreational water environments*. 1999. p. 136-58.
- [10] Whitton BA, Potts M. Introduction to the cyanobacteria. Vol. 9789400738, *Ecology of Cyanobacteria II: Their Diversity in Space and Time*. 2012. p. 1-13.
- [11] Whitton BA, Potts M. The Ecology of Cyanobacteria [Internet]. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*. Kluwer Academic Publishers; 2000. Available from: <http://publications.lib.chalmers.se/records/fulltext/245180/245180.pdf%0Ahttps://hdl.handle.net/20.500.12380/245180%0Ahttp://dx.doi.org/10.1016/j.jsames.2011.03.003%0Ahttps://doi.org/10.1016/j.gr.2017.08.001%0Ahttp://dx.doi.org/10.1016/j.precamres.2014.12>
- [12] Hoiczky E, Hansel A. Cyanobacterial cell walls: News from an unusual prokaryotic envelope. *J Bacteriol*. 2000;182(5):1191-9.
- [13] Kumar K, Mella-Herrera RA, Golden JW. Cyanobacterial heterocysts. *Cold Spring Harb Perspect Biol*. 2010;2(4):1-19.
- [14] Zahra Z, Choo DH, Lee H, Parveen A. Cyanobacteria: Review of current potentials and applications. *Environ - MDPI*. 2020;7(2).
- [15] Toepel J, Welsh E, Summerfield TC, Pakrasi HB, Sherman LA. Differential transcriptional analysis of the cyanobacterium *Cyanothece* sp. strain ATCC 51142 during light-dark and continuous-light growth. *J Bacteriol*. 2008;190(11):3904-13.

- [16] Sandh G, El-Shehawy R, Díez B, Bergman B. Temporal separation of cell division and diazotrophy in the marine diazotrophic cyanobacterium *Trichodesmium erythraeum* IMS101. FEMS Microbiol Lett. 2009;295(2): 281-8.
- [17] Singh R, Parihar P, Singh M, Bajguz A, Kumar J, Singh S, et al. Uncovering potential applications of cyanobacteria and algal metabolites in biology, agriculture and medicine: Current status and future prospects. Front Microbiol. 2017;8(APR):1-37.
- [18] Steinhoff FS, Karlberg M, Graeve M, Wulff A. Cyanobacteria in Scandinavian coastal waters - A potential source for biofuels and fatty acids? Algal Res [Internet]. 2014;5(1):42-51. Available from: <http://dx.doi.org/10.1016/j.algal.2014.05.005>
- [19] Cardozo KHM, Guaratini T, Barros MP, Falcão VR, Tonon AP, Lopes NP, et al. Metabolites from algae with economical impact. Vol. 146, Comparative Biochemistry and Physiology - C Toxicology and Pharmacology. Elsevier Inc.; 2007. p. 60-78.
- [20] Martínez-Francés E, Escudero-Oñate C. Cyanobacteria and Microalgae in the Production of Valuable Bioactive Compounds. In: IntechOpen [Internet]. 2018. p. 105-28. Available from: <http://dx.doi.org/10.1039/C7RA00172J%0Ahttps://www.intechopen.com/books/advanced-biometric-technologies/liveness-detection-in-biometrics%0Ahttp://dx.doi.org/10.1016/j.colsurfa.2011.12.014>
- [21] Carmichael WW. Cyanobacteria secondary metabolites—the cyanotoxins. J Appl Bacteriol. 1992;72(6):445-59.
- [22] Chen B, Wan C, Mehmood MA, Chang JS, Bai F, Zhao X. Manipulating environmental stresses and stress tolerance of microalgae for enhanced production of lipids and value-added products—A review. Bioresour Technol [Internet]. 2017;244:1198-206. Available from: <http://dx.doi.org/10.1016/j.biortech.2017.05.170>
- [23] Amadu AA, Qiu S, Ge S, Addico GND, Ameka GK, Yu Z, et al. A review of biopolymer (Poly- β -hydroxybutyrate) synthesis in microbes cultivated on wastewater. Sci Total Environ [Internet]. 2021;756:143729. Available from: <https://doi.org/10.1016/j.scitotenv.2020.143729>
- [24] Ge S, Champagne P, Plaxton WC, Leite GB, Marazzi F. Microalgal cultivation with waste streams and metabolic constraints to triacyl glycerides accumulation for biofuel production. Biofuels, Bioprod Biorefining. 2017;11(4):325-43.
- [25] El-Baz FK, El-Senousy WM, El-Sayed AB, Kamel MM. In vitro antiviral and antimicrobial activities of *Spirulina platensis* extract. J Appl Pharm Sci. 2013;3(12):52-6.
- [26] Soule T, Palmer K, Gao Q, Potrafka RM, Stout V, Garcia-Pichel F. A comparative genomics approach to understanding the biosynthesis of the sunscreen scytonemin in cyanobacteria. BMC Genomics. 2009;10:1-10.
- [27] Rosic NN. Mycosporine-like amino acids: Making the foundation for organic personalised sunscreens. Mar Drugs. 2019;17(11):1-17.
- [28] Ranaivoarisoa TO, Singh R, Rengasamy K, Guzman MS, Bose A. Towards sustainable bioplastic production using the photoautotrophic bacterium *Rhodopseudomonas palustris* TIE-1. J Ind Microbiol Biotechnol [Internet]. 2019;46(9-10):1401-17. Available from: <https://doi.org/10.1007/s10295-019-02165-7>
- [29] Kamravamanesh D, Lackner M, Herwig C. Bioprocess Engineering

Aspects of Sustainable Polyhydroxy alkananoate Production in Cyanobacteria. *Bioengineering*. 2018;5(4):111-29.

[30] Jiang XR, Wang H, Shen R, Chen GQ. Engineering the bacterial shapes for enhanced inclusion bodies accumulation. *Metab Eng [Internet]*. 2015;29:227-37. Available from: <http://dx.doi.org/10.1016/j.ymben.2015.03.017>

[31] Carpine R, Olivieri G, Hellingwerf KJ, Pollio A, Marzocchi A. Industrial production of poly- β -hydroxybutyrate from CO₂: Can cyanobacteria meet this challenge? *Processes*. 2020;8(3):1-23.

[32] Rossi F, Li H, Liu Y, De Philippis R. Cyanobacterial inoculation (cyanobacterisation): Perspectives for the development of a standardized multifunctional technology for soil fertilization and desertification reversal. *Earth-Science Rev [Internet]*. 2017;171:28-43. Available from: <http://dx.doi.org/10.1016/j.earscirev.2017.05.006>

[33] Singh JS, Kumar A, Rai AN, Singh DP. Cyanobacteria: A precious bio-resource in agriculture, ecosystem, and environmental sustainability. *Front Microbiol*. 2016;7(APR):1-19.

[34] Martins J, Peixe L, Vasconcelos VM. Unraveling Cyanobacteria Ecology in Wastewater Treatment Plants (WWTP). *Microb Ecol*. 2011;62(2):241-56.

[35] Jochum M, Moncayo LP, Jo Y. Microalgal cultivation for biofertilization in rice plants using a vertical semi-closed airlift photobioreactor. *PLoS One*. 2018;13(9):1-13.

[36] Shweta S, Samuel K. Study of Cyanobacteria as biofertilizer from the rice field. *world J Pharm Res*. 2015;4(3):1696-706.

[37] Sahu D, Priyadarshani I, Rath B. Cyanobacteria -As Potential Biofertilizer. *CIBTech J Microbiol* Jul-Sept Oct-Dec. 2012;1(3):2319-386720.

[38] Hussain A, Hasnain S. Phytostimulation and biofertilization in wheat by cyanobacteria. *J Ind Microbiol Biotechnol*. 2011;38(1):85-92.

[39] Bocchi S, Malgioglio A. Azolla-Anabaena as a Biofertilizer for Rice Paddy Fields in the Po Valley , a Temperate Rice Area in Northern Italy. 2010;2010.

[40] Misra S, Kushik D. Growth Promoting Substances of Cyanobacteria. II. Detection of Amino Acids, Sugars and Auxins. *Proc Indian natn Sc Acad [Internet]*. 1989;B55(5&6):499-504. Available from: http://www.insa.nic.in/writereaddata/UpLoadedFiles/PINSA/Vol55B_1989_5and6_Art29.pdf

[41] Schwartz RE, Hirsch CF, Sesin DF, Flor JE, Chartrain M, Fromtling RE, et al. Pharmaceuticals from cultured algae. *J Ind Microbiol*. 1990;5(2-3):113-23.

[42] Burja AM, Banaigs B, Abou-Mansour E, Grant Burgess J, Wright PC. Marine cyanobacteria - A prolific source of natural products. *Tetrahedron*. 2001;57(46):9347-77.

[43] Berry JP. Cyanobacterial Toxins as Allelochemicals with Potential Applications as Algaecides, Herbicides and Insecticides. *Mar Drugs*. 2008;6(2):117-46.

[44] Rastogi RP, Sinha RP. Biotechnological and industrial significance of cyanobacterial secondary metabolites. *Biotechnol Adv [Internet]*. 2009;27(4):521-39. Available from: <http://dx.doi.org/10.1016/j.biotechadv.2009.04.009>

[45] Doamekpor LK, Onwona-Agyeman R, Ameka GK. Bioenergy :

Biodiesel from Freshwater Cyanobacterium Occurring in Ghana Green Microalgae and a. West African J Appl Ecol. 2019;27(2):51-60.

[46] Liu X, Miao R, Lindberg P, Lindblad P. Modular engineering for efficient photosynthetic biosynthesis of 1-butanol from CO₂ in cyanobacteria. Energy Environ Sci. 2019;12(9): 2765-77.

[47] Nozzi NE, Oliver JWK, Atsumi S. Cyanobacteria as a Platform for Biofuel Production. Front Bioeng Biotechnol. 2013;1(August).

[48] Mazard S, Penesyan A, Ostrowski M, Paulsen TI, Egan S. Tiny Microbes with a Big Impact : The Role of Cyanobacteria and Their Metabolites in Shaping Our Future. Mar Drugs. 2016;14(97):1-19.

[49] Algenol Biotech [Internet]. [cited 2021 Feb 14]. Available from: www.algenol.com

[50] Joule Unlimited [Internet]. [cited 2021 Feb 14]. Available from: <http://www.jouleunlimited.com>

[51] Morone J, Alfeus A, Vasconcelos V, Martins R. Revealing the potential of cyanobacteria in cosmetics and cosmeceuticals — A new bioactive approach. Algal Res [Internet]. 2019;41(May):101541. Available from: <https://doi.org/10.1016/j.algal.2019.101541>

[52] Okeyoshi K, Okajima MK, Kaneko T. The cyanobacterial polysaccharide sacran: characteristics, structures, and preparation of LC gels. Polym J [Internet]. 2021;53(1):81-91. Available from: <http://dx.doi.org/10.1038/s41428-020-00426-2>

[53] Chronakis IS, Madsen M. Algal proteins. In: Handbook of food proteins [Internet]. Woodhead Publishing Limited; 2011. p. 353-94. Available from:

<http://dx.doi.org/10.1533/9780857093639.353>

[54] Zainuddin EN, Mundt S, Wegner U, Mentel R. Cyanobacteria a potential source of antiviral substances against influenza virus. Med Microbiol Immunol. 2003;191(3-4):181-2.

[55] Silva T, S. Salomon P, Hamerski L, Walter J, B. Menezes R, Siqueira JE, et al. Inhibitory effect of microalgae and cyanobacteria extracts on influenza virus replication and neuraminidase activity. PeerJ. 2018;6:e5716.

[56] Vijayakumar S, Menakha M. Pharmaceutical applications of cyanobacteria-A review. J Acute Med [Internet]. 2015;5(1):15-23. Available from: <http://dx.doi.org/10.1016/j.jacme.2015.02.004>

[57] Abed RMM, Dobretsov S, Sudesh K. Applications of cyanobacteria in biotechnology. J Appl Microbiol. 2009;106(1):1-12.

[58] Pagels F, Guedes AC, Amaro HM, Kijjoa A, Vasconcelos V. Phycobili proteins from cyanobacteria: Chemistry and biotechnological applications. Biotechnol Adv [Internet]. 2019;37(3):422-43. Available from: <https://doi.org/10.1016/j.biotechadv.2019.02.010>

[59] Telford WG, Moss MW, Morseman JP, Allnut FCT. Cyanobacterial stabilized phycobilisomes as fluorochromes for extracellular antigen detection by flow cytometry. J Immunol Methods. 2001;254(1-2):13-30.

[60] Earthrise. What is Linablue [Internet]. [cited 2021 Jan 11]. Available from: <https://www.earthrise.com/what-is-linablue>

[61] Algae World News. The first algae-based ink is derived from spirulina instead of petrochemicals [Internet].

2019 [cited 2020 Dec 26]. Available from: <https://news.algaeworld.org/2019/12/the-first-algae-based-ink-is-derived-from-spirulina-instead-of-petrochemicals/>

[62] LGSonic. Ultrasonic Algae Control - Treat Large Water Surfaces - LG Sonic [Internet]. Available from: <https://www.lgsonic.com/ultrasonic-algae-control/>

[63] Cyano Biotech GmbH [Internet]. CyanoBiotech. Available from: <http://www.cyano-biotech.com/content/home/index.php>

[64] Photanol. Photanol [Internet]. Available from: <https://photanol.com/>

[65] Biotechnologia A. Algae biotechnologia [Internet]. Available from: <http://www.algae.com.br/site/pt/>

[66] Algenuity. Making algae work [Internet]. Available from: <https://www.algenuity.com/>

[67] LivingInkTech — About Living Ink [Internet]. Available from: <https://livingink.co/about>

[68] Spira. Spira Inc [Internet]. Available from: <https://www.spirainc.com/>