We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Loss of Smell and Taste as Clinical Onset of COVID-19

Nihal Seden

Abstract

Initially, symptoms of COVID-19 associated with Ear-Nose-Throat were thought to be flulike symptoms in the foreground. Such as fever, chills, cough, dyspnoea, myalgia, headache, sore throat. Olfactory and gustatory dysfunction was not a noticeable symptom at first. As the number of cases has risen worldwide, sudden onset hyposmia/anosmia has received increasing attention as a symptom of COVID-19. The reported incidence of anosmia varies internationally: as low as 30% in South Korea, and as high as 88% in Europe. The loss of smell that occurs in COVID-19 infection its general character is sudden onset anosmia. There is currently no specific treatment for COVID-19 related anosmia. Olfactory dysfunction can heal spontaneously. However, not a small number of patients may have permanent impairment.

Keywords: olfactory dysfunction, postviral anosmia, COVID-19 related anosmia, ENT

1. Introduction

Like all healthcare workers, Ear-Nose-Throat (ENT) specialists did not hesitate to take part in the forefront of the epidemic, and investigated the issues where they could benefit both in terms of their expertise and as primary physicians in combating the pandemic.

Initially, symptoms of COVID-19 associated with ENT were thought to be flulike symptoms in the foreground. Such as fever, chills, cough, dyspnoea, myalgia, headache, sore throat, etc. [1]. Olfactory and gustatory dysfunction was not a noticeable symptom at first. In the first studies reported from China, there were no evidence of patients with symptoms of changes and/or loss of smell and taste [2–4]. As the number of cases has risen worldwide, sudden onset hyposmia/anosmia was received increasing attention as a symptom of COVID-19. Due to the efforts of the American Academy of Otolaryngology-Head and Neck Surgery and the British Association of Otorhinolaryngology-Head and Neck Surgery, sudden onset hyposmia and anosmia were accepted as symptoms of COVID-19 by the Centers for Disease Control and Prevention (CDC) and the World Health Organization on 17 April 2020 and 4 May 2020, respectively [5–7]. The reported incidence of anosmia varies internationally: as low as 30% in South Korea, and as high as 88% in Europe [8]. Various hypotheses are on the agenda as to what might cause this difference. One hypothesis is focused on the ethnicity-host factor. A meta-analysis, reported on nearly 40,000 patients across 104 studies found that anosmia (and ageusia) is more prevalent in Caucasians than Asians (54.8 vs. 17.7%, respectively) [9]. In another hypothesis, spike protein mutations - pathogenic factor - are questioned as the cause of the difference in smell loss [10].

In a multicenter European study, a total of 357 patients (85.6%) had olfactory dysfunction related to COVID-19 infection. Among them, 284 (79.6%) patients were anosmic, and 73 (20.4%) were hyposmic. Phantosmia and parosmia were noted in 12.6% and 32.4% of the patients during the disease course, respectively [11]. As we leave behind a year of the pandemic today, the sudden onset of odor loss and taste disturbance are now among the most important ENT-related symptoms, and olfactory disorder is the best predictor of COVID-19 status of all the associated symptoms [12].

2. Pathophysiology

Coronaviruses are known to cause odor loss from previous studies [13]. However, the pathophysiological mechanism of COVID-19 that causes odor and taste disorders has not been fully clarified yet.

2.1 Pathophysiology of gustatory dysfunction

In humans, the sense of taste is carried by three cranial nerves. Facial nerve (7th cranial nerve), glossopharyngeal nerve (9th cranial nerve) and vagus (10th cranial nerve). When the terminal branches are stimulated, the sense of taste reaches to the nucleus solitarius in the brainstem and then it is carried to the thalamus. Hypogeusia can develop through the involvement of one of these three nerves, the nucleus solitarius or tract, or any of the thalamus nuclei. Angiotensin-Converting Enzyme 2 (ACE2) receptors, which allow SARS-CoV-2 to attach to the tissue, are widely expressed in the mucous membrane of the entire oral cavity, especially in the tongue [14–16]. The role of ACE2 in modulating taste perception has been emphasized in many studies analyzing the chemosensitive side effects of ACE2 inhibitors and angiotensin II blockers [16, 17]. Taste disturbance usually regresses after cessation of treatment. Also, a condition recently identified for SARS-CoV-2 is that it can bind to sialic acid receptors [18]. Sialic acid is an essential component of saliva mucin and protects glycoproteins that transport taste molecules into taste pores from early enzymatic degradation [16]. A decrease in sialic acid in saliva is associated with an increase in the threshold of taste [19]. Although it has been suggested that the deterioration in the perception of smell may also cause the loss of taste function due to the close functional link between these two chemosensory systems, the sense of taste seems to be more affected in recent publications.

2.2 Pathophysiology of olfactory dysfunction

Chen et al. reported that ACE2 immunohistochemical expression was 200 to 700 times greater in the sustentacular cells of the olfactory neuroepithelium than nasal or tracheal epithelia [20]. An animal study, showed increase in macrophages in the olfactory epithelium and lamina propria after SARS-CoV-2 infection [21]. Generally, most data indicate that the main targets of SARS-CoV-2 are sustentacular cells in the olfactory epithelium [20–24].

These results show that hyposmia or anosmia is mainly caused by nasal epithelial infection and not a result of general malaise.

3. Diagnosis of olfactory and taste disorders in COVID-19

In many studies on the transmission method of the SARS-CoV-2 virus, it has been shown that this virus is transmitted by droplets. Objective olfactory tests may pose a risk of contamination and therefore require extra precaution.

Commonly used objective olfactory tests are the Connecticut Chemosensory Clinical Research Center (CCCRC) test, University of Pennsylvania Smell Identification test (UPSIT), the Sniffin' Sticks method and the Odor Stick Identification test (OSIT). The UPSIT has four "scratch and sniff" booklets that each contain 10 microcapsule fragrances. After people open the capsule, they smell the page in the booklet [25, 26]. In the CCCRC test, fragrances are offered in bottles that are not transparent [27]. Sniffin' Sticks are felt tip pens impregnated with scents that are handed to the patient to smell [28]. In the OSIT test, the researcher folds a piece of fragrant paraffin paper in half to crush the microcapsule and then offers it to the participant. The participant then opens and smells the paper [29]. For the Sniffin' Sticks and CCCRC tests, the odor threshold is considered along with the ability for odor discrimination, while discrimination alone is assessed in the UPSIT and OSIT.

As for the taste, since taste tests can be performed with disposable strips, they can be used safely in patients with taste disorders.

Many studies have been published on COVID-19 and odor disorders since the beginning of the pandemic, and because of the contamination risk, the vast majority were subjective reports based on questionnaires or self-reports [30–32]. However, in studies that will be done by performing objective odor tests on patients, reports about the importance of obtaining data started to increase; for example Leichen et al. examined 86 patients for anosmia and hyposmia rates by testing with Sniffin' Sticks test, which is a psychophysical odor test. A total of 33 (38%) patients who reported that they had a loss of smell were normosmic according to the Sniffin' Sticks test. In the anosmic group, 78.8% of the patients stated that they had a loss of smell [33]. We had managed to evaluate olfactory objectively, without any risk of contamination, by a method that we described at our study [34].

4. Clinical features

The general feature of loss of smell in COVID-19 infection is sudden onset anosmia. Gane et al., stated in their case series, the isolated sudden onset anosmia syndrome (ISOA), could be the only finding of COVID-19 without any other symptoms [35]. Usually, olfactory disorder is not accompanied by nasal congestion or rhinorrhea.

Olfactory dysfunction due to COVID-19 infection seems to effect females and young individuals more commonly.

Is this odor dysfunction completely reversible, or how long should a patient wait for full recovery of olfactory function? There have been many studies on the alleviation of the loss of smell in COVID-19 [36–45].

Complete recovery of olfactory dysfunction varies between 11-49% and up to 25% of the patients seems to show no improvement at all.

5. Treatment

There is currently no specific treatment for COVID-19 related anosmia. Olfactory dysfunction can heal spontaneously. However, not a small number of patients may have permanent impairment. Efficacy of the treatments are unknown due to lack of data; treatments targeting post-infectious olfactory dysfunction could potentially be beneficial for COVID-19. While the use of systemic steroids is not recommended in patients with odor loss, the use of spray form and long applicators has been found appropriate for nasal steroids [46]. Fragrance therapy gives

successful results in postviral odor loss, especially when it is started early and used for at least three months. Therefore, in the loss of smell associated with COVID-19, fragrance therapy, including rose, lemon, clove and eucalyptus fragrances, each scent is sniffed for twenty seconds twice a day [47].

6. Other ENT related symptoms of COVID-19

In patients without pneumonia, the COVID-19 clinic is similar to some diseases frequently encountered by ENT physicians in the outpatient clinic. COVID-19 can be confused with the flu or common cold in terms of initial symptoms, causing patients to apply to the ENT clinic in the foreground. There may be serous nasal discharge, hyperemic oropharynx, watery eyes and tearing. Since hoarseness is not among the symptoms mentioned, it may be thought that there is no cause of laryngitis or findings similar to laryngitis are not found on examination. Sore throat is a common symptom of COVID-19. Most of pharyngitis or sore throat is of viral origin and these pathogens are predominantly rhinovirus, influenza, adenovirus, coronavirus (previously isolated types) and parainfluenza virus [48].

Other human corona viruses are common cause of viral pharyngitis. Considering that, Sars-CoV-2 is likely to cause pharyngitis and/or similar clinical features.

Both possibilities should be kept in mind in a patient presenting with hyperemia in the oropharynx and difficulty swallowing. Unfortunately, in the current situation, definitive diagnosis distinction can only be made by laboratory testing. The rapid progression and sudden onset of the disease should be thought-provoking for COVID-19.

It should be kept in mind that COVID-19 can also cause rhinitis. It may have similar symptoms with allergic rhinitis. Although fatigue and weakness due to sleep disturbance in allergic rhinitis are reported; muscle pain, malaise and sore throat are not expected. Watery and red eyes may occur in both diseases, but itching in the eyes, palate and nose can be a guide for allergic diseases. Likewise, flu and cold are other common diseases that can be confused with COVID-19. While runny nose is more common in colds than other diseases, in influenza and Sars-CoV-2 infection dry cough is in the foreground.

7. Conclusions

In the fight against COVID-19, it would not be wrong to say that the most important and alarming symptoms specifically to ENT are odor and taste disorders. It should be kept in mind that these deficits do not heal completely in all patients.

Conflict of interest

None.

Thanks

I dedicate this book chapter to all healthcare professionals who work devotedly in combating the epidemic, and especially to colleagues who have lost their lives.

With respect and mercy.

IntechOpen

IntechOpen

Author details

Nihal Seden
Ear Nose Throat Clinic, Istanbul Training and Research Hospital, University of
Health Sciences, Istanbul, Turkey

*Address all correspondence to: nihalseden@hotmail.com

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Symptoms of Coronavirus. <https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/symptoms.html> [Accessed: 2020-03-08]
- [2] Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395(10223):507-513. doi:10.1016/S0140-6736(20)30211-7
- [3] Chen T, Dai Z, Mo P, et al. Clinical Characteristics and Outcomes of Older Patients with Coronavirus Disease 2019 (COVID-19) in Wuhan, China: A Single-Centered, Retrospective Study. *J Gerontol A Biol Sci Med Sci*. 2020;75(9):1788-1795. doi:10.1093/gerona/glaa089
- [4] Guan WJ, Ni ZY, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med*. 2020;382(18):1708-1720. doi:10.1056/NEJMoa2002032
- [5] AAO-HNS. Anosmia, hyposmia, and dysgeusia symptoms of coronavirus disease. <https://www.entnet.org/content/aa-hns-anosmia-hyposmia-and-dysgeusia-symptoms-coronavirus-disease>; [Accessed: 2020-03-22]
- [6] Q&A on coronaviruses (COVID-19) 17 April 2020 Q&A. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/question-and-answers-hub/q-a-detail/q-a-coronaviruses> [Accessed: 2020-04-17]
- [7] Hopkins, C., and Kumar, N. Loss of sense of smell as marker of COVID-19 infection, ENT UK. <https://www.entuk.org/sites/default/files/files/Loss%20of%20sense%20of%20smell%20as%20marker%20of%20COVID.pdf> (accessed 2020-3-20).
- [8] Strauss SB, Lantos JE, Heier LA, Shatzkes DR, Phillips CD. Olfactory Bulb Signal Abnormality in Patients with COVID-19 Who Present with Neurologic Symptoms. *AJNR Am J Neuroradiol*. 2020;41(10):1882-1887. doi:10.3174/ajnr.A6751
- [9] von Bartheld, C. S., Hagen, M. M., and Butowt, R. (2020). Prevalence of chemosensory dysfunction in COVID-19 patients: a systematic review and meta-analysis reveals significant ethnic differences. *ACS Chem. Neurosci*. 11, 2944-2961. doi:10.1021/acscchemneuro.0c00460
- [10] Butowt R, Bilinska K, Von Bartheld CS. Chemosensory Dysfunction in COVID-19: Integration of Genetic and Epidemiological Data Points to D614G Spike Protein Variant as a Contributing Factor. *ACS Chem Neurosci*. 2020;11(20):3180-3184. doi:10.1021/acscchemneuro.0c00596
- [11] Lechien JR, Chiesa-Estomba CM, De Siati DR, et al. Olfactory and gustatory dysfunctions as a clinical presentation of mild-to-moderate forms of the coronavirus disease (COVID-19): a multicenter European study. *Eur Arch Otorhinolaryngol* 2020;277(8):2251-61. <https://doi.org/10.1007/s00405-020-05965-1>.
- [12] Menni C, Valdes AM, Freidin MB, et al. Real-time tracking of self-reported symptoms to predict potential COVID-19. *Nat Med*. 2020;26(7):1037-1040. doi:10.1038/s41591-020-0916-2
- [13] Suzuki M, Saito K, Min Wp, et al. Identification of viruses in patients with postviral olfactory dysfunction. *Laryngoscope*. 2007. doi:10.1097/01.mlg.0000249922.37381.1e.
- [14] Xu H, Zhong L, Deng J, et al. High expression of ACE2 receptor of 2019-nCoV on the epithelial cells of oral mucosa. *Int J Oral Sci*. 2020;12(1):8. Published 2020 Feb 24. doi:10.1038/s41368-020-0074-x

- [15] Vaira LA, Salzano G, Fois AG, Piombino P, De Riu G. Potential pathogenesis of ageusia and anosmia in COVID-19 patients. *Int Forum Allergy Rhinol*. 2020;10(9):1103-1104. doi:10.1002/alr.22593
- [16] Tsuruoka S, Wakaumi M, Araki N, Ioka T, Sugimoto K, Fujimura A. Comparative study of taste disturbance by losartan and perindopril in healthy volunteers. *J Clin Pharmacol*. 2005; 45(11):1319-1323. doi:10.1177/0091270005280445
- [17] Suliburska J, Duda G, Pupek-Musialik D. The influence of hypotensive drugs on the taste sensitivity in patients with primary hypertension. *Acta Pol Pharm*. 2012;69(1):121-127.
- [18] Milanetti M, Miotto M, Di Rienzo L, Monti M, Gosti G, Ruocco G. In-Silico evidence for two receptors based strategy of SARS-CoV-2. *ArXiv*: 2003.11107.
- [19] Pushpass RG, Pellicciotta N, Kelly C, Proctor G, Carpenter GH. Reduced Salivary Mucin Binding and Glycosylation in Older Adults Influences Taste in an In Vitro Cell Model. *Nutrients*. 2019;11(10):2280. Published 2019 Sep 24. doi:10.3390/nu11102280
- [20] Chen M, Shen W, Rowan NR, Kulaga H, Hillel A, Ramanathan M Jr, Lane AP. Elevated ACE2 expression in the olfactory neuroepithelium: implications for anosmia and upper respiratory SARS-CoV-2 entry and replication. *bioRxiv [Preprint]*. 2020 May 9:2020.05.08.084996. doi: 10.1101/2020.05.08.084996.
- [21] Bryce B, St Albin A, Murri S, et al. Massive transient damage of the olfactory epithelium associated with infection of sustentacular cells by SARS-CoV-2 in golden Syrian hamsters. *Brain Behav Immun*. 2020;89:579-586. doi:10.1016/j.bbi.2020.06.032
- [22] Piques A, Brondel L and Pénicaud L (2021) COVID 19-Induced Smell and Taste Impairments: Putative Impact on Physiology. *Front. Physiol*. 11:625110. doi: 10.3389/fphys.2020.625110
- [23] Kirschenbaum D, Imbach LL, Ulrich S, et al. Inflammatory olfactory neuropathy in two patients with COVID-19. *Lancet*. 2020;396(10245):166. doi:10.1016/S0140-6736(20)31525-7
- [24] Brann DH, Tsukahara T, Weinreb C, et al. Non-neuronal expression of SARS-CoV-2 entry genes in the olfactory system suggests mechanisms underlying COVID-19-associated anosmia. *Sci Adv*. 2020;6(31):eabc5801. doi:10.1126/sciadv.abc5801
- [25] Doty RL. Olfactory dysfunction and its measurement in the clinic and workplace. *Int Arch Occup Environ Health* 2006;79(4):268-82. <https://doi.org/10.1007/s00420-005-0055-6>.
- [26] Doty RL, Shaman P, Kimmelman CP, Dann MS. University of Pennsylvania Smell Identification Test: a rapid quantitative olfactory function test for the clinic. *Laryngoscope* 1984; 94(2 Pt 1):176-8. <https://doi.org/10.1288/00005537-198402000-00004>.
- [27] Cain WS, Gent JF, Goodspeed RB, Leonard G. Evaluation of olfactory dysfunction in the Connecticut Chemosensory Clinical Research Center. *Laryngoscope* 1988;98 (1):83-8. <https://doi.org/10.1288/00005537-198801000-00017>.
- [28] Hummel T, Sekinger B, Wolf SR, Pauli E, Kobal G. 'Sniffin' sticks': olfactory performance assessed by the combined testing of odor identification, odor discrimination and olfactory threshold. *Chem Senses* 1997;22(1):39-52. <https://doi.org/10.1093/chemse/22.1.39>
- [29] Kobayashi M, Reiter ER, DiNardo LJ, Costanzo RM. A new clinical olfactory function test: cross-cultural influence. *Arch Otolaryngol Head Neck Surg* 2007;133 (4):331-6. <https://doi.org/10.1001/archotol.133.4.331>.

- [30] Speth MM, Singer-Cornelius T, Oberle M, Gengler I, Brockmeier SJ, Sedaghat AR. Olfactory dysfunction and Sinonasal symptomatology in COVID-19: prevalence, severity, timing, and associated characteristics. *Otolaryngol Head Neck Surg* 2020; 163(1):114-20. <https://doi.org/10.1177/0194599820929185>.
- [31] Baig AM, Khaleeq A, Ali U, Syeda H. Evidence of the COVID-19 virus targeting the CNS: tissue distribution, host-virus interaction, and proposed neurotropic mechanisms. *ACS Chem Neurosci* 2020;11(7):995-8. <https://doi.org/10.1021/acscchemneuro.0c00122>.
- [32] Yan CH, Faraji F, Prajapati DP, Boone CE, DeConde AS. Association of chemosensory dysfunction and COVID-19 in patients presenting with influenza-like symptoms. *Int Forum Allergy Rhinol* 2020;10(7):806-13. <https://doi.org/10.1002/alr.22579>.
- [33] Lechien JR, Cabaraux P, Chiesa-Estomba CM, et al. Objective olfactory evaluation of self-reported loss of smell in a case series of 86 COVID-19 patients. *Head Neck* 2020;42(7):1583-90. <https://doi.org/10.1002/hed.26279>.
- [34] Seden N, Yiğit E, Yiğit Ö, Kaygısız İ. Objective evaluation of odor loss in COVID-19 and other suspected cases. *Am J Otolaryngol*. 2021;42(1):102761. doi:10.1016/j.amjoto.2020.102761
- [35] Gane SB, Kelly C, Hopkins C. Isolated sudden onset anosmia in COVID-19 infection. A novel syndrome? *Rhinology*. 2020. doi: 10.4193/Rhin20.114.
- [36] Amer MA, Elsherif HS, Abdel-Hamid AS, Elzayat S. Early recovery patterns of olfactory disorders in COVID-19 patients; a clinical cohort study. *Am J Otolaryngol*. 2020 Nov-Dec;41(6):102725. doi: 10.1016/j.amjoto.2020.102725.
- [37] Hopkins C, Surda P, Whitehead E, Kumar BN. Early recovery following new onset anosmia during the COVID-19 pandemic - an observational cohort study. *J Otolaryngol Head Neck Surg*. 2020 May 4;49(1):26. doi: 10.1186/s40463-020-00423-8.
- [38] Kosugi EM, Lavinsky J, Romano FR, et al. Incomplete and late recovery of sudden olfactory dysfunction in COVID-19. *Braz J Otorhinolaryngol*. 2020 July-Aug;86(4):490-496. doi: 10.1016/j.bjorl.2020.05.001.
- [39] Panda S, Mohamed A, Sikka K, et al. Otolaryngologic manifestation and long-term outcome in mild COVID-19: experience from a tertiary care centre in India. *Indian J Otolaryngol Head Neck Surg*. 2020;1-6. doi:10.1007/s12070-020-02217-w
- [40] Chiesa-Estomba CM, Lechien JR, Radulesco T, et al. Patterns of smell recovery in 751 patients affected by the COVID-19 outbreak. *Eur J Neurol*. 2020;27(11):2318-2321. doi:10.1111/ene.14440
- [41] Le Bon SD, Pisarski N, Verbeke J, et al. Psychophysical evaluation of chemosensory functions 5 weeks after olfactory loss due to COVID-19: a prospective cohort study on 72 patients. *Eur Arch Otorhinolaryngol*. 2020;1-8. doi:10.1007/s00405-020-06267-2
- [42] Iannuzzi L, Salzo AE, Angarano G, et al. Gaining back what is lost: recovering the sense of smell in mild to moderate patients after COVID-19. *Chem Senses* 2020 Dec 5;45(9):875-881. doi: 10.1093/chemse/bjaa066. PMID: 33033827; PMCID: PMC7665358.
- [43] Vaira LA, Hopkins C, Petrocelli M, et al. Smell and taste recovery in coronavirus disease 2019 patients: a 60-day objective and prospective study. *J Laryngol Otol*. 2020;134(8):703-709. doi:10.1017/S0022215120001826
- [44] Otte MS, Eckel HNC, Poluschkin L, Klusmann JP, Luers JC. Olfactory

dysfunction in patients after recovering from COVID-19. *Acta Otolaryngol.* 2020 Dec;140(12):1032-1035. doi: 10.1080/00016489.2020.1811999. Epub 2020 Aug 27. PMID: 32852240.

[45] Boscolo-Rizzo P, Menegaldo A, Fabbris C, Spinato G, Borsetto D, Vaira LA, Calvanese L, Pettorelli A, Sonego M, Frezza D, Bertolin A, Cestaro W, Rigoli R, D'Alessandro A, Tirelli G, Da Mosto MC, Menini A, Polesel J, Hopkins C. Six-Month Psychophysical Evaluation of Olfactory Dysfunction in Patients with COVID-19. *Chem Senses.* 2021 Jan 1;46:bjab006. doi: 10.1093/chemse/bjab006.

[46] Vroegop AV, Eeckels AS, Rompaey V Van, et al. COVID-19 and olfactory dysfunction – an ENT perspective to the current COVID-19 pandemic. *B-ENT.* 2020. doi:10.5152/b-ent.2020.20127.

[47] Whitcroft KL, Hummel T. Olfactory Dysfunction in COVID-19: Diagnosis and Management. *JAMA.* 2020. doi:10.1001/jama.2020.8391.

[48] Wolford RW, goyal A, Belgam Syed SY, Schaefer TJ. pharyngitis. [Updated 2020 Mar 30]. In: Statpearls [Internet]. Treasure Island (FL): Statpearls publishing; 2020. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK519550/>