We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Health Risks Associated with the Consumption of Legumes Contaminated with Pesticides and Heavy Metals

Motunrayo Ganiyat Akande

Abstract

Legumes have high nutritional value and they are important sources of protein, carbohydrates, fats and dietary fiber. The contamination of legumes with pesticides and heavy metals has been reported in scientific literature. Human beings are mainly exposed to the residues of pesticides and heavy metals through the dietary route. The purpose of this review chapter is to highlight the acute and chronic health risks that human beings may be exposed to as a result of the ingestion of legumes polluted with pesticides and heavy metals. Additionally, the mechanisms through which pesticides and heavy metals engender different undesirable health outcomes in human beings were stated. Scientific literature were perused and the information contained in them were collated to derive this chapter. Pesticides cause short-term health effects including hypersensitivity and mortality, while heavy metals induce acute effects like seizures and death. Some chronic untoward effects of pesticides are congenital disabilities and neurological damage. Heavy metals elicit disorders like anemia, hypertension and cancer. It is envisaged that the findings documented in this review will create awareness of the health risks posed by the contamination of legumes with the residues of pesticides and heavy metals so that food safety measures can be enforced globally.

Keywords: Legumes, Health risks, Pesticides, Heavy metals, Contamination

1. Introduction

Legumes are plants that belong to the family, *Fabaceae* or *Leguminosae*, and they include chickpeas, cowpea, lentils, soy, etc. [1]. They are highly nutritious and can be consumed as food by human beings and animals [2].

Pesticides are chemical substances that are manufactured for the control of pests in domestic, agricultural and industrial settings, among others [3]. They are classified as herbicides, fungicides, bactericides and insecticides mainly for agricultural applications [4]. Insecticides comprise carbamates, organochlorines, organophosphates, neonicotinoids, pyrethroids; herbicides include benzothiazolyl urea, carbamic and sulfanilic acids, isoxazolyl urea, phenylpyrazole and pyridinium; while fungicides contain carboxamides, carbamates, dithiocarbamates, etc. [5, 6].

Heavy metals (HMs) are a group of metals that possess high atomic weights and densities beyond 5 g/cm^3 [7]. They are usually derived from agricultural, mining and industrial activities, as well as effluents [8]. It has been observed that HMs for instance, lead (Pb), cadmium (Cd), zinc (Zn), nickel (Ni) and copper (Cu), build up in the soil and in plant uptake structures and evoke injurious effects on the environment and the health status of human beings [7, 9, 10]. Besides, HMs have the propensity to be toxic when the populace are exposed to them or when they are consumed in proportions beyond the acceptable daily limits [11].

Different food stuffs such as legumes, cereals, fruits, vegetables, etc. have been contaminated with pesticides and HMs in various parts of the world. It is noteworthy that the common environmental pollutants, pesticides and HMs, increasingly cause numerous health hazards to the populace because of their permeation and upsurge in the food chain, as well as their persistence in the bionetwork [12].

In 2020, [13] identified the organophosphate pesticides (malathion, parathion, ethion and carbophenothion) in cowpea (an African legume) from Gwagwalada market in Abuja, Nigeria, through the use of Gas Chromatography–Mass Spectrometry. The levels of the pesticides found in the cowpea samples exceeded the maximum residue limits set by the European Union and the Agency for Toxic Substances and Disease Registry. Additionally, [14] confirmed the contamination of cowpea by residues of organochlorine (endosulfan and lindane), and organophosphorus (malathion) pesticides in high levels in Northern Cameroun.

Furthermore, heavy metals including nickel, cadmium, manganese and cobalt were detected in cowpea by [15]. In the research, the concentrations of nickel, cobalt and manganese discovered in the legumes were below the acceptable limits by FAO/WHO, while the cadmium concentration was beyond the FAO/WHO limit permitted. However, in a study conducted in Saudi Arabian markets, it was discovered that kidney beans and haricots contained high levels of Mn, while peas had elevated levels of zinc beyond the standard permissible levels [16]. Hence, regular monitoring of food stuffs for HM content was advocated.

It has been observed that fertilizers and pesticides are responsible for the increased level of soil pollution by non-essential micronutrients such as arsenic, cadmium, mercury, nickel and lead [17]. These non-essential micronutrients in the soil are conveyed through various plants as they accumulate in the edible portions [18]. Subsequently, the general population and animals may be exposed to the residues of these environmental contaminants (pesticides and HMs) when they consume plants that have been polluted with them.

The aim of this review chapter is to underscore the acute and chronic health risks that human beings may be exposed to during their life time as a result of the consumption of legumes contaminated with pesticides and HMs. In addition, the mechanisms through which pesticides and HMs engender different undesirable health outcomes in biological systems were highlighted.

It is envisaged that the information in this review chapter will stimulate and enhance concerted efforts towards the regular monitoring of legumes and other foodstuffs for pollutants in order to attain food safety.

2. Mechanisms through which pesticides induce adverse health effects

When human beings consume pesticides in legumes and other food sources through the dietary route, the pesticides may cause acute and chronic health risks through diverse mechanisms. For instance, organophosphate pesticides (chlorpyrifos, diazinon, dichlorvos, fenitrothion, malathion, parathion, etc.) bring about deleterious health effects by inhibiting the function of acetylcholinesterase,

reducing the secretion of insulin, and perturbation of the metabolism of nutrients in living systems [19–21].

Moreover, organophosphate pesticides stimulate the release of reactive oxygen species and this phenomenon might cause oxidative stress [22, 23]. Oxidative stress refers to a disparity between the levels of prooxidants and antioxidants thereby culminating in damage to vital molecules including DNA, RNA, lipids and proteins in biological systems [24]. Oxidative stress has been identified as an important mechanism through which different kinds of pesticides cause biological injuries to human beings and animals.

The organochlorine pesticides are chlorinated hydrocarbons that persist in the environment and they include methoxychlor, dieldrin, chlordane, mirex and lindane, among others [25]. They cause the stimulation of the nervous system through the perturbation of action potentials, thereby leading to paralysis and death [26]. Dietary exposure brings about the bioaccumulation of organochlorine pesticides in the body, and this may terminate in derangements in human health [27].

Carbamates such as methiocarb, carbaryl, aldicarb, propoxur and carbofuran reversibly inhibit acetylcholinesterase in mammals and humans [19]. This group of pesticides are capable of causing apoptosis and necrotic changes in the cells of the immune system [28]. Consequently, the immune system becomes compromised and affected individuals become susceptible to various diseases when they are exposed to these pollutants in food and other sources.

Pyrethroids (e.g. allethrin, permethrin, cypermethrin and deltamethrin) are neurotoxins and they disrupt the muscular structures and modify voltage-dependent sodium channels in the body [29]. They exhibit low acute toxicity to mammals and avian species. Conversely, at low levels, pyrethroids elicit acute toxicity to diverse aquatic species and arthropods [30].

3. Acute health risks associated with dietary exposure to pesticides

Acute health risks may become visible instantaneously or within 24 hours sequel to exposure to a pesticide [31]. Pesticides can have access to human bodies through the dermal [32], ocular [33], oral and respiratory [34] routes. According to [35], the exposure of the populace to pesticide residues through the oral route is about five orders of enormity compared to the other routes. It has been affirmed that pesticides can produce acute health disorders such as hypersensitivity, asthma and mortality in people [8, 36–38].

4. Chronic health risks related to dietary exposure to pesticides

It has been reported that chronic detrimental health risks do not manifest in human beings even within a day subsequent to pesticide exposure [31]. Chronic health risks evoked by exposure to pesticides in food such as legumes include congenital disabilities and decreased birth weight [36, 37]. Additionally, many organophosphate insecticides bring about declines in sperm counts, viability, density and motility; inhibition of spermatogenesis, reductions in testes weights, and sperm DNA damage in males [38]. Other chronic adverse effects of pesticides entail hindering the activities of hormones, their time of release, or mimicking these hormones, thereby culminating in reduced fertility and deformities in the male and female reproductive tracts [39]. These harmful effects may result in declines in the population of human beings affected. Also, pesticides perturb the immune system function and elicit carcinogenicity [40, 41].

Some investigators have asserted that organochlorine pesticides and their active metabolites are linked to neurological aberrations, cancers, hypertension, cardiovascular and dermatological disorders in humans [42–44]. Besides, the exposure of the populace to organophosphate, organochlorine and carbamate pesticides may evoke chronic neurological disorders such as Alzheimer's and Parkinson's disease [45]. According to [46], pesticides affect neuronal function negatively by hyperphosphorylation and the disruption of microtubules thereby ultimately causing Alzheimer's disease.

5. Mechanisms by which heavy metals induce health risks in the human population

Heavy metals can accumulate in the bones or adipose tissues of human beings through dietary intake (for example, the consumption of legumes and other crops), thereby leading to the diminution of critical nutrients and undermined immune defenses [8].

Lead induces oxidative stress in tissues and it is plausible that it brings about deleterious effects in humans that consume legumes and other plants contaminated with it through this mechanism [47]. Also, lead, a ubiquitous lethal HM, may activate the development of tumors through the production of reactive oxygen species, as well as the promotion of damage to DNA and its repair mechanisms in living organisms [48].

Furthermore, it has been documented that arsenic sets off neoplasms through the alteration of the genome of human beings, perturbation of DNA and induction of oxidative injury [49, 50]. Another common HM, mercury, generates carcinogenesis through oxidative damage and interference with the structure, mending and preservation of DNA in living organisms [51].

These myriads of mechanisms may be responsible for the untoward acute and chronic health hazards reported in human beings who have consumed foodstuffs such as legumes, cereals, fruits, vegetables, etc. that have been tainted with HMs over a period of time.

6. Acute health risks associated with dietary exposure of individuals to heavy metals

Heavy metal contamination is a health menace to both adults and children [52]. When HMs build up in human tissues internally, for instance, when legumes polluted with HMs are eaten for some time, they may harm the central nervous system, and produce seizures, headache and coma [8].

There is scientific evidence that lead may engender neurotoxicity, nephrotoxicity and impaired haeme synthesis [53]. When children have contact with cadmium through a range of routes, for example the oral route (by feeding on legumes such as chickpeas, cowpea, etc.), they become exceedingly prone to lead intoxication and permanent neurological abnormalities may ensue [54]. Other acute impacts of lead exposure documented in children are inattention, hyperactivity, increased dullness, irritability, headache, convulsion, coma and death [55, 56].

Copper is important for the normal functioning of the brain, but it can be noxious if the cellular concentration surpasses the metabolic requirement [57]. Elevated levels of copper can cause dysfunctions in the working memory of individuals [58], while short-term exposure to copper through dietary and other routes have been linked with stomach pain, haematemesis, melena, jaundice, anorexia

and vomiting [59]. Moreover, rhabdomyolysis, cardiac and renal failure, hepatic necrosis, haemolysis, methemoglobinemia, encephalopathy, as well as mortality can occur in severe copper toxicity [60]. These harmful acute impacts of high copper levels may be attributed to its induction of oxidative stress, DNA damage and lessening of cell proliferation [61].

Moreover, the distortion of the concentrations of high-density lipoproteins and impairment of the immune system have been ascribed to excessive concentrations of zinc in the bodies of human beings [62]. People are usually exposed to zinc through the ingestion of contaminated food, for instance, legumes, pulses, among others [63].

7. Chronic health risks linked with the exposure of people to heavy metals through the oral route

Chronic health effects like diabetes, neurodegenerative diseases, renal damage, bone disorders and tumors of the breast, prostate and lungs have been elicited by Cd in people [64–66].

There are existing reports that indicate that the long-term effects of lead exposure through food and other routes are manifested as anemia, abdominal colic, miscarriages, male infertility, birth defects, renal diseases and behavioral dysfunctions in children [67, 68]. It has also been observed that lead may elicit decreased circulating maternal thyroid hormone thereby influencing growth patterns adversely [67, 68].

It is known that tremendous amounts of arsenic in the soil, food crops (including legumes, cereals, vegetables, etc) and groundwater can stimulate cancer and dermal aberrations, as well as disorders in the heart, stomach, intestines, liver, kidneys and brain, among others [62, 69–73].

Furthermore, excessive copper consumption can activate hepatic damage and other gastric-related problems in people [62, 74, 75]. Wilson's disease, a form of chronic Cu toxicity in human beings, is characterized by alterations in mental states, motor disorders, dysphagia, incoordination, haemolytic anemia, renal and hepatic dysfunctions [72]. It has been shown that human beings become susceptible to chronic copper poisoning when they ingest food items (for instance, legumes, pulses, cereals, etc.) contaminated with copper probably through polluted irrigation sources [76]. Additionally, persistent bronchitis, emphysema, pulmonary disorders and fibrosis were reported in people following long-term exposure to nickel through dietary sources [77].

8. Conclusion

This review chapter highlighted the mechanisms through which the widespread pollutants, pesticides and HMs, evoke acute and chronic health disorders in human beings. Pesticides are usually applied in agricultural settings for the control of vectors and the enhancement of crop yield, while most HMs are utilized for industrial purposes.

Human beings may be susceptible to various health risks engendered by pesticides and HMs through the consumption of food crops, for example, legumes, contaminated with them. This may eventually pose serious threats to the wellbeing and survival of the populace except if regular monitoring of food items for the residues of these ubiquitous contaminants are conducted globally by the appropriate agencies.

It is envisioned that the information contained in this review chapter will provide a springboard for scientists, researchers and agriculturists, among others, to create innovative techniques for the minimization of human exposures to the residues of pesticides and HMs in order to forestall their pernicious effects in the general population.

Acknowledgements

The author is grateful to the members of staff of the Faculty of Veterinary Medicine, University of Abuja, Federal Capital Territory, Nigeria, for their support.

Conflict of interest

The author declares that there is no conflict of interest.

Author details

Motunrayo Ganiyat Akande
Department of Veterinary Pharmacology and Toxicology, Faculty of Veterinary Medicine, University of Abuja, Federal Capital Territory, Nigeria

*Address all correspondence to: akande.ganiyat@uniabuja.edu.ng

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Stagnari F, Pisante M. Managing faba bean residues to enhance the fruit quality of the melon (*Cucumis melo* L.). *Crop Sci Hort*. 2010; 126: 317-323. <https://doi.org/10.1016/j.scienta.2010.07.004>.
- [2] Adepoju FA, Adepoju AF, Fadina OO. Heavy metal levels in beans (*Vigna unguiculata*) in selected markets in Ibadan, Nigeria. *J Exp Agric Int*. 2018; 25: 1-8. DOI: 10.9734/jeai/2018/35900.
- [3] Damalas CA, Eleftherohorinos IG. Pesticide exposure, safety issues, and risk assessment indicators. *Int J Environ Res Public Health*. 2011; 8: 1402-1419. DOI: 10.3390/ijerph8051402.
- [4] Abdel Khalek ST, Mostafa ZK, Hassan HA, Abd El-Bar MM, Abu El-Hassan GMM. A new list to the entomofauna associated with faba bean, *Vicia faba* L. (*Fabales: Fabaceae*) grown in El-Kharga Oasis, New Valley Governorate, Egypt. *Egypt. Acad J Biol Sci*. 2018; 11: 95-100.
- [5] Kaur R, Mavi GK, Raghav S, Khan I. Pesticides classification and its impact on environment. *Int J Curr Microbiol Appl Sci*. 2019; 8: 1889-1897. <https://doi.org/10.20546/ijcmas.2019.803.224>.
- [6] Lippmann M, Leikauf G, editors. *Environmental Toxicants Human Exposures and Their Health Effects*. 4th ed. New Jersey; 2020. p. 1-40.
- [7] Zhang X, Yan L, Liu J, Zhang Z, Tan C. Removal of different kinds of heavy metals by novel PPG-nZVI beads and their application in simulated stormwater infiltration facility. *Appl Sci*. 2019; 9: 4213. DOI:10.3390/app9204213.
- [8] Alengebawy A, Abdelkhalek ST, Qureshi SR, Wang MQ. Heavy metals and pesticides toxicity in agricultural soil and plants: ecological risks and human health implications. *Toxics*. 2021; 9: 42. <https://doi.org/10.3390/toxics9030042>.
- [9] Su C, Jiang L, Zhang W. A review on heavy metal contamination in the soil worldwide: situation, impact and remediation techniques. *Environ Skept Critics*. 2014; 3: 24-38.
- [10] Tóth G, Hermann T, Da Silva MR, Montanarella L. Heavy metals in agricultural soils of the European Union with implications for food safety. *Environ Int*. 2016; 88: 299-309. <https://doi.org/10.1016/j.envint.2015.12.017>.
- [11] Onakpa MM, Njan AA, Kalu OC. A review of heavy metal contamination of food crops in Nigeria. *Annals of Global Health*. 2018; 84: 488-494. DOI: <https://doi.org/10.29024/aogh.2314>.
- [12] Verger PJP, Boobis AR. Reevaluate pesticides for food security and safety. *Science*. 2013; 341: 717-718. <https://doi.org/10.1016/j.envint.2015.12.017>.
- [13] Akande MG, Sanni FS, Enefe NG. Human health risk evaluation of organophosphate insecticide residues in post-harvest cowpea in Gwagwalada, Abuja, Nigeria. *J Health Pollution*. 2020; 10: 201203. DOI: 10.5696/2156-9614-10.28.201203.
- [14] Jean S, Benoit NM, Bosco TJ, Srivastava AK, Srivastava LP. Contamination of cowpea and by-products by organophosphorous pesticide residues in Ngaoundere markets: dietary risk estimation and degradation study. *Afr J Food Sci*. 2013; 7: 92-102. <https://doi.org/10.5897/AJFS2013.1013>.
- [15] Akande MG, Sanni FS, Enefe NG. Assessment of the concentrations and health risk of some heavy metals in cowpea (*Vignus unguiculata*) in Gwagwalada, Nigeria. *Drug and*

Chemical Toxicology. 2019; doi:
10.1080/01480545.2019.1621334.

[16] Ali MHH, Al-Qahtani KM. Assessment of some heavy metals in vegetables, cereals and fruits in Saudi Arabian markets. *Egyptian Journal of Aquatic Research*. 2012; 38: 31-37. <https://doi.org/10.1016/j.ejar.2012.08.002>.

[17] Gunatilake SK, Samaratunga SS, Rubasinghe RT. Chronic kidney disease (CKD) in Sri Lanka - current research evidence justification: a review. *Sabaragamuwa University Journal*. 2014; 13: 31-58. DOI:10.4038/susljv13i2.7680.

[18] Edirisinghe EMRKB, Jinadasa BKKK. Arsenic and cadmium concentrations in legumes and cereals grown in the North Central Province, Sri Lanka and assessment of their health risk. *International Journal of Food Contamination*. 2019; 6:3. <https://doi.org/10.1186/s40550-019-0073-x>.

[19] Shah MAA, Srivastava AK Garg. Toxicology of Agrochemicals. In: Garg SK, editor. 1st ed. CBS Publishers & Distributors Pvt. Ltd: New Delhi; 2002. p. 146-179.

[20] Jaga K, Dharmani C. Sources of exposure to and public health implications of organophosphate pesticides. *Rev Panam Salud Publica*. 2003; 14:171-185. DOI: 10.1590/s1020-49892003000800004.

[21] Karami-Mohajeri S, Abdollahi M. Toxic influence of organophosphate, carbamate, and organochlorine pesticides on cellular metabolism of lipids, proteins, and carbohydrates: a systematic review. *Hum Exp Toxicol*. 2011; 30(9):1119-1140. DOI: 10.1177/0960327110388959.

[22] Abdollahi M, Ranjbar A, Shadnia S, Nikfar S, Rezaie A. Pesticides and oxidative stress: a review. *Med Sci*

Monit. 2004; 10: 141-147. PMID: 15173684.

[23] Sharma D, Kaur GS, Khera KS. Triazophos-induced oxidative stress and histomorphological changes in ovary of female Wistar rats. *Pestic Biochem Physiol*. 2014; 117: 9-18. DOI: 10.1016/j.pestbp.2014.09.004.

[24] Conti V, Izzo V, Corbi G, Russomanno G, Manzo V, De Lise F, Di Donato A, Filippelli A. Antioxidant supplementation in the treatment of aging-associated diseases. *Front Pharmacol*. 2016; 7:24. DOI: 10.3389/fphar.2016.00024.

[25] Jones KC, de Voogt P. Persistent organic pollutants (POPs): state of the science. *Environmental Pollution*. 1999; 100: 209-221. [https://doi.org/10.1016/S0269-7491\(99\)00098-6](https://doi.org/10.1016/S0269-7491(99)00098-6).

[26] Aliu YO. *Veterinary Pharmacology*. 1st ed. Wusasa-Zaria: Tamaza Publishing Company Limited; 2007. 466 p.

[27] Snedeker S. Pesticides and breast cancer risk: a review of DDT, DDE and dieldrin. *Environ Health Perspect*. 2001; 109 (suppl 1): 35-47. DOI: 10.1289/ehp.01109s135.

[28] Li Q, Kobayashi M, Kawada T. Ziram induces apoptosis and necrosis in human immune cells. *Arch Toxicol*. 2011; 85: 355-361. <https://doi.org/10.1007/s00204-010-0586-9>.

[29] Ray DE, Fry JR. A reassessment of the neurotoxicity of pyrethroid insecticides. *Pharmacol Ther*. 2006; 111: 174-193. DOI: 10.1016/j.pharmthera.2005.10.003.

[30] Pérez-Fernández V, Garcia MA, Marina ML. Characteristics and enantiomeric analysis of chiral pyrethroids. *J Chromatogr A*. 2010; 1217: 968-989. DOI: 10.1016/j.chroma.2009.10.069.

- [31] Singh NS, Sharma R, Parween T, Patanjali PK. Pesticide contamination and human health risk factor. In: Oves M, et al. editors. *Modern Age Environmental Problems and their Remediation*. International Publishing AG: Springer; 2018. p. 50-68. DOI: 10.1007/978-3-319-64501-8_3.
- [32] Anderson SE, Meade BJ. Potential health effects associated with dermal exposure to occupational chemicals. *Environ Health Insights*. 2014; 8s1, EHI. S15258. DOI: 10.4137/EHI.S15258.
- [33] Fareed M, Kesavachandran CN, Pathak MK, Bihari V, Kuddus M, Srivastava AK. Visual disturbances with cholinesterase depletion due to exposure of agricultural pesticides among farm workers. *Toxicol Environ Chem*. 2012; 94: 1601-1609. <https://doi.org/10.1080/02772248.2012.718780>.
- [34] Amaral AFS. Pesticides and asthma: challenges for epidemiology. *Front Public Health*. 2014; 2: 6. DOI: 10.3389/fpubh.2014.00006.
- [35] Olutona GO, Aderemi MA. Organochlorine pesticide residue and heavy metals in leguminous food crops from selected markets in Ibadan, Nigeria. *Legume Science*. 2019;1-9. <https://doi.org/10.1002/leg3.3>.
- [36] Sharon M, Bhawana M, Anita S, Gothecha VK. A short review on how pesticides affect human health. *Int J Ayurvedic Herb Med*. 2012; 2: 935-946.
- [37] Wickerham EL, Lozoff B, Shao J, Kaciroti N, Xia Y, Meeker JD. Reduced birth weight in relation to pesticide mixtures detected in cord blood of full-term infants. *Environ Int*. 2012; 47: 80-85. DOI: 10.1016/j.envint.2012.06.007.
- [38] Mehrpour O, Karrari P, Zamani N, Tsatsakis AM, Abdollahi M. Occupational exposure to pesticides and consequences on male semen and fertility: a review. *Toxicol Lett*. 2014; 230: 146-156. DOI: 10.1016/j.toxlet.2014.01.029.
- [39] Kabir ER, Rahman MS, Rahman I. A review on endocrine disruptors and their possible impacts on human health. *Environ Toxicol Pharmacol*. 2015; 40: 241-258. DOI: 10.1016/j.etap.2015.06.009.
- [40] Garry VF. Pesticides and children. *Toxicol Appl Pharmacol*. 2004; 198: 152-163. DOI: 10.1016/j.taap.2003.11.027.
- [41] Özkara A, Akyıl D, Konuk M. Pesticides, environmental pollution and health. In: Larramendy ML, Soloneski S, editors. *Environmental Health Risk-Hazardous Factors to Living Species*: IntechOpen; 2016. DOI: 10.5772/63094.
- [42] Walbel H, Henckes C, editors. *Pesticides use and policies in Ghana: an economic analysis of current practice and factors influencing pesticide use*. Pesticide Policy Project Publication Series No. 10. Hannover: Germany; 2001.
- [43] Mansour SA. Pesticide exposure: Egyptian scene. *Toxicology*. 2004; 198: 91-115. DOI: 10.1016/j.tox.2004.01.036.
- [44] Parbhu B, Rodgers G, Sullivan JE. Death in a toddler following endosulfan ingestion. *Clin Toxicol*. 2009; 47: 899-901. DOI: 10.3109/15563650903328879.
- [45] Keifer MC, Firestone J. Neurotoxicity of pesticides. *J Agromed*. 2007; 1:17-25. DOI: 10.1300/J096v12n01_03.
- [46] Zaganas I, Kapetanaki S, Mastorodemos V, Kanavouras K, Colosio C, Wilks MF, Tsatsakis AM. Linking pesticide exposure and dementia: what is the evidence? *Toxicology*. 2013; 10: 3-11. DOI: 10.1016/j.tox.2013.02.002.

- [47] Jaishankar M, Tseten T, Anbalagan N, Mathew BB, Beeregowda KN. Toxicity, mechanism and health effects of some heavy metals. *Interdiscip Toxicol.* 2014; 7: 60-72. DOI: 10.2478/intox-2014-0009.
- [48] Silbergeld EK, Waalkes M, Rice JM. Lead as a carcinogen: Experimental evidence and mechanisms of action. *Am J Ind Med.* 2000; 38: 316-323. DOI: 10.1002/1097-0274(200009)38:3<316::aid-ajim11>3.0.co;2-p.
- [49] Martinez VD, Vucic EA, Becker-Santos DD, Gil L, Lam WL. Arsenic exposure and the induction of human cancers. *J Toxicol.* 2011; 2011: 1-13. DOI: 10.1155/2011/431287.
- [50] Bjorklund G, Aaseth J, Chirumbolo S, Urbina MA, Uddin R. Effects of arsenic toxicity beyond epigenetic modifications. *Environ Geochem Health.* 2017; 39: 1-11. DOI: 10.1007/s10653-017-9967-9.
- [51] Crespo-Lopez ME, Macedo GL, Pereira SI, Arrifano GP, Picanco-Diniz DL, do Nascimento JL, Herculano AM. Mercury and human genotoxicity: Critical considerations and possible molecular mechanisms. *Pharmacological Research.* 2009; 60(4):212-220. DOI: 10.1016/j.phrs.2009.02.011.
- [52] Lu X, Zhang X, Li LY, Chen H. Assessment of metals pollution and health risk in dust from nursery schools in Xi'an, China. *Environ Res.* 2014; 128: 27-34. <https://doi.org/10.1016/j.envres.2013.11.007>.
- [53] Malik RN, Zeb N. Assessment of environmental contamination using feathers of *Bubulcus ibis* L., as a biomonitor of heavy metal pollution, Pakistan. *Ecotoxicology.* 2009; 18: 522-536. DOI: 10.1007/s10646-009-0310-9.
- [54] Maret W. The bioinorganic chemistry of lead in the context of its toxicity. In: Sigel A, Sigel H, Sigel RKO, editors. *Lead: its effects on environment and health.* Volume 17. Berlin: De Gruyter; 2017. p. 1-20.
- [55] Pfadenhauer LM, Burns J, Rohwer A, Rehfuess EA. A protocol for a systematic review of the effectiveness of interventions to reduce exposure to lead through consumer products and drinking water. *Syst Rev.* 2014; 3: 36. DOI: 10.1186/2046-4053-3-36
- [56] Evens A, Hryhorczuk D, Lanphear BP, Rankin KM, Lewis DA, Forst L, Rosenberg D. The impact of low-level lead toxicity on school performance among children in the Chicago Public Schools: A population-based retrospective cohort study. *Environ Health.* 2015; 14: 21. <https://doi.org/10.1186/s12940-015-0008-9>.
- [57] Johnson WT. Nutritional neuroscience. In: Lieberman HR, Kanarek RB, Prasad C, editors. *Nutritional Neuroscience.* Boca Raton: CRC Press (Taylor and Francis Group); 2005. p. 289-305.
- [58] Zhou G, Ji X, Cui N, Cao S, Liu C, Liu J. Association between serum copper status and working memory in schoolchildren. *Nutrients.* 2015; 7: 7185-7196. DOI: 10.3390/nu7095331.
- [59] Gamakaranage CSSK, Rodrigo C, Weerasinghe S, Gnanathasan A, Puvanaraj V, Fernando H. Complications and management of acute copper sulphate poisoning: a case discussion. *J Occup Med Toxicol.* 2011; 6: 34. DOI: 10.1186/1745-6673-6-34.
- [60] Harris ED. Cellular copper transport and metabolism. *Annu Rev Nutr.* 2000; 20: 291-310. DOI: 10.1146/annurev.nutr.20.1.291.
- [61] Oe S, Miyagawa K, Honma Y, Harada M. Copper induces hepatocyte

- injury due to the endoplasmic reticulum stress in cultured cells and patients with Wilson disease. *Exp Cell Res*. 2016; 347: 192-200. DOI: 10.1016/j.yexcr.2016.08.003.
- [62] Zhou H, Yang WT, Zhou X, Liu L, Gu JF, Wang WL, Zou JL, Tian T, Peng PQ, Liao BH. Accumulation of heavy metals in vegetable species planted in contaminated soils and the health risk assessment. *Int J Environ Res Public Health*. 2016; 13: 289. DOI: 10.3390/ijerph13030289.
- [63] Hough RL, Breward N, Young SD, Crout NM, Tye AM, Moir AM, Thornton I. Assessing potential risk of heavy metal exposure from consumption of home produced vegetables by urban populations. *Environ. Health Perspect*. 2004; 112: 215-221. DOI: 10.1289/ehp.5589.
- [64] Avenant-Oldewage A, Marx HM. Bioaccumulation of chromium, copper and iron in the organs and tissues of *Clarias gariepinus* in the Olifants River, Kruger National Park. *Water SA*. 2000; 26: 569-582.
- [65] Jiang LF, Yao TM, Zhu ZL, Wang C, Ji LN. Impacts of Cd (II) on the conformation and self-aggregation of Alzheimer's tau fragment corresponding to the third repeat of microtubule-binding domain. *Biochim Biophys Acta-Proteins Proteomics*. 2007; 1774: 1414-1421. DOI: 10.1016/j.bbapap.2007.08.014.
- [66] Chunhabundit, R. Cadmium exposure and potential health risk from foods in contaminated area, Thailand. *Toxicol. Res*. 2016; 32: 65-72. DOI: 10.5487/TR.2016.32.1.065.
- [67] Gundacker C, Hengstschläger M. The role of the placenta in fetal exposure to heavy metals. *Wien Med Wochenschr*. 2012; 162: 201-206. DOI: 10.1007/s10354-012-0074-3.
- [68] Wani AL, Ara A, Usmani JA. Lead toxicity: a review. *Interdiscip Toxicol*. 2015; 8: 55-64. DOI: 10.1515/intox-2015-0009.
- [69] Chiou HY, Hsueh YM, Liaw KF, Horng SF, Chiang MH, Pu YS, Lin JS, Huang CH, Chen CJ. Incidence of internal cancers and ingested inorganic arsenic: a seven-year follow-up study in Taiwan. *Cancer Res*. 1995; 55: 1296-1300. PMID: 7882325.
- [70] Kapaj S, Peterson H, Liber K, Bhattacharya P. Human health effects from chronic arsenic poisoning - a review. *J Environ Sci Health A Tox Hazard Subst Environ Eng*. 2006; 41: 2399-2428. DOI: 10.1080/10934520600873571.
- [71] Lin HJ, Sunge T, Cheng CY, Guo HR. Arsenic levels in drinking water and mortality of liver cancer in Taiwan. *J Hazard Mater*. 2013; 262: 1132-1138. DOI: 10.1016/j.jhazmat.2012.12.049.
- [72] Islam S, Rahman MM, Rahman MA, Naidu R. Inorganic arsenic in rice and rice-based diets: health risk assessment. *Food Control*. 2017; 82: 196-202. <https://doi.org/10.1016/j.foodcont.2017.06.030>.
- [73] El-Kady AA, Abdel-Wahhab MA. Occurrence of trace metals in foodstuffs and their health impact. *Trends Food Sci Technol*. 2018; 75: 36-45. <https://doi.org/10.1016/j.tifs.2018.03.001>.
- [74] Gaetke LM, Chow CK. Copper toxicity, oxidative stress, and antioxidant nutrients. *Toxicology*. 2003; 189: 147-163. DOI: 10.1016/S0300-483X(03)00159-8.
- [75] Rahman MA, Rahman MM, Reichman SM, Lim RP, Naidu R. Heavy metals in Australian grown and imported rice and vegetables on sale in Australia: health hazard. *Ecotoxicol Environ Saf*. 2014; 100: 53-60. <https://doi.org/10.1016/j.ecoenv.2013.11.024>.

[76] El-Kady AA, Abdel-Wahhab MA. Occurrence of trace metals in foodstuffs and their health impact. Trends Food Sci. Technol. 2018; 75: 36-45. <https://doi.org/10.1016/j.tifs.2018.03.001>.

[77] United States Department of Health and Human Services, Agency for Toxic Substances and Disease Registry. 2005; 1-397.