We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Genetic Resources of The Universal Flavor, Vanilla

Minoo Divakaran and N.T. Fathima Rafieah

Abstract

Commercially cultivated vanilla (*V. planifolia*) is native to Mexico and its cultivation and breeding programmes face major bottlenecks. This study reports presence of important agronomic characters in two important and endangered species of Vanilla, *V. aphylla* and *V. pilifera*, indigenous to India. *V. aphylla* was tolerant to Fusarium wilt and had longer flower life than the cultivated vanilla. *V. pilifera* flowers were fragrant, showed signs of insect pollination and had large fruit size. The species were amenable to interspecific hybridization and successful reciprocal crosses were done. Sequence similarity studies indicated the clustering of leafy and leafless species separately.

Keywords: interspecific hybridization, *V. aphylla*, *V. pilifera*, sequence similarity

1. Introduction

The genus Vanilla includes about 110 species and the species have been treated in various monographic works [1, 2] including the life history of *V. planifolia* [3]. *Vanilla planifolia* (Salisb.) Ames (syn. *V. fragrans* Andrews.), is a tropical climbing orchid known for yielding the delicate popular flavor, vanilla [4] and is the second most expensive spice traded in the world market [5] (Spices Board 2000). The major vanilla producing countries are Madagascar, Comoro, Indonesia, Mexico and the Reunion, of which, Madagascar holds the prominent position.

Vanilla was introduced to Europe from Mexico, in about 1500 and its reputation of being an aphrodisiac followed it to countries where it was introduced. The importance of vanilla since early times in Mexico, is evident by the mention of offering vanilla as a medicinal beverage as part of a tribute during reign of Itzcoatl (Aztec Emperor) in 1427 and citing vanilla as a remedy for fatigue in Badianus manuscript in 1552 [6]. *Vanilla planifolia*, which yields the vanilla of commerce, is native to Mexico and parts of Central America and the history of origin of cultivated vanilla suggests that the entire stock outside Mexico may be from a single genetic source. For the last 400 years, humans have been playing important role in the dispersal and spread of vanilla in the New World.

2. Species of Vanilla

Studies of divergence among species of agronomic importance have been receiving greater attention. Genomics-based tools are efficient to characterize and

identify genetic diversity in Vanilla and act as a significant tool for genomics-assisted plant breeding [7]. RAPD polymorphism was used to estimate the level of genetic diversity and interrelationships among few related species *Vanilla planifolia*, including both leafy and leafless types such as *V. tahitensis*, *V. andamanica*, *V. pilifera* and *V. aphylla*. Studies revealed that there is very limited variation within collections of *V. planifolia*, indicative of its narrow genetic base [8]. The British introduced *Vanilla planifolia* into India about 200 years ago whereas five other species are native viz., *V. pilifera* Holt., *V. andamanica* Rolfe., *V. aphylla* Blume., *V. walkeriae* Wight. and *V. wightiana* Lindl.

V. pilifera originally described from Malaya, recorded in Thailand is found in the Mikir hills of Northeast India. *V. aphylla*, an endangered species, previously known from Thailand is found in South India [9]. *V. andamanica* is endemic to Andaman group of Islands and is believed to be same as *V. albida* [10]. *V. tahitensis*, which is commonly exploited throughout the tropics, is indigenous to the Tahiti Islands. The presence and absence of leaves, and floral characters (colors of flower, lip, hairs on lip and ovary-pedicel etc.), morphologically distinguish these species (Table 1).

A preliminary analysis of the various characters of Vanilla species including the above species, showed presence and absence of leaves formed an important part in the classification of the genus which in general had the basic chromosome number $x = 16$. Most of the Indian species were leafless, except *V. pilifera* which was intermediate in character, i.e., leafless in early stages and long narrow leaves at maturity and the chromosome number in *V. aphylla* is $2n = 64$, whereas the cultivated vanilla and *V. tahitensis* had a somatic chromosome number of $2n = 32$ [10, 11]. Differences in floral characters existed in flower color and lip characters (Table 2). In *V. pilifera* vines, leaves developed as the vine grew with flowers that were narrower (2.8 x 0.8 cm) with distinct pure white ovary-pedicel (Figures 1 and 2), pale green tepals, purplish violet and longer (6 mm approx.) hairs on white lip. *V. aphylla* is leafless (with scales-1.8 cm) and yellowish-cream flowers (petal size 3 x 1.2 cm approx.) having tuft of hairs that are cream near tip, deep reddish inside (2–3 mm) and light green ovary-pedicel (Figures 3 and 4).

Species	Leaf type		Internode	Median ridge
	Shape	Size		
<i>V. aphylla</i>	Scale leaves to leafless	2.1 cm in fresh shoots	6.5 cm	Absent in fresh shoots
<i>V. pilifera</i>	Narrow (intermediate to <i>V. planifolia</i> and <i>V. aphylla</i>)	L (8.5–16.5 cm) B (1.6–3.1 cm)	7 cm	Present all along the stem

Table 1.
Vegetative characters of *Vanilla aphylla* and *V. pilifera*.

Species	Petal color	Ovary-pedicel	Tuft of hair on the lip	Nature	Fruit size after pollination – 4 weeks
<i>V. aphylla</i>	Yellowish cream, L-3 cm, B-1.2 cm	Light green	Cream near tip, reddish brown inside (2–3 mm)	Longer life	14 cm (L) 3 cm (B)
<i>V. pilifera</i>	Pale green, narrower, L-2.8 cm, B-0.8 cm	White	Violet and longer (6 mm)	More brittle	11.5 cm (L) 3.3 cm (B)

L, Length; B, Breadth.

Table 2.
Variations in floral characters.

Figure 1.
Members of V. aphylla inflorescence arranged sequentially (Inset: Close-up of an opened flower).

Figure 2.
Members of V. pilifera inflorescence arranged sequentially (Inset: Close-up of an opened flower).

Figure 3.
L.S. of flowers of *V. aphylla* (L) and *V. pilifera* (R).

Figure 4.
Comparison of dissected out flowers of *V. aphylla* (L) and *V. pilifera* (R).

3. Biotechnological applications

Micropropagation and *in vitro* conservation techniques for the different species of Vanilla [12] and interspecific hybridization as a tool for gene flow of desirable characters from wild species into cultivated species, through pollen, have been reported [13]. Genetic interrelationships studies, using RAPD profiles [8], among different species revealed that the leafless forms of vanilla, *V. aphylla* and *V. pilifera* formed a separate sub-cluster. All the other leafy vanilla types formed a separate sub-cluster. *V. pilifera*, which showed an intermediate leaf character, showed only 50–56.1% similarity to *V. planifolia* but closely resembled *V. aphylla* (76.8%). Thus, the present study reveals the presence of important agronomic characters for introgression into cultivated vanilla and which can be utilized to overcome major bottlenecks in vanilla breeding. The presence of fragrance which attracts insects, coupled with signs of fruit set without hand pollination, holds *V. pilifera* as a potential candidate for breeding programmes, to overcome the problem of lack of natural seed set in vanilla. *V. aphylla* which was tolerant to *Fusarium oxysporum* [8] and its crossability to cultivated vanilla can be utilized as a bridging species and to help wipe out diseases arising out of monoculture. Interspecific hybridization has been reported and hence transfer of these desirable traits into cultivated vanilla,

V. planifolia, may not be hindered. The advent of biotechnological tools, offers techniques for transfer of these characters into *V. planifolia*, thus making the dream of transforming vanilla into a fragrant, natural seed setting, disease tolerant commercially important orchid can be turned into a reality.

The identification of a hydratase/lyase type enzyme as being a vanillin synthase offers new opportunities for the Vanilla pod-based industries. The accumulation of vanillin glucoside in the capsules of cultivated vines in response to environmental challenges may now be assessed at the molecular level. Likewise, the basis for development of genetic markers for the selection of vanilla orchid varieties with improved aromatic properties has now been laid down. Vanillin produced biologically is termed 'natural' vanillin and has a high economic value compared with chemically synthesized vanillin. Likewise, in the transition towards a bio-based economy, it is important to develop sustainable production systems to replace those currently based on fossil fuels. The demonstration that a single enzyme in the vanilla pod catalyzes the conversion of ferulic acid and ferulic acid glucoside into vanillin and vanillin glucoside provides several options for biotechnological applications [14].

4. Materials and methods

4.1 Genomic DNA isolation

Genomic DNA was isolated from approximately 100 mg fresh leaves by grinding in a pestle and mortar using liquid Nitrogen and following the procedure using DNeasy® Plant Mini Kit (Qiagen, USA). The ground sample powder (100 mg) was transferred to microfuge tubes. Followed by addition of 400 µl AP1 buffer and 4 µl RNase A and mixed by vortex. The tubes were incubated at 65°C for 10 min in a water bath with intermittent mixing 2–3 times by inverting the tubes. Added 130 µl buffer P3 to the tube, mixed and incubated for 5 min on ice. The lysate was centrifuged for 5 min at 14,000 rpm. The samples were then loaded onto the QIAshredder spin columns and centrifuged at 14,000 rpm for 2 min. The flow-through was transferred to a new tube without disturbing the pellet. Added 1.5 volume of buffer AW1 and mixed by pipetting. The contents were then loaded in 650 µl fractions onto the DNeasy mini spin column and centrifuged at 8000 rpm for 1 minute. The flow-through was discarded. The spin column was placed into a new 2 ml collection tube and added 500 µl buffer AW2, followed by centrifugation for 1 min at 8000 rpm. This last step with buffer AW2 step was repeated, with centrifugation at 14,000 rpm for 2 min. The spin columns were placed in fresh microfuge tubes and 100 µl AE buffer was added onto the membranes and incubated at room temperature for 5 min. The tubes were then centrifuged at 8000 rpm for 1 min. This step was repeated with another 100 µl of AE buffer. The eluted samples were stored at –20°C.

4.2 Measurement of purity and DNA concentration

Quality and quantity of genomic DNA was monitored by using UV/Vis. Spectrophotometry and quality was confirmed by using 0.8% Agarose Gel Electrophoresis. Each of the sample DNA was diluted to 5 ng/µL in double distilled water for use as a PCR template.

4.3 PCR amplification of DNA barcoding region and sequencing

PCR reactions were carried out using universal primers for the DNA barcode regions matK, nrDNA-ITS, rbcL and trnH-psbA. All the specific locus primers were

purchased with universal M13 primer sequence at their 5' ends, thus enabling the direct sequencing of the PCR products using the universal M13 primers. The PCR amplification was performed in a 20 µl reaction mixture, consisting of 1X PCR buffer (2 mM Mgcl2), 200 µM each of dATP, dCTP, dGTP, dTTP; 0.5 µM of each forward and reverse primers and 1 U of Taq polymerase (TakaRa-Taq), and (5–20 ng) DNA template. DNA amplification was performed in a thermal cycler (Eppendoff, Germany). When the reaction has finished, the tubes were stored at 4°C. PCR products were separated by agarose gel electrophoresis (1.8%). The list of primers, their nucleotide sequences, annealing temperature and the specific PCR cycling conditions are shown in **Table 3**. A large volume PCR reaction (100 µl) per sample loci was done and PCR purification was done using Nucleospin Gel and PCR Clean-up kit (Macherey-Nagel, Germany). The purified PCR products were sequenced using M13 universal primers (M13 forward and M13 reverse primers) on ABI 3730xl

S. No	DNA region	Primers Name	Sequence (5'-3')	Reference	PCR reaction conditions
1	matK	390F	CGATCTATTCAATCAATATTTTC	[15]	95°C for 4 min. 95°C for 45 sec. 48°C for 30 sec 72°C for 50 sec. 72°C for 8 min. } 35X
		1326R	TCTAGCACACGAAAGTCGAAGT		
3	ITS	ITS4	TCCTCCGCTTATTGATATGC	[16]	94°C for 4 min. 94°C for 40 sec. 55 for 40 sec. 72°C for 1 min. 72°C for 8 min. } 36x
		ITS5	GGAAGTAAAGTCGTAACAAGG	[17]	
4	ITS	ITS-P5	CCTTATCAYTTAGAGGAAGGAG	[18]	94°C for 4 min. 94°C for 30 sec. 55°C for 40 sec. 72°C for 1 min. 72°C for 10 min. } 34x
		ITS-u4	RGTTTCTTTTCCTCCGCTTA		
5	rbcL	rbcL_1F	ATGTCACCACAAACAGAAAC	[19]	95°C for 4 min. 95°C for 45 sec. 55°C for 30 sec. 72°C for 50 sec. 72°C for 8 min. } 35x
		rbcL_724R	TCGCATGTACCTGCAGTAGC	[20]	
6	rbcL	rbcL_1F	ATGTCACCACAAACAGAAAC	[19]	95°C for 1 min. 95°C for 30 sec. 51°C for 30 Section 72°C for 1 min. 72°C for 5 min. } 35x
		rbcLa_r	CTTCTGCTACAAATAAGAATCGATCTC	[21]	
7	rbcL	rbcL_1F	ATGTCACCACAAACAGAAAC	[19]	95°C for 1 min. 95°C for 30 sec. 51°C for 30 sec. 72°C for 1 min. 72°C for 5 min. } 35x
		rbcLa_SI_Rev	GTAAATCAAGTCCACCRCG	[22]	
8	rbcL	rbcL_1F	ATGTCACCACAAACAGAAAC	[19]	95°C for 1 min. 95°C for 30 sec. 51°C for 30 sec. 72°C for 1 min. 72°C for 5 min. } 35x
		rbcLaj634R	GAAACGGTCTCTCCAACGCAT	[23]	
9	trnH-psbA	psbA3_f	GTTATGCATGAACGTAATGCTC	[24]	92°C for 4 min. 94°C for 1 min. 52°C for 1 min. 64°C for 1 min. 64°C for 8 min. } 35x
		trnHf_05	CGCGCATGGTGGATTCACAATCC	[25]	

Table 3.
List of primers used for amplification of different loci and their PCR conditions.

DNA sequencer at AgriGenome labs facility, Kochi, India. Each DNA barcode region was sequenced.

The different universal primers used in this study for the amplification are shown in **Table 4** along with the amplified product size.

The final edited sequences are provided in FASTA format below for each of the successful sequencing reactions:

>Vanilla S matk.

```
TCTCACATTTAAATTATGTGTCAGATCTACTAATACCCTATCCCATACATC
TGGAAATCTTAGTTCAAATTCTTCAATGCTGGGTCAAAGATGTTCTTTCTTTG
CATTTATTGCGATTGTTTTTTCACGAATATCATAATTTGAATAGTCTCGTTAC
TTCAAAGAAATCTATTTATGTCTTTTCAAAAATAAATAAAAGATTTTTTTTAT
TCCTACATAATTTTTATGTATATGAATCCGAATATCTATTCCTGTTTCTTCGT
AAACAGTCTTCTTATTTACGATCAACATCTTCTGGAGTGTTTCTTGAACAAA
CACATTTCTATGTAAAAATAGAACATATTCATCTTATAGTAGTAGTGTTG
TAATTCCTTCAAAGGGACCTATGGTTTCTCGAAGATCCTTTCATGCATTAT
GTTTCGATATCAAGGAAAAGCTATTCTGGGTCAAAGGAACTCTTATTCTGG
TGAATAAATGGAAATATTATCTTATTAATTTTTGGCAATCTTATTTTCACTTT
TGGTCTCAACCAGATAGGATCTATAGAAAGCAATTCTCCGACTATTCCTTTT
CTTTCCTGGGGTATTTTTCAAGTGTATTAATAAATACTTTGGTAGTCAGAAA
TCAAATGCTAGAGAATTGCTTCTCATAAATACTCCGACTCAGAAATTAGAT
ACCATAGCCCCGGTTATTTCTCTTATTGGATCCTTGTCGAAGGCAAAATTTT
GTACGTTAATGGGTCATCCCATTAGTAAACCGATCTGGACCGATTATCGGA
TTCTGAGATTATTGATCGATTTTGTCTGAATATGTAGAAATCTTTGTCTGTTATC
ACAGTGGATCCTCAAAAAAACAGGTTT.
```

>VG matk.

```
TTCTCACATTTAAATTATGTGTCAGATCTACTAATACCCTATCCCATACAT
CTGGAAATCTTAGTTCAAATTCTTCAATGCTGGGTCAAAGATGTTCTTTCTTT
GCATTTATTGCGATTGTTTTTTCACGAATATCAGAATTTGAATAGTCTCGTTA
CTTCAAAGAAATCTATTTATGTCTTTTCAAAAAAATAAAAGATTCTTTTTA
TTCCTACATAATTTTTATGTATATGAATTCGAATATCTATTCATGTTTCTTCG
TAAACAGTCTTCTTATTTACGATCAACATCTTCTGGAGTGTTTCTTGAACAAA
CACATTTTTATGGAAAAATAGAACATATTCATCTTATAGTAGTAGTGTTT
TAATTCCTTAAAAAGCGACCTATGGTTTCTCGAAGATCCTTTCATGCATTAT
GTTTCGATATCAAGGAAAAGCTATTCTGGGTCAAAGGAACTCTTATTCTGT
TGAATAAATGGAAATATTATATTATTTATTTTTGCAATCTTATTTTCACTTT
TGGTCTCAACCAGATAGGATCTATAGAAAGCAATTCTCTGACTATTCCTTTT
CTTTCCTGGGGTATTTTTCAAGTGTATTAATAAATACTTTGGTAGTCAGAAA
TCAAATGCTAGGGAATTGCTTCTCATAAATATTCCGATTCAGAAATTAGAT
ACCACAGCCCCGGTGATTTCTCTTATTGGATCCTTGTCGAAGGCAAAATTTT
GTACGTTAATGGGTCATCCCATTAGTAAACCGATCTGGACTGATTTATCGGA
TTCTGAGATTATTGATCGATTTTGTCTGAATATGTAGAAATCTTTGTCTGTTATC
ACAGTGGA.
```

>VP matk.

```
TTCTCACATTTAAATTATGTGTCAGATCTACTAATACCCTATCCCATACAT
CTGGAAATCTTAGTTCAAATTCTTCAATGCTGGGTCAAAGATGTTCTTTCTTT
GCATTTATTGCGATTGTTTTTTCACGAATATCAGAATTTGAATAGTCTCGTTA
CTTCAAAGAAATCTATTTATGTCTTTTCAAAAAAATAAAAGATTCTTTTTA
TTCCTACATAATTTTTATGTATATGAATTCGAATATCTATTCATGTTTCTTCG
TAAACAGTCTTCTTATTTACGATCAACATCTTCTGGAGTGTTTCTTGAACAAA
CACATTTTTATGGAAAAATAGAACATATTCATCTTATAGTAGTAGTGTTT
TAATTCCTTAAAAAGCGACCTATGGTTTCTCGAAGATCCTTTCATGCATTAT
GTTTCGATATCAAGGAAAAGCTATTCTGGGTCAAAGGAACTCTTATTCTGT
TGAATAAATGGAAATATTATATTATTTATTTTTGCAATCTTATTTTCACTTT
```


S. No	Plant Name	Matk	ITS		rbcL Primers				trnH-psbA
		(product size)	(product size)	(product size)	(product size)	(product size)	(product size)	(product size)	(product size)
1	V1	390F/1326R (1000 bp)	ITS4/ITS5 (800 bp)	—	rbcL_1F/rbcL_724R (NA*)	rbcLa_f/rbcLa_r (700 bp)	rbcLa_f/rbcLa_SI_Rev (600 bp)	rbcLa_f/rbcLaj634R (650 bp)	psbA3_f/trnHf_05 (750 bp)
2	VG	390F/1326R (1000 bp)	ITS4/ITS5 (800 bp)	ITS-P5/ITS-u4 (800 bp)	rbcL_1F/rbcL_724R (NA*)	rbcLa_f/rbcLa_r (NA*)	rbcLa_f/rbcLa_SI_Rev (600 bp)	rbcLa_f/rbcLaj634R (650 bp)	psbA3_f/trnHf_05 (800 bp)
3	VP	390F/1326R (1000 bp)	ITS4/ITS5 (800 bp)	ITS-P5/ITS-u4 (800 bp)	rbcL_1F/rbcL_724R (NA*)	rbcLa_f/rbcLa_r (NA*)	rbcLa_f/rbcLa_SI_Rev (600 bp)	rbcLa_f/rbcLaj634R (650 bp)	psbA3_f/trnHf_05 (800 bp)

*NA, No amplification.
The Bold text indicates successful sequencing was done for these samples.
The ITS region was very problematic while sequencing and only V1 was completed, while sequencing is pending for VG and VP samples.
Loci rbcL and trnH-psbA has been successfully amplified but has not yet been sent for sequencing.

Table 4.
List of primers pairs used for amplification of different barcode loci and its estimated product sizes in agarose gel, for the Vanilla species under study.

TGGTCTCAACCAGATAGGATCTATAGAAAGCAATTCTCTGACTATTCCTTTT
CTTTCCTGGGGTATTTTTCAAGTGTATTAATAAATACTTTGGTAGTCAGAAA
TCAAATGCTAGGGAATTGCTTTCTCATAAATATTCCGATTCAGAAATTAGAT
ACCACAGCCCCGGTGATTTCTCTTATTGGATCCTTGTCGAAGGCAAAATTTT
GTACGTTAATGGGTCATCCCATTAGTAAACCGATCTGGACTGATTTATCGGA
TTCTGAGATTATTGATCGATTTTGTCTGAATATGTAGAAATCTTTGTCTGTTATC
ACAGTGA.

>VS ITS.

AGTGGAAGTAAAAGTCGTAACAAGGTTTCCGTAGGTGAACCTGCGGAAG
GATCATTGACGAGAGCTATGACTGATCGAGTGATCTGTGCAACCTGTGGGG
GTGCGACGGCTGTTTGATGTCGCATTCTTCCATCGCAGAGCTCCTGCTTCCA
GGGGGAGCTCGATGCTGTGGGGGGGATAAACAACAGCCTATGGGCGTGGTCA
TACGCCAAGGGAGAGCAAATGTAAAGCCGCCAACGGGTGTGTTGTGCGTCG
CCAGGCCCAGTGGGGTATGGCAAACGAACACTGAACGACTCTCGACAACGG
ATATCTTGCTCTCGCATCGATGAAGAACGCAGCGAAATGCGATACGTGTTG
TGAATTGTAGAATCCCGTGAACCATCCATTTTTTTGAACGCAAGTTGCGCCCG
AGGATGCAAGCCGAGGGCACTCCTGCATGGGTGTAATGCGTTCTGTCTGCTC
CTCGCGCAGGCATGGAATCGTTGGTTTAGATCAGCGGGCCCCTCGCCAGGAT
GCGATCGATGGCACCCTGTGCTACGGCATGGCGTGTTCAAGCGTTGGGCGA
TGGTCGGCTGTAGACACGGCAAGAGGTGGATGCCACCGAGTGTTGTGGTGT
TGGCCAGTAGGAACCGATGTTGCAGTGCGACAAGGTGATGCCCCTTGCAAA
TCCAACCTCCATGCTCCATGGTGTGGAATCGTGACCCCATGTTAGGTGAGGCT
ACCCGCTGAGTTTAAGCATATCAATAAGCGGA.

5. Result and discussion

5.1 Presence of important agronomic characters

Among the different species of vanilla studied *Vanilla aphylla* Blume and *V. pilifera* Holtt., flowered synchronously (**Figure 5**). *V. aphylla* occurs naturally in South India (**Figure 6**) and *V. pilifera* (**Figure 7**) in Assam, Northeast India. Flowers of both the species opened sequentially and lasted for one day in *V. pilifera*, whereas it lasted for 2 days in *V. aphylla*. In the former, signs of fruit set were observed even without pollination (**Figures 1 and 8**) whereas *V. aphylla* flowers did not set fruit (**Figure 3**), ruling out the possibility of natural fruit set in this species, which is thus similar to *V. planifolia* (**Table 5**).

Figure 5.
V. pilifera (T) inflorescence in comparison with that of *V. aphylla* (B). Arrow Indicates signs of natural fruit set without pollination in *V. pilifera*.

Figure 6.
V. pilifera flower with a leaf (which develops at maturity).

Figure 7.
Vine of V. aphylla in bloom.

Figure 8.
Cross section of the ovary pedicel of *V. aphylla* without pollination, and after 24 hrs.

Species	Disease resistance	Fragrance	Natural seed set	Crossability	Fruit size	Flower life (hours)
<i>V. aphylla</i>	Tolerant to <i>Fusarium oxysporum</i>	—	Not seen	Crossable to <i>V. pilifera</i> and <i>V. planifolia</i>		> 36
<i>V. pilifera</i>	—	Highly fragrant	Symptoms seen	Crossable to <i>V. aphylla</i>	Larger	~24
<i>V. planifolia</i>	Susceptible		Not seen	Crossable to <i>V. aphylla</i>		< 24

Table 5.
Important agronomic characters.

Cross sections of the ovary pedicel were observed after closing of the flowers. Persistent perianth is characteristic to the genus and also indicative of effective pollination. In flowers where pollination is not effected, the perianth is shed after the flower closes. Perianth in *V. pilifera* were found to persist even without pollination and the cross sections indicated initiation of seed set (**Figures 9 and 10**), whereas *V. aphylla* did not show any indications (**Figures 11 and 12**). Since rostellum is present in both the species, natural pollination without an aid is ruled out. It can be suspected, that the fragrance of the *V. pilifera* flowers attracts insects (which were found to frequent the flowers often) to visit them and bring about effective pollination.

Pollinations both self and interspecific hybridizations between the two species were done and fruits set was observed (**Figure 13**).

5.2 Sequence analysis

General observations from the experiment

- 1. The matK sequences of VG and VP are identical.

Figure 9.
Cross section of the ovary pedicel of V. aphylla without pollination, and after 24 hrs.

Figure 10.
C.S. of ovary-pedicel of V. pilifera without pollination, showing indications of seed set.

Figure 11.
C.S. of ovary-pedicel of V. pilifera without pollination, showing indications of seed set.

Figure 12.
Flowers of V. pilifera in comparison with V. aphylla.

Figure 13.
V. aphylla inflorescence with fruit set after interspecific hybridization.

2. The matK sequence of V1 was different from VG/VP at 21 nucleotide positions as shown in **Table 6** below.

Nucleotide position	1	1	2	2	2	3	3	3	3	3	4	4	4	5	5	6	6	6	6	6	7
	3	9	0	3	5	2	2	6	7	8	7	9	9	0	6	4	6	6	8	9	7
	7	0	3	8	2	3	8	7	7	3	1	2	8	5	3	0	1	7	5	5	3
VS matk	T	T	T	C	C	C	T	G	C	G	G	C	A	G	C	A	C	C	T	T	C
VG matk	G	A	C	T	A	T	G	T	A	C	T	A	T	T	T	G	T	T	C	G	T
VP matk	G	A	C	T	A	T	G	T	A	C	T	A	T	T	T	G	T	T	C	G	T

Table 6.
Matk sequence analysis.

3. Blast search of the matk sequences of VG/VP in the NCBI blast search gave 100% match with *Vanilla planifolia* (Accession No. KJ566306.1), as in the NCBI search results shown below.

Description	Max score	Total score	Query cover	E value	Ident	Accession
<i>Vanilla planifolia</i> chloroplast, complete genome	1563	1563	100%	0.0	100.00%	KJ566306.1
<i>Vanilla planifolia</i> chloroplast, complete genome	1546	1546	100%	0,0	99.65%	MF197310.1
<i>Vanilla planifolia</i> tRNA-Lys (trnK) gene, partial sequence; and maturase K (matK) gene, compl	1524	1524	100%	0.0	99.17%	JN181462.1
<i>Vanilla planifolia</i> maturase K (matK) chloroplast pseudogene, partial sequence	1507	1507	100%	0.0	98.82%	AF263687.1
<i>Vanilla planifolia</i> chloroplast matK pseudogene	1507	1507	100%	0.0	98.82%	AJ310079.1
<i>Vanilla planifolia</i> plastid partial matK gene for maturase K, specimen voucher Chase O-199 K	1423	1423	95%	0.0	98.40%	AJ581443.1
<i>Vanilla somae</i> voucher KFBG290 maturase K (matK) gene, partial cds: plastid	1419	1419	100%	0.0	96.93%	KY966974.1
<i>Vanilla aphylla</i> chloroplast DNA, complete genome	1419	1419	99%	0.0	97.03%	LC085348.1
<i>Vanilla piliifera</i> voucher V5 maturase K (matK) gene, partial cds: chloroplast	1354	1354	99%	0,0	95.64%	FJ816099.1
<i>Vanilla siamensis</i> voucher V2 maturase K (matK) gene, partial cds: chloroplast	1315	1315	97%	0,0	95.23%	FJ816097.1
<i>Vanilla planifolia</i> voucher SBB-0324 maturase K (matK) gene, partial cds: chloroplast	1314	1314	84%	0.0	100.00%	JN004635.1
<i>Vanilla planifolia</i> isolate AD7LN25 maturase K (matK) qene. Partial cds: chloroplast	1284	1284	89%	0.0	97.11%	MF349972.1

4. Blast search of the matk sequences of VS in the NCBI blast search gave maximum identity with *Vanilla somae* (Accession No.KY966974.1). See the NCBI search results below.

Description	Max score	Total score	Query cover	E value	Ident	Accession
<i>Vanilla somae</i> voucher KFBG290 maturase K (matK) gene, partial cds: plastid	1570	1570	99%	0.0	99.54%	KY966974.1
<i>Vanilla aphylla</i> chloroplast DNA, complete genome	1570	1570	99%	0.0	99.54%	LC085348.1
<i>Vanilla piliifera</i> voucher V5 maturase K (matK) gene, partial cds: chloroplast	1489	1489	100%	0.0	97.81%	FJ816099.1

Description	Max score	Total score	Query cover	E value	Ident	Accession
<i>Vanilla pompona</i> chloroplast. Complete genome	1474	1474	100%	0,0	97.45%	MF197310.1
<i>Vanilla planifolia</i> chloroplast. Complete genome	1469	1469	100%	0.0	97.34%	KJ566306.1
<i>Vanilla planifolia</i> tRNA-Lys (trnK) gene, partial sequence: and maturase K (matK) gene, complel	1452	1452	100%	0.0	96.99%	JN181462.1
<i>Vanilla siamensis</i> voucher V2 maturase K (matK) gene, partial cds: chloroplast	1447	1447	98%	0.0	97.33%	FJ816097.1
<i>Vanilla planifolia</i> maturase K (matK) chloroplast pseudogene, partial sequence	1435	1435	100%	0.0	96.64%	AF263687.1
<i>Vanilla planifolia</i> chloroplast matK pseudogene	1435	1435	100%	0.0	96.64%	AJ310079.1
<i>Vanilla planifolia</i> isolate AD7LN25 maturase K (matK) gene, partial cds: chloroplast	1408	1408	90%	0.0	99.23%	MF349972.1
<i>Vanilla roscheri</i> voucher NMK:838.10216 maturase K (matK) gene, partial cds: chloroplast	1395	1395	93%	0.0	97.89%	KU748308.1
<i>Vanilla aphylla</i> voucher V1 maturase K (matK) gene, partial cds: chloroplast	1391	1391	96%	0.0	96.79%	FJ816096.1

5. Blast search of the ITS sequences of VS in the NCBI blast search gave maximum identity with *Vanilla shenzhenica* (Accession No. JF796930.1). See the NCBI search results below.

Description	Max score	Total score	Query cover	E value	Ident	Accession
<i>Vanilla somae</i> voucher KFBG290 maturase K (matK) gene, partial cds: plastid	1570	1570	99%	0.0	99.54%	KY966974.1
<i>Vanilla aphylla</i> chloroplast DNA, complete genome	1570	1570	99%	0.0	99.54%	LC085348.1
<i>Vanilla pilifera</i> voucher V5 maturase K (matK) gene, partial cds: chloroplast	1489	1489	100%	0.0	97.81%	FJ816099.1
<i>Vanilla pompona</i> chloroplast. Complete genome	1474	1474	100%	0,0	97.45%	MF197310.1
<i>Vanilla planifolia</i> chloroplast. Complete genome	1469	1469	100%	0.0	97.34%	KJ566306.1
<i>Vanilla planifolia</i> tRNA-Lys (trnK) gene, partial sequence: and maturase K (matK) gene, complel	1452	1452	100%	0.0	96.99%	JN181462.1
<i>Vanilla siamensis</i> voucher V2 maturase K (matK) gene, partial cds: chloroplast	1447	1447	98%	0.0	97.33%	FJ816097.1
<i>Vanilla planifolia</i> maturase K (matK) chloroplast pseudogene, partial sequence	1435	1435	100%	0.0	96.64%	AF263687.1
<i>Vanilla planifolia</i> chloroplast matK pseudogene	1435	1435	100%	0.0	96.64%	AJ310079.1

Description	Max score	Total score	Query cover	E value	Ident	Accession
Vanilla planifolia isolate AD7LN25 maturase K (matK) gene, partial cds: chloroplast	1408	1408	90%	0.0	99.23%	MF349972.1
Vanilla roscheri voucher NMK:838.10216 maturase K (matK) gene, partial cds: chloroplast	1395	1395	93%	0.0	97.89%	KU748308.1
Vanilla aphylla voucher V1 maturase K (matK) gene, partial cds: chloroplast	1391	1391	96%	0.0	96.79%	FJ816096.1

6. The ITS sequences matching with different Vanilla sp. were downloaded and subjected to analysis using MEGA7.0 software (Table 7).

	VS ITS	JF796930.1_Vanilla_shezenhica	KY966687.1_Vanilla_somae	AF151006.1_Vanilla_aphylla	JF825978.1_Vanilla_siamensis	FJ425830.1_Vanilla_imperialis	FJ425840.1_Vanilla_roscheri	FJ425835.1_Vanilla_barbellata	FJ425834.1_Vanilla_africana	EU498163.1_Vanilla_bahiana	GQ867241.1_Vanilla_planifolia	GQ867237.1_Vanilla_pompona	AF391785.1_Vanilla_hirsuta	EU498165.1_Vanilla_edwallii
VS ITS	0													
JF796930.1_Vanilla_shezenhica	20	0												
KY966687.1_Vanilla_somae	20	0	0											
AF151006.1_Vanilla_aphylla	27	19	19	0										
JF825978.1_Vanilla_siamensis	42	39	39	41	0									
FJ425830.1_Vanilla_imperialis	40	33	33	41	47	0								
FJ425840.1_Vanilla_roscheri	41	34	34	45	53	40	0							
FJ425835.1_Vanilla_barbellata	55	48	48	55	66	57	58	0						
FJ425834.1_Vanilla_africana	63	51	51	56	66	56	60	76	0					
EU498163.1_Vanilla_bahiana	96	89	89	96	101	97	99	107	100	0				
GQ867241.1_Vanilla_planifolia	107	100	100	107	119	106	113	110	110	45	0			
GQ867237.1_Vanilla_pompona	99	92	92	99	104	98	103	107	104	20	45	0		
AF391785.1_Vanilla_hirsuta	101	92	92	102	114	101	102	108	100	39	31	43	0	
EU498165.1_Vanilla_edwallii	132	133	133	136	143	136	143	141	146	139	150	136	146	0

Table 7. Estimates of evolutionary divergence between sequences.

The number of base differences per sequence from between sequences are shown. The analysis involved 14 nucleotide sequences. All ambiguous positions were removed for each sequence pair. There was a total of 657 positions in the final dataset. Evolutionary analyses were conducted in MEGA7 (Figure 14).

Figure 14. Phylogenetic analysis of the ITS sequences inferred using the neighbor-joining method, computed using the Kimura 2-parameter method and are in the units of the number of base substitutions per site. The analysis involved 14 nucleotide sequences. All ambiguous positions were removed for each sequence pair. There was a total of 657 positions in the final dataset. The analyses were conducted in MEGA7.

6. Conclusions

The analysis involved 14 nucleotide sequences. All ambiguous positions were removed for each sequence pair. There was a total of 657 positions in the final dataset. The analyses were conducted in MEGA7. The phylogeny analysis also revealed the separate clustering offer leafy and leafless species. *Vanilla siamensis*, a leafy species, indicating signs of self-pollination in its wild, in Thailand, clustered with leafless *V. aphylla* species.

The studies further reveal the complexity of the biosynthesis of the natural vanillin synthesis. However, it is to be further analyzed whether leafy character is associated with enhanced photosynthetic products that indirectly affect the vanillin synthesis too. This reiterates the need for conservation of the genetic resources [12] of Vanilla across the continents, for implementing meaningful breeding programs, to enhance vanillin productivity in addition to disease resistance and reproductive behavior.

Acknowledgements

The first author acknowledges the former Directors of Indian Institute of Spices Research, Kozhikode, Kerala from where she initiated these studies. The authors are thankful for the financial support rendered by University Grants Commission (MRP and CPE funds), DST-FIST funds, for taking up these research initiatives, in their laboratory and express gratitude to Scientists at Centre for Medicinal Plant Research, Kottakkal, Malappuram, for support in Sequencing work.

Author details

Minoo Divakaran* and N.T. Fathima Rafieah
Department of Botany, Providence Women's College, Kozhikode, Kerala, India

*Address all correspondence to: minoodivakaran@gmail.com

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Correll, D. S. (1953). *Vanilla*-its botany, history, cultivation and economic importance. *Economic Botany*, 7(4), 291–358. DOI:10.1007/bf02930810
- [2] Bouriquet G (1954) *Le vanillier et la vanille dans le monde*. Encyclope'die Biologique XLVI. Lechevalier, Paris VI
- [3] Swamy, B. G. L. (1947). On the life-history of *Vanilla planifolia*. *Botanical Gazette*, 108(3), 449–456. DOI:10.1086/335429
- [4] Purseglove JW, Brown EG, Green CL, Robbins SRJ (eds.) 1981. *Spices*. Vol. 2. Longman Inc., New York, pp 644-735.
- [5] Spices Board 2000. *Vanilla Status paper*. Spices Board, Cochin, India, pp. 33.
- [6] Lubinsky P (2004) *Vanilla diversity in Mexico*. In: Bakto Fla-vors, Rutgers University (eds) *Vanilla 2004 Europe*, Second International Congress, Cannes, France, 30 Sept–01 Oct 2004.
- [7] Hu, Y., Resende, M. F. R., Jr., Bombarely, A., Brym, M., Bassil, E., and Chambers, A. H. (2019). Genomics-based spanersity analysis of *Vanilla* species using a *Vanilla planifolia* draft genome and genotyping-by-sequencing. *Scientific Reports*, 9(1). DOI:10.1038/s41598-019-40144-1
- [8] Minoo, D., Jayakumar, V. N., Veena, S. S., Vimala, J., Basha, A., Saji, K. V., Nirmal Babu, K., and Peter, K. V. (2007). Genetic variations and interrelationships in *Vanilla planifolia* and few related species as expressed by RAPD polymorphism. *Genetic Resources and Crop Evolution*, 55(3), 459–470. DOI: 10.1007/s10722-007-9252-3
- [9] Satish K and Manilal K.S. 1993. *Vanilla aphylla* Blume. *Amer. Orch. Soc. Bull.* 62, 394–397.
- [10] Seidenfaden G. 1978. Orchid genera in Thailand VI Neottioideae. *Dansk. Bot. Ark.* 32, 138–146.
- [11] Franklin, W.M., 1963. Chromosome number and behavior in a *Vanilla* hybrid and several *Vanilla* species. *Bull. Torrey Bot. Club* 90, 416–417
- [12] Minoo, D., Nirmal Babu, K., and Peter, K. V. (2006). Conservation of *Vanilla* species, in vitro. *Scientia Horticulturae*, 110(2), 175–180. DOI: 10.1016/j.scienta.2006.07.003
- [13] Minoo, D., Nirmal Babu, K., Ravindran, P. N., and Peter, K. V. (2006). Interspecific hybridization in vanilla and molecular characterization of hybrids and selfed progenies using RAPD and AFLP markers. *Scientia Horticulturae*, 108(4), 414–422. DOI: 10.1016/j.scienta.2006.02.018
- [14] Gallage, N. J., Hansen, E. H., Kannangara, R., Olsen, C. E., Motawia, M. S., Jørgensen, K., Holme, I., Hebelstrup, K., Grisoni, M., and Møller, B. L. (2014). Vanillin formation from ferulic acid in *Vanilla planifolia* is catalysed by a single enzyme. *Nature Communications*, 5(1). DOI:10.1038/ncomms5037
- [15] Cuenoud, P., Savolainen, V., Chatrou, L. W., Powell, M., Grayer, R. J., and Chase, M. W. (2002). Molecular phylogenetics of Caryophyllales based on nuclear 18S rDNA and plastid rbcL, atpB, and matK DNA sequences. *American Journal of Botany*, 89(1), 132–144. DOI:10.3732/ajb.89.1.132
- [16] White, T. J., Bruns, T., Lee, S., and Taylor, J. (1990). AFmplification and direct sequencing of fungal ribosomal rna genes for phylogenetics. In *PCR Protocols* (pp. 315–322). Elsevier. doi: 10.1016/b978-0-12-372180-8.50042-1
- [17] Sang, T., Crawford, D. J., and Stuessy, T. F. (1995). Documentation of

- reticulate evolution in peonies (*Paeonia*) using internal transcribed spacer sequences of nuclear ribosomal DNA: Implications for biogeography and concerted evolution. *Proceedings of the National Academy of Sciences*, 92(15), 6813–6817. DOI:10.1073/pnas.92.15.6813
- [18] Cheng, T., Xu, C., Lei, L., Li, C., Zhang, Y., and Zhou, S. (2015). Barcoding the kingdom Plantae: New PCR primers for ITS regions of plants with improved universality and specificity. *Molecular Ecology Resources*, 16(1), 138–149. DOI:10.1111/1755-0998.12438
- [19] Olmstead, R. G., Michaels, H. J., Scott, K. M., and Palmer, J. D. (1992). Monophyly of the Asteridae and identification of their major lineages inferred from DNA sequences of *rbcL*. *Annals of the Missouri Botanical Garden*, 79(2), 249. DOI:10.2307/2399768
- [20] Fay, M. F., Swensen, S. M., and Chase, M. W. (1997). Taxonomic affinities of *Medusagyne oppositifolia* (Medusagynaceae). *Kew Bulletin*, 52(1), 111. DOI:10.2307/4117844
- [21] Kress, W. J., and Erickson, D. L. (2007). A two-locus global DNA barcode for land plants: The coding *rbcL* gene complements the non-coding *trnH-psbA* spacer region. *PLoS ONE*, 2(6), e508. DOI:10.1371/journal.pone.0000508
- [22] Kress, W. J., Erickson, D. L., Jones, F. A., Swenson, N. G., Perez, R., Sanjurjo, O., and Bermingham, E. (2009a). Plant DNA barcodes and a community phylogeny of a tropical forest dynamics plot in Panama. *Proceedings of the National Academy of Sciences*, 106(44), 18621–18626. DOI:10.1073/pnas.0909820106
- [23] Fazekas, A. J., Burgess, K. S., Kesanakurti, P. R., Graham, S. W., Newmaster, S. G., Husband, B. C., Percy, D. M., Hajibabaei, M., and Barrett, S. C. H. (2008). Multiple multilocus DNA barcodes from the plastid genome discriminate plant species equally well. *PLoS ONE*, 3(7), e2802. DOI:10.1371/journal.pone.0002802
- [24] Sang, T., Crawford, D. J., and Stuessy, T. F. (1997). Chloroplast DNA phylogeny, reticulate evolution, and biogeography of *Paeonia* (Paeoniaceae). *American Journal of Botany*, 84(8), 1120–1136. DOI:10.2307/2446155
- [25] Tate, J.A., Simpson, B.B. (2003). Paraphyly of *Tarasa* (Malvaceae) and diverse origins of the Polyploid species. *Systematic Botany*. 28. 723-737. DOI: 10.1043/02-64.1