We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Introductory Chapter: The Important Physiological Characteristics and Industrial Applications of Acidophiles

Linxu Chen, Jianqun Lin and Jianqiang Lin

1. The definition of acidophiles

Acidophiles are an important category of extremophiles that are defined by the environmental conditions in which they grow optimally. Acidophile is a broad definition that organisms can grow preferentially in environments with a pH at below 6. In 2007, Johnson proposed a generally accepted classification standard according to the optimal pH. The organisms with optimal pH at 3 or below are classified as extreme acidophiles, and those with an optimal pH of 3–5 are moderate acidophiles [1]. Although some organisms can grow at a pH lower than 5, they are recognized as acid-tolerant species because of their pH optima above 5. The research history of acidophiles started in the discovery of a sulfur-oxidizing bacteria isolated from a compost sample mixed with sulfur, rock phosphate, and soil by Waksman and Joffe in 1922 [2]. This bacterium has an optimal growth pH at 2.0–2.8 and is a strict autotroph that obtains energy by oxidizing inorganic sulfur substances (elemental sulfur, thiosulfate, and hydrogen sulfide). This bacterium was named as *Thiobacillus thiooxidans* by Waksman and Joffe, and later was reclassified as *Acidithiobacillus thiooxidans* by Kelly and Wood in 2000 [3]. With the development of microbiology and gene sequencing technology, more and more acidophiles have been discovered, identified, and sequenced. Until now, the most acidophilic organisms are from an archaeal genus of *Picrophilus*, firstly isolated from acidic hot springs and dry hot soil in Hokkaido in Japan [4]. Members in *Picrophilus* have optimal pH at 0.7 and the ability to grow at a pH of 0. Moreover, acidophiles are involved not in the domains of *Bacteria* and *Archaea*, but also in the *Eukarya* domains, such as some acidophilic fungi, algae, and yeast distributed in the acid mine environments.

2. The typical acidophilic bacteria and the applications of acidophiles

Acidithiobacillus is a kind of extensive research and wide application of gram-negative acidophiles. Members in this genus are broadly existed in the sulfur-containing acidic environments on land or in the sea, such as acid mine drainage (AMD), iron-sulfur mineral mines, hot springs, and sediments [5–10]. *Acidithiobacillus* spp., as the important sulfur and iron-oxidizers, participate in the element cycles of sulfur and iron, and promote the acid environment generation and acid microecosystem formation. All *Acidithiobacillus* strains have the capability of oxidizing various reduced inorganic sulfur compounds (RISCs)

Trait	<i>A. ferrooxidans</i>	<i>Acidithiobacillus ferrivorans</i>	<i>A. ferrophilus</i>	<i>A. ferridurans</i>	<i>Acidithiobacillus thiooxidans</i>	<i>Acidithiobacillus caldus</i>	<i>A. albertensis</i>
Gram stain	—	—	—	—	—	—	—
Cell size (µm)	1.0 × 0.5	2.4 × 0.5	1–2	1–2	1.0–2.0 × 0.5	1.2–1.9 × 0.7	1–2 × 0.4–0.6
Motility	+/–	+	+	+	+	+	+
Growth pH (optimum)	1.3–4.5 (2.0–2.5)	1.9–3.4 (2.5)	1.5– (2.0)	1.4–3.0 (2.1)	0.5–5.5 (2.0–3.0)	1.0–3.5 (2.0–2.5)	0.5–6.0 (3.5–4.0)
Growth T/°C (optimum)	10–37 (30–35)	4–37 (28–33)	5–33 (30)	10–37 (29)	10–37 (28–30)	32–52 (40–45)	10–40 (25–30)
Oxidation of S ⁰ , S ₄ O ₆ ²⁻ , S ₂ O ₃ ²⁻	+	+	+	+	+	+	+
Oxidation of Fe ²⁺	+	+	+	+	—	—	—
Growth on sulfide minerals	+	+	+	+	—	—	—
Growth on hydrogen	+	(+)	—	+	—	+	NR
Anaerobic growth with Fe ³⁺	+	+	+	+	—	—	—
N ₂ fixation	+	+	NR	NR	—	—	—
Mol% G + C	58–59	55–56	57.4	58.4	52	63–64	61.5
Thiosulfate- metabolic pathways	TSD enzyme; S ₄ I pathway.	Sox system; TSD enzyme; S ₄ I pathway.	NR	TSD enzyme; S ₄ I pathway.	Sox system; S ₄ I pathway.	Sox system; S ₄ I pathway.	Sox system; S ₄ I pathway.

+, positive; –, negative; +/–, the positive or negative result from different reports; (+), some strains have the ability to oxidize hydrogen; NR, not reported; Tm, temperature.

Table 1.
Taxonomic traits of species in the genus of *Acidithiobacillus*.

and elemental sulfur, and some of them also have ferrous iron oxidation ability [11]. By the oxidation of sulfur and ferrous, *Acidithiobacillus* spp. obtains electrons to generate the bioenergy (ATP) and reducing power (NADH/NADPH) to fix carbon dioxide for autotrophic growth. More and more species have been identified based on their physiological characters and 16S rRNA gene sequences (**Table 1**) [2, 12–16]. Species in *Acidithiobacillus* can be divided into two groups according to their energy-substrates: the sulfur-oxidizing only species, including *A. thiooxidans*, *Acidithiobacillus caldus* and *A. albertensis*, and the sulfur and ferrous-oxidizing species, including *A. ferrooxidans*, *Acidithiobacillus ferrivorans*, *A. ferriphilus*, and *A. ferridurans* (**Table 1**).

Acidithiobacillus spp. and other chemoautotrophic acidophilic bacteria have an important application in bioleaching. The bioleaching technology is originated from the biohydrometallurgy industry, and has become a great potential and broad-prospects in non-ferrous metal extraction (golden, silver, copper et al.) from various sulfide ores. *Acidithiobacillus* spp. have the remarkable capabilities of metabolizing the sulfur and iron in ores and adapting to extremely acidic environments, thus they have become the most active and preponderant bacteria used in biomining [17, 18]. *A. ferrooxidans*, *A. thiooxidans*, and *A. caldus* are the wide used ore leaching species in biomining for mineral extraction from ores [19, 20]. In recent years, based on their abilities to produce acid and heavy leaching metals, *Acidithiobacillus* spp. have been used from biohydrometallurgy to the treatment of wastes containing heavy metals, such as sewage sludge, spent household batteries, mine tailings, and printed circuit boards [21–25]. Moreover, these bacteria have been widely studied in microbial desulfurization of coal and gas [26–28]. In a word, the great application values of *Acidithiobacillus* spp. have been exploited from the biohydrometallurgy industry to the environmental pollution treatments.

3. The physiological feature of chemoautotrophic acidophiles

Sulfur oxidation is a characteristic physiological feature for many acidophilic microorganisms and is an important biochemical process that promotes the generation of the acid environment and the formation of acidophilic microbial communities. *Acidithiobacillus* spp., as the first-discovered and the most widespread used acidophile, has been attracted extensive attention and has been used as model sulfur-oxidizing bacteria to research microbial sulfur metabolism [11, 29–40]. The oxidation states of element sulfur are range from -2 to $+6$, resulting in different kinds of RISCs (tetrathionate ($S_4O_6^{2-}$), thiosulfate ($S_2O_3^{2-}$), sulfite (SO_3^{2-}), sulfide (S^{2-}) et al.), and elemental sulfur (S^0). Thus, many microbes, particularly autotrophic sulfur-oxidizing microbes, have evolved a variety of enzymes and proteins participating in the oxidation of RISCs and S^0 . Research shows *Acidithiobacillus* spp. have a high-efficient and sophisticated sulfur-metabolizing network that could oxidize RISCs and S^0 to sulfate. Based on metabolic substrates, the sulfur-metabolic enzymes in *Acidithiobacillus* spp. could be categorized as elemental sulfur oxidation enzymes, enzymes in thiosulfate oxidation pathways, sulfide oxidation enzymes, and sulfite oxidation enzymes. These enzymes work cooperatively in different cellular compartments to oxidize the RISCs and S^0 to the final product sulfate (**Figures 1** and **2**) [11]. As shown in **Figure 1**, the extracellular elemental sulfur (S_8) oxidation in *A. caldus* starts from the activation and transportation of S_8 by special outer-membrane proteins (OMP), generating the persulfide sulfane sulfur in the periplasm; then the persulfide sulfane sulfur is oxidized to sulfite that can directly enter the sulfur oxidizing enzyme (Sox) system or form $S_2O_3^{2-}$ via a nonenzymatic reaction; the generated thiosulfate is then metabolized by the truncated Sox pathway or catalyzed by thiosulfate:quinol oxidoreductase (TQO or DoxDA) to

Figure 1. The model of sulfur oxidation in *Acidithiobacillus caldus*. OMP, outer-membrane proteins; TQO, thiosulfate quinone oxidoreductase; TetH, tetrathionate hydrolase; SQR, sulfide:Quinone oxidoreductase; PDO, persulfide dioxigenase; SOR, sulfur oxygenase reductase; TST, rhodanese; HDR, Hdr-like complex; SAT, ATP sulfurylase; bd, bo₃, terminal oxidases; QH₂, quinol pool; NADH, NADH dehydrogenase complex I.

Figure 2. The model of sulfur oxidation in *A. ferrooxidans*. OMP, outer-membrane proteins; TQO, thiosulfate quinone oxidoreductase; TSD, thiosulfate dehydrogenase; TetH, tetrathionate hydrolase; SQR, sulfide:Quinone oxidoreductase; PDO, persulfide dioxygenase; HDR, Hdr-like complex; SAT, ATP sulfurylase; bd, *bo*₃, terminal oxidases; QH₂, quinol pool; NADH, NADH dehydrogenase complex I.

generate $\text{S}_4\text{O}_6^{2-}$; $\text{S}_4\text{O}_6^{2-}$ is further hydrolyzed by tetrathionate hydrolase (TetH); the H_2S produced during the activation of S_8 can be oxidized by sulfide:quinone oxidoreductase (SQR) located in the inner membrane; the periplasmic elemental sulfur (S_0) produced from Sox pathway, tetrathionate hydrolysis and sulfide oxidation, could be re-activated at the outer membrane region, or be mobilized into the cytoplasm

where Sn could be used by cytoplasmic elemental sulfur oxidation enzyme persulfide dioxygenase (PDO) and Sulfur oxygenase reductase (SOR); the metabolites from the reaction of PDO and SOR could be utilized by cytoplasmic sulfur-metabolic enzymes, including the $\text{S}_2\text{O}_3^{2-}$ metabolism via by rhodanese (TST) and the Hdr-like complex (HDR), the degradation of SO_3^{2-} via the APS pathway and the oxidation of S^{2-} by SQR. During the sulfur metabolic process, the periplasmic sulfur-oxidizing pathways (Sox and TetH) are responsible for electron acquisition, thus they are important for the sulfur metabolism in *A. caldus*. Different from '*A. caldus*' like sulfur metabolism network, some sulfur-oxidizers, such as *A. ferrooxidans*, did not have the Sox pathway, but rather a thiosulfate dehydrogenase (TSD) (**Figure 2**). Interestingly, *A. ferrivorans* possesses both Sox system and TSD enzyme (**Table 1**). The proposal of sulfur metabolism models provides new knowledge and insights in understanding the metabolism and adaptation mechanisms of acidophilic sulfur-oxidizing microorganisms in extreme environments.

4. The significance of studying and understanding acidophiles

Acidophiles, as important extremophiles, have presented important scientific significance and industrial application values. Researches on acidophiles do not only help us understand the diversity and adaptation of life on earth, but also be conducive in developing various new biotechnologies to resolve the problems of resource exploitation, pollution treatment, and human health. This book provides some new breakthroughs and insights on the researches of acidophiles: the two-component system (TCS) in the regulation of sulfur metabolic process; the adaptation mechanisms of acidophiles to low pH; the regulation mechanism and the application strategy of quorum sensing in bioleaching bacteria; *Lactobacillus acidophilus* and its application in the human health.

Author details

Lin Xu Chen*, Jianqun Lin and Jianqiang Lin*
State Key Laboratory of Microbial Technology, Shandong University,
Qingdao, China

*Address all correspondence to: linxuchen@sdu.edu.cn
and jianqianglin@sdu.edu.cn

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Johnson JD. Physiology and ecology of acidophilic microorganisms. In: Gerday C, Glansdorff N, editors. *Physiology and Biochemistry of Extremophiles*. Washington, DC, USA: American Society of Microbiology Press; 2007. pp. 255-270
- [2] Waksman SA, Joffe J. Microorganisms concerned in the oxidation of sulfur in the soil: II. *Thiooxidans*, a new sulfur-oxidizing organism isolated from the soil. *Journal of Bacteriology*. 1922;7(2):239
- [3] Kelly DP, Wood AP. Reclassification of some species of *Thiobacillus* to the newly designated genera *Acidithiobacillus* gen. nov., *Halothiobacillus* gen. nov. and *Thermithiobacillus* gen. nov. *International Journal of Systematic and Evolutionary Microbiology*. 2000;50(2):511-516
- [4] Schleper C et al. *Picrophilus* gen. nov., fam. nov.: A novel aerobic, heterotrophic, thermoacidophilic genus and family comprising archaea capable of growth around pH 0. *Journal of Bacteriology*. 1995;177(24):7050-7059
- [5] Harrison AP Jr. The acidophilic thiobacilli and other acidophilic bacteria that share their habitat. *Annual Review of Microbiology*. 1984;38(1):265-292
- [6] Schrenk MO et al. Distribution of *Thiobacillus ferrooxidans* and *Leptospirillum ferrooxidans*: Implications for generation of acid mine drainage. *Science*. 1998;279(5356):1519-1522
- [7] Jones DS et al. Community genomic analysis of an extremely acidophilic sulfur-oxidizing biofilm. *The ISME Journal*. 2012;6(1):158
- [8] Hua Z-S et al. Ecological roles of dominant and rare prokaryotes in acid mine drainage revealed by metagenomics and meta transcriptomics. *The ISME Journal*. 2015;9(6):1280
- [9] Sharmin S et al. Characterization of a novel thiosulfate dehydrogenase from a marine acidophilic sulfur-oxidizing bacterium, *Acidithiobacillus thiooxidans* strain SH. *Bioscience, Biotechnology, and Biochemistry*. 2016;80(2):273-278
- [10] Nuñez H et al. Molecular systematics of the genus *Acidithiobacillus*: Insights into the phylogenetic structure and diversification of the taxon. *Frontiers in Microbiology*. 2017;8:30
- [11] Wang R et al. Sulfur oxidation in the acidophilic autotrophic *Acidithiobacillus* spp. *Frontiers in Microbiology*. 2019;9:3290
- [12] Temple KL, Colmer AR. The autotrophic oxidation of iron by a new bacterium: *Thiobacillus ferrooxidans*. *Journal of Bacteriology*. 1951;62(5):605
- [13] Hallberg KB, Lindström EB. Characterization of *Thiobacillus caldus* sp. nov., a moderately thermophilic acidophile. *Microbiology*. 1994;140(12):3451-3456
- [14] Xia J-L et al. A new strain *Acidithiobacillus albertensis* BY-05 for bioleaching of metal sulfides ores. *Transactions of Nonferrous Metals Society of China*. 2007;17(1):168-175
- [15] Liljeqvist M, Rzhapishchevska OI, Dopson M. Gene identification and substrate regulation provides insights into sulfur accumulation during bioleaching with the psychrotolerant *Acidithiobacillus ferrivorans*. *Applied and Environmental Microbiology*. 2013;79(3):951-957
- [16] Falagán C, Johnson DB. *Acidithiobacillus ferriphilus* sp. nov., a facultatively anaerobic iron-and

- sulfur-metabolizing extreme acidophile. International Journal of Systematic and Evolutionary Microbiology. 2016;**66**(1): 206-211
- [17] Rawlings D, Tributsch H, Hansford G. Reasons why 'Leptospirillum'-like species rather than *Thiobacillus ferrooxidans* are the dominant iron-oxidizing bacteria in many commercial processes for the biooxidation of pyrite and related ores. Microbiology. 1999;**145**(1):5-13
- [18] Rawlings DE. Characteristics and adaptability of iron-and sulfur-oxidizing microorganisms used for the recovery of metals from minerals and their concentrates. Microbial Cell Factories. 2005;**4**(1):13
- [19] Valdés J et al. *Acidithiobacillus ferrooxidans* metabolism: From genome sequence to industrial applications. BMC Genomics. 2008;**9**(1):597
- [20] Valdés J et al. Comparative genome analysis of *Acidithiobacillus ferrooxidans*, *A. thiooxidans* and *A. caldus*: Insights into their metabolism and ecophysiology. Hydrometallurgy. 2008;**94**(1-4):180-184
- [21] Pathak A, Dastidar M, Sreekrishnan T. Bioleaching of heavy metals from sewage sludge. Journal of Environmental Management. 2009;**90**(8):2343-2353
- [22] Bayat B, Sari B. Comparative evaluation of microbial and chemical leaching processes for heavy metal removal from dewatered metal plating sludge. Journal of Hazardous Materials. 2010;**174**(1-3):763-769
- [23] Arshadi M, Mousavi S. Simultaneous recovery of Ni and Cu from computer-printed circuit boards using bioleaching: Statistical evaluation and optimization. Bioresource Technology. 2014;**174**:233-242
- [24] Nguyen VK et al. Bioleaching of arsenic and heavy metals from mine tailings by pure and mixed cultures of *Acidithiobacillus* spp. Journal of Industrial and Engineering Chemistry. 2015;**21**:451-458
- [25] Rastegar S et al. Bioleaching of V, Ni, and Cu from residual produced in oil fired furnaces using *Acidithiobacillus ferrooxidans*. Hydrometallurgy. 2015;**157**:50-59
- [26] Azizan A, Najafpour G, Harun A. Microbial desulfurization of coal with mixed cultures of *Thiobacillus ferrooxidans* and *Thiobacillus thiooxidans*. In: Proceedings of the 2nd International Conference on Advances in Strategic Technologies-ICAST; 2000
- [27] Charnnok B et al. Oxidation of hydrogen sulfide in biogas using dissolved oxygen in the extreme acidic biofiltration operation. Bioresource Technology. 2013;**131**:492-499
- [28] He H et al. Biodesulfurization of coal with *Acidithiobacillus caldus* and analysis of the interfacial interaction between cells and pyrite. Fuel Processing Technology. 2012;**101**:73-77
- [29] Suzuki I, Werkman C. Glutathione and sulfur oxidation by *Thiobacillus thiooxidans*. Proceedings of the National Academy of Sciences. 1959;**45**(2): 239-244
- [30] London J. Cytochrome in *Thiobacillus thiooxidans*. Science. 1963;**140**(3565):409-410
- [31] London J, Rittenberg S. Path of sulfur in sulfide and thiosulfate oxidation by thiobacilli. Proceedings of the National Academy of Sciences. 1964;**52**(5):1183-1190
- [32] Suzuki I, Chan C, Takeuchi T. Oxidation of elemental sulfur to sulfite by *Thiobacillus thiooxidans* cells. Applied and Environmental Microbiology. 1992;**58**(11):3767-3769

[33] Hallberg KB, Dopson M, Lindström EB. Reduced sulfur compound oxidation by *Thiobacillus caldus*. Journal of Bacteriology. 1996;**178**(1):6-11

[34] Quatrini R et al. Extending the models for iron and sulfur oxidation in the extreme acidophile *Acidithiobacillus ferrooxidans*. BMC Genomics. 2009; **10**(1):394

[35] Chen L et al. *Acidithiobacillus caldus* sulfur oxidation model based on transcriptome analysis between the wild type and sulfur oxygenase reductase defective mutant. PLoS One. 2012;**7**(9): e39470

[36] Yin H et al. Whole-genome sequencing reveals novel insights into sulfur oxidation in the extremophile *Acidithiobacillus thiooxidans*. BMC Microbiology. 2014;**14**(1):179

[37] Li L-F et al. The σ 54-dependent two-component system regulating sulfur oxidization (Sox) system in *Acidithiobacillus caldus* and some chemolithotrophic bacteria. Applied Microbiology and Biotechnology. 2017;**101**(5):2079-2092

[38] Wang Z-B et al. The two-component system RsrS-RsrR regulates the tetrathionate intermediate pathway for thiosulfate oxidation in *Acidithiobacillus caldus*. Frontiers in Microbiology. 2016;**7**:1755

[39] Wu W et al. Discovery of a new subgroup of sulfur dioxygenases and characterization of sulfur dioxygenases in the sulfur metabolic network of *Acidithiobacillus caldus*. PLoS One. 2017;**12**(9):e0183668

[40] Yu Y et al. Construction and characterization of tetH overexpression and knockout strains of *Acidithiobacillus ferrooxidans*. Journal of Bacteriology. 2014;**196**(12):2255-2264