We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Epidemiology of Bovine Mastitis and Its Diagnosis, Prevention, and Control

S.D. Audarya, D. Chhabra, R. Sharda, R. Gangil, R. Sikroodia, J. Jogi and N. Shrivastava

Abstract

Mastitis is an inflammation of mammary glands that is prevalent in dairy bovines. It causes a significant proportion of economic losses to the dairy farmers in India. Cattle and buffalo farming contribute significantly to the economy of the state. Various infectious agents such as bacteria, fungi, and algae may cause mastitis. Hence, it is essential to understand the etiological agents and predisposing factors that lead to mastitis in susceptible bovine populations in Madhya Pradesh state so that appropriate prevention and control strategies can be implemented. In this chapter, epidemiology, diagnosis, prevention, and control measures of mastitis in general and in India, the state of Madhya Pradesh, in particular, will be presented.

Keywords: mastitis, cattle, buffalo, bovine, Madhya Pradesh, epidemiology, diagnosis, prevention, control

1. Introduction

Mastitis, an inflammation of the mammary gland, is a very common disease in bovines. Among all the pathogens, bacteria are commonly implicated as the cause of mastitis [1]. Mastitis is characterized by inflammatory changes in milk and udder tissue. Mastitis is prevalent worldwide in dairy farming, causing heavy economic losses to the dairy industry. In those countries having well-developed dairy industry, morbidity of mastitis in dairy cows is 40% [2]. Affected bovines may lose their total milk production. Infectious agents like bacteria residing on the udder tissue or from the environment can enter through the teat canal. Milk from bovines with mastitis is unfit for human consumption because some mastitis-causing bacteria are zoonotic and can cause human infection [3]. Once the susceptible dairy bovine develops mastitis, it loses its milk production capacity significantly. Milk from a cow with mastitis is discarded due to inferior milk quality. Dairy animal owners have to bear extra costs for treating and maintaining such infected animals [4]. Clinically, there are two forms of mastitis: clinical form and subclinical form. The clinical form is characterized by local visible inflammatory changes in milk and udder tissue with or without systemic clinical signs, whereas subclinical form does not manifest clinical signs of mastitis but increased somatic cell counts with the presence of the causative agent. The diagnosis of the subclinical form of mastitis requires cow side test such as California mastitis test (CMT) or various laboratory tests including somatic cell

count (SCC) and milk bacteriological culture. Early diagnosis of subclinical form of mastitis is very much essential for successful treatment and control of infection [5]. Generally, high milk-producing cows are more suffered from mastitis than low milk producers [6]. This chapter highlights the epidemiology of mastitis and available diagnostic methods, prevention, and control measures with major focus on India in general and the state of Madhya Pradesh in particular.

1.1 The dairy sector and bovine population in the state of Madhya Pradesh, India

Madhya Pradesh is the second largest state in India with an area of 3,08,000 sq. km. The state is the part of the peninsular plateau of the country situated in the north-central part. It is one of the landlocked states in India (Figure 1).

It has three major seasons; summer, monsoon, and winter. The temperature of the entire state during summer (March–June) ranges above 29.4°C. Monsoon starts mid-June, and between June and September, the state gets the majority of its share of rainfall. In the winter season (November–February), the temperature remains low in the northern parts of the state compared to the southern parts. The state has over 70 million (7 crores) human population. The majority of the population of Madhya Pradesh (75%) resides in the villages, and most of them have income from agriculture. Tribal population accounts for 20% of the total population of Madhya Pradesh [7]. Livestock rearing provides them extra income and food security. Madhya Pradesh state Livestock and Poultry development corporation envision to increase the income levels of farmers involved in animal husbandry, particularly women by adopting a series of measures including a) increase in milk production, b) protect farmers from economic losses, and c) educate farmers about better management practices [8]. Madhya Pradesh state cooperative dairy federation limited has set up three-tier structures for dairy cooperatives. Primary village cooperatives (I tier), regional milk unions (II tier), and the apex federation (III tier) work in tandem for smooth functioning at field and plant operation levels and also for marketing the

Figure 1.
Madhya Pradesh state in India.

Animal	Livestock population (in million)				Change in percent
	India (2019)		Madhya Pradesh		
	Total	Female	2012	2019	
Cattle	192.49	145.12	19.6	18.7	−4.42
Buffalo	109.85	100.57	8.2	10.3	25.88

Table 1.
Bovine population in India and Madhya Pradesh.

branded products (Sanchi and alike). During 2018–2019, there were 517 milk routes and 4698 functional dairy cooperative societies. A total of U.S. \$ 148.71 million (Indian ₹ 1096.92 crores) has been paid to milk producers [9]. The total bovine population in 2019 increased by 1% to that of the total bovine population recorded in the previous census. According to 20th livestock census (provisional) statistics, the total bovine population in India stands at 302.79 (Cattle, Buffalo, Mithun, and Yak) million in 2019 [10]. However, in the state of Madhya Pradesh, only cattle and buffalo are reared. The cattle and buffalo population of the country and the state of Madhya Pradesh from the above-mentioned census is presented in **Table 1**.

Mastitis is an economically important infectious disease of cattle and buffalo in the state. It is essential to know its epidemiology, diagnosis, prevention, and control measures in India in general and in the state of Madhya Pradesh in particular.

1.2 Institutes working on mastitis in the state of Madhya Pradesh

Madhya Pradesh state at present is having three constituent veterinary colleges covering the western (Dr. Ambedkar Nagar-Mhow) and north-central and north-eastern region (Jabalpur and Rewa) of the state (**Figure 2**) under the auspices of Nanaji Deshmukh Veterinary Science University, Jabalpur. The laboratories there are well equipped for the diagnosis of cases of mastitis. Besides, the state also boasts of a vibrant veterinary service where disease diagnostic laboratories are the mainstay to avert any serious issues about infectious diseases.

Figure 2.
Locations of institutes working for diagnosis and prevention and control of mastitis in the state of Madhya Pradesh (red dot—Dr. Ambedkar Nagar-Mhow, pink dot—Jabalpur, green dot—Rewa).

Work published on mastitis from the western and north-central regions from the state of Madhya Pradesh is available. However, there are very few published reports presently available from the north-eastern region of the state. Although recently a study was conducted for the assessment of subclinical mastitis in bovines in and around Rewa district [11], in this study, a 31.4% prevalence rate for subclinical mastitis in cattle was reported. It is essential to improve awareness of agriculture and livestock owners in general and dairy animal owners in particular about prevalent bovine diseases including mastitis in the state of Madhya Pradesh to minimize production losses. So, information and toolkits regarding mastitis, its diagnosis, prevention, and control measures are being distributed.

2. Economic impact due to mastitis

Among the two forms of mastitis, the clinical mastitis is readily diagnosed by clinical manifestations (**Figure 3**) and affected animals can be treated but the subclinical mastitis is asymptomatic and often diagnosed late, after days or weeks of infection. In Asia, incidence rates of clinical and subclinical mastitis ranged from 1-8% and 55-60%, respectively. The dairy industry in India in general and in the state of Madhya Pradesh, in particular, contributes to the economy by uplifting people from poverty and also helps in earning a regular income. Mastitis is responsible for nearly 70% of milk loss. Economic losses due to mastitis in India were estimated to be around U.S. \$ 971.39 million (Indian ₹ 7165 crore) [12].

Figure 3.
Cow affected with clinical form of mastitis showing visible udder swelling.

3. Epidemiology of mastitis in India and Madhya Pradesh

3.1 Clinical and subclinical mastitis

Clinical mastitis is diagnosed readily by visible clinical signs and changes in the milk. In one of the studies conducted in India, in buffaloes from rural and urban dairy farmers, there was 18.74% prevalence of clinical mastitis [13]. In one of the other studies, 4.77% incidence of clinical mastitis was reported in the state of Madhya Pradesh [14]. Out of 260 cases of bovines in the Jammu region of the

country, 30 cases (11.54%) were positive for clinical mastitis. Prevalence of clinical mastitis in bovines ranged from 4.77 to 18.74% [13–15].

A higher level of incidence of subclinical mastitis (47.79%) was reported in the state of Madhya Pradesh [14]. In subclinical mastitis, there are no visible changes in milk or udder appearances in the affected bovines. Incidence of subclinical mastitis ranged between 19.2 and 83% in cows of Punjab state in India [16]. Recently, 40% of the overall incidence of subclinical mastitis was recorded in buffaloes in the state of Madhya Pradesh [17]. However, in lactating cows, the overall occurrence of subclinical mastitis was 27.81% [18]. In a recent study carried out in 2020, the reported overall prevalence of subclinical mastitis in cows was 31.55%. The prevalence of subclinical mastitis was highest in 5–7 years of age group (38.50%). The prevalence percentages of subclinical mastitis in organized (scientifically reared animals with adequate floor space availability) and unorganized dairy farms (animals reared in open space by livestock owners) were 29.82 and 41.66%, respectively [19]. In one of the other studies conducted in India, in buffaloes, there was 32.9% prevalence of subclinical mastitis [13].

When compared to clinical mastitis, subclinical mastitis is 15 to 40 times more prevalent [13, 20]. The prevalence of subclinical mastitis in dairy herds varied from 5 to 75% [21].

Subclinical mastitis results in greater economic losses to the farmers rearing dairy cattle and buffalo. No visible clinical signs are noticed in subclinically affected animals. In India, about 70–80% of economic losses have been attributed to subclinical mastitis alone. It occurs worldwide and also has adverse impacts on animal health and the quality of milk produced. Besides culling chronically infected animals from herds, decreased fecundity in affected animals and cost of treatment for mastitis lead to major economic losses. Subclinically infected cattle and buffalo can be a source of infection to other susceptible populations in the herd [22].

The prevalence of subclinical mastitis in bovines in India ranged from 9.88 to 86.87% [23]. In Madhya Pradesh, the prevalence of subclinical mastitis was reported highest in Jersey Cross (86.87%) than Holstein Friesian (75%), Malvi (57.35%), Sahiwal (75%), and Gir (80%) [24]. Among the various bacterial causative agents, staphylococci and streptococci were reported to be the most common pathogens of mastitis in India [25].

3.2 Etiology of mastitis

Mastitis is caused by physical, chemical, and biological agents. Generally, bacterial infections are the main causes of mastitis. Among many of the different microorganisms isolated from cases of bovine mastitis, the most common are Staphylococci (*Staphylococcus aureus*), Streptococci (*Streptococcus agalactiae*, *Streptococcus dysgalactiae*, *Streptococcus uberis*), and members of the family Enterobacteriaceae (*Escherichia coli*, *Klebsiella pneumoniae*) and other less common causative agents are *Pseudomonas aeruginosa*, *Mycoplasma* species, *Mycobacterium* species, *Nocardia asteroides*, *Candida* species, *Cryptococcus* species, and *Aspergillus* species. Rarely viruses are implicated in producing mastitis in bovines. The mastitis may be acute, per acute, subacute, chronic, and subclinical. Entry of the pathogen via teat canal into the mammary gland is characterized by increased leucocyte count in the milk [26].

Staphylococci, streptococci, *Escherichia coli*, *Pseudomonas* species, *Corynebacterium* species, *Mycoplasma* species, *Streptococcus agalactiae*, *Streptococcus dysgalactiae*, and *Mycobacterium tuberculosis* are isolated from buffalo suffering from mastitis. Among all the pathogens of bovine mastitis, staphylococci, streptococci, micrococci, *Corynebacterium* species, and *Escherichia coli* were isolated in Madhya

Pradesh. *Listeria* organisms were isolated from raw milk. Incidence of streptococci (32.53%), micrococci (5.74%), *Corynebacterium* (1.91%), and *Escherichia coli* (0.95%) was reported in cases of bovine mastitis. However, recently, 17.19% mastitis cases in buffalo were due to infection with *Escherichia coli* [27–31].

Depending on the mode of transmission of mastitis pathogens from their natural habitat to the mammary glands, there are contagious and environmental mastitis pathogens. Contagious mastitis pathogens exist on the udder or teat surface of infected cows. These are the primary source of infection from where they are transmitted during milking to uninfected cows. Major contagious bacterial mastitis pathogens are coagulase-positive *Staphylococcus aureus*, *Streptococcus agalactiae*, and *Mycoplasma bovis*. *Corynebacterium bovis* is the less common contagious mastitis pathogen [26]. *Staphylococcus aureus*, a bacterial species, is most commonly isolated from cases of bovine mastitis [32]. The reported prevalence of *Staphylococcus aureus* was 58.85% in Madhya Pradesh. *Staphylococcus aureus* is known to acquire antimicrobial resistance very quickly. Methicillin resistance *Staphylococcus aureus* (MRSA) is also a causative agent of mastitis in dairy cattle. Detection of MRSA has serious public health significance. In dairy cattle, the reported prevalence of MRSA mastitis was 16.47% in the Jabalpur region of Madhya Pradesh [33].

Environmental mastitis pathogens are in the environment of dairy cows and they can transfer to the mammary glands at any time. Major environmental mastitis pathogens include environmental streptococci (*Streptococcus uberis* and *Streptococcus dysgalactiae*) and coliform bacteria (*Escherichia coli*, *Klebsiella* species, *Enterobacter* species, and *Citrobacter* species). Some of them are opportunistic pathogens and infect mainly the immunocompromised host [26, 34]. Coliform bacteria (*Escherichia coli*) do not normally live on the skin or in the udder. These organisms can enter the teat canal when the animal comes in contact with a contaminated environment. Contaminated bedding materials, soil, manure, and organic matter in the environment can be a source of *Escherichia coli* that can lead to environmental mastitis. This kind of environmental mastitis was reported from several countries. *Escherichia coli* are one of the most frequently isolated bacteria from clinical infections. In severe, naturally occurring clinical cases of mastitis due to *Escherichia coli* there can be necrosis of the mammary epithelium [35].

4. Mastitis risk factors

Mastitis risk factors include managemental factors and cow factors. In management, the most common measures that can be used to avoid mastitis at the farm level are regular floor cleaning, use of appropriate milking techniques, and udder washing before milking and pre- and post-milking teat dipping in anti-septic solutions. Culling is hardly practiced in the country and also in the state of Madhya Pradesh. Regular screening of bovines for mastitis is not practiced at the larger scale as it needs to be in the state. Generally, dairy owners approach to the testing facilities after the cases of mastitis in bovines are clinically visible. Cow factors include the stage of lactation, breed, history of mastitis, and parity. Even if udder defense mechanisms are there, microbial infection overpowers at times and causes mastitis. Additionally, at times due to inadequate livestock management and husbandry practices such as unhygienic maintenance of livestock, inadequate floor space available to the animals, improper ventilation, and faulty milking techniques used by milkman also contribute as predisposing factors for mastitis. Physical injury to teat skin, teat canal, and mammary cistern are also important predisposing factors for entry of microbial pathogens in the udder to cause mastitis [36].

5. Diagnosis

Rapid cow side tests that are used to diagnose mastitis are highly required to implement prevention and control strategies. Some of the tests used in the diagnosis of clinical and subclinical mastitis are described below.

5.1 Cow side test—strip cup test

The physical appearance of milk is checked by a test named strip cup test. Strip cup or strip plate-based visual examination is routinely used for detection of clinical mastitis in individual and herd animals. The quality of the foremilk is examined visually after squirting few stripes of milk on the strip cup for gross examination of blood, flakes, clots, wateriness.

5.2 California mastitis test (CMT)

California mastitis test (CMT) is a simple rapid screening test, based on the estimation of the number of somatic cells in the milk sample. The somatic cell population consists of 75% leucocytes and 25% epithelial cells. The rise in the leucocytes indicates mastitis. The CMT reagent is mixed with the milk samples and the reagent causes lysis of somatic cells and release of DNA that form a gel. The CMT test result is qualitatively estimated (**Figure 4**). The average SCC of 2,00,000 cells/mL is considered as normal milk. For bulk tank milk $>2,00,000$ cells/mL of milk shows the presence of mastitis with a significant loss in milk production; for composite milk, from all the four quarters of a cow $>2,00,000$ cell/mL is considered mastitis whereas for milk from a single quarter of a cow $>1,00,000$ cells/mL is considered mastitis milk [37].

5.3 Milk bacteriological culturing and identification

For isolation and identification of bacteria-causing mastitis, the milk samples for bacteriological examination are first centrifuged and the resulting sediment is streaked on ordinary, selective, or differential media and incubated aerobically. Attempts should be made to isolate mycotic and anaerobic organisms. Milk sample

Figure 4.
California mastitis test paddle showing gel formation.

is streaked on sheep blood agar (0.05–1% aesculin), which supports the growth of most of the pathogenic bacteria-causing mastitis. The growth of bacteria can be further confirmed by primary and secondary biochemical tests and also by molecular detection methods.

After isolation of the bacterial pathogen, further identification is carried out using phenotypic and genotypic methods. Isolation and identification of the bacterial pathogen from cases of bovine mastitis are of paramount importance for testing effective antibacterial drugs against the bacterial isolate. Phenotypic characterization of bacterial pathogen includes: a) evaluation of bacterial morphology and growth characteristics, b) testing the ability of bacteria to metabolize substrates (biochemical tests), and c) testing of the antimicrobial sensitivity. Many of the commercial bacterial identification testing kits are developed using these phenotypic traits. These tests are easy to perform, readily available in the markets and cost effective. However, growing mastitis causing bacteria or fungi in the laboratory requires time and manpower. The major limitation for using tests based on phenotypic characterization is the variable expression of phenotypic characteristics by bacteria, even if the bacterial isolates are from the same species. This led to difficulty in identifying the bacteria correctly. However, culture systems at the farm site are increasingly being used for the diagnosis of mastitis [38].

Staphylococci—Colonies of *Staphylococcus aureus* (gram-positive cocci, generally appearing as irregularly arranged clusters) are round, shiny, golden yellow. These colonies are surrounded by a double zone of hemolysis on blood agar. Baired Parker agar and Mannitol salt agar are the selective growth media used to grow this species. Coagulase test (slide agglutination test and tube agglutination test) gives positive result in case there is a pathogenic strain of staphylococci. Streptococci (gram-positive cocci, generally arranged irregularly in chains) produce small translucent colonies on blood agar (with alpha, beta, and gamma hemolysis) [26, 37].

Streptococci-Edward's medium is a selective as well as an indicator medium for haemolysis and aesculin hydrolysis. Darkening of colonies shows hydrolysis of aesculin. *Streptococcus uberis* and *Enterococcus faecalis* hydrolyze aesculin. *Streptococcus agalactiae* and *Streptococcus dysgalactiae* are hydrolysis negative. Only *Enterococcus* grows on MacConkey agar and produces red pinpoint colonies. Christie-Atkins-Munch-Peterson (CAMP) test is used to identify hemolytic streptococci [26, 37].

Coliform bacteria—*Escherichia coli*, *Klebsiella pneumoniae*, and *Enterobacter aerogenes* are the most common species of bacteria isolated from milk samples collected from bovines suffering due to mastitis. MacConkey agar is used to grow coliform bacteria. *Escherichia coli* gives a metallic sheen appearance on Eosin Methylene Blue (EMB) agar. Colonies of *Escherichia coli* are generally non-mucoid. *Klebsiella pneumoniae* (non-motile) and *Enterobacter aerogenes* colonies are generally mucoid [26, 37].

5.4 Species and strain determination for a bacterium isolated from a case of mastitis

Nowadays, species and strain determination for a bacterium are carried out using a technique based on matrix-assisted laser desorption ionization–time of flight mass spectrometry (MALDI-TOF MS). It is a reliable, easy to use, and cost-effective technique. This technique can have 100% sensitivity and specificity for identifying the infectious agent from the cases of mastitis [39–42]. The major limitation of MALDI-TOF MS is, it can only have existing bacterial protein profiles for any specific interpretation. The technology is not available to common laboratories that have mastitis diagnosis facilities in the state of Madhya Pradesh. Genotypic methods use nucleic acid for the identification of species as well as strain typing [43]. Polymerase chain reaction

Molecular method	Species level	Strain level	Reference
AFLP	✓		[44]
RFLP		✓	[45]
MLVA		✓	[46]
Ribotyping	✓		[47]
tDNAiSLP	✓	✓	[48, 49]
PFGE		✓	[50]
DNA Sequencing	✓	✓	[51]
<i>AFLP, Amplified fragment length polymorphism; RFLP, Restriction fragment length polymorphism; MLVA, Multiple locus variable-number tandem repeat analysis; tDNAiSLP, Transfer DNA intergenic spacer length polymorphism analysis; PFGE, Pulsed field gel electrophoresis.</i>			

Table 2.
Molecular methods for identification of infectious agents of mastitis.

(PCR) is highly sensitive and the most commonly used molecular method to amplify target nucleic acid of a specific infectious agent causing mastitis. Virulent strains of organisms and those organisms that are not grown in culturing are also identified by using these molecular diagnostic methods (**Table 2**).

5.5 Electrical conductivity test

An increase in conductivity of milk from bovines suffering from mastitis is due to an increase in the salt concentration, which can be measured by an electrical conductivity test. Mastitis led to changes in ion concentrations [52], which impacts on the electrical conductivity of milk. Electrical conductivity can be measured [53] and the electrical conductivity rises with the increase in the concentration of sodium chloride in milk. Therefore, the measurement of electrical conductivity is used as a simple physical method to diagnose mastitis. An electrical conductivity meter is used to determine the electrical conductivity (EC). The EC of milk is expressed in the unit of milliSiemens (mS). This test has the following advantages: 1) One-time marginal investment is enough, 2) no special training is needed, and 3) easy to do and results are readily available.

5.6 Somatic cell count (SCC)

An increase in somatic cell count (SCC) in milk samples from bovines suffering from mastitis is measured by various tests such as CMT, white slide test, direct microscopic count, catalase test and anti-trypsin test, Brabant mastitis test (BMT), and Wisconsin mastitis test (WMT). Because of inflammation, the composition of milk in the animal suffering from mastitis is changed from normal to abnormal [54]. Somatic cell count is used to diagnose subclinical mastitis. The SCC includes direct microscopic somatic cell count (DMSCC), the bulk milk somatic cell count (BMSCC), and individual cow somatic cell count (ICSCC). The BMSCC is the universally accepted screening test for mastitis. In DMSCC, milk sample is smeared on a clean glass slide in the area of 1 cm² and stained with 1% methylene blue to examine 60 fields under the microscope for the count. The average number of cells in the fields is multiplied by the multiplication factor of the microscope to obtain the number of cells/ml of milk sample. Electronic somatic cell counters are used for DMSCC. A count of less than 1,00,000 signifies normal udder [1, 55, 56].

5.7 Chlorine test

An increase in chloride concentration of milk is an indicator of mastitis and it can be detected chemically. The presence of an increased quantity of chlorides in mastitic milk forms the basis of the test whereas normal milk contains about 0.07% chlorides. In mastitis, there is decreased amount of lactose and an increased amount of sodium chloride to maintain the normal milk osmotic pressure; hence during inflammation, there is an increase in the chloride content (> 0.14 percent) [57].

5.8 N-acetyl-B-D-glucosaminidase (NAGASE) test

This is a enzyme-based test that measure cell-associated enzyme N-acetyl-B-D-glucosaminidase in the milk. The highest level of the enzyme indicates a high cell count. It is a simple, effective, and most reliable test for the detection of subclinical mastitis. The readings for normal milk with 0.5×10^4 cells/ml and for mastitis milk with 1.5×10^4 cells/ml are 0.0053 and 0.034 moles/min/ml, respectively [58].

5.9 Indicators

Indicators of inflammation used to diagnose mastitis are well documented [59]. Recently, acute phase proteins were compared for diagnosis of subclinical mastitis in cross-bred cows in India [60].

5.10 Serological tests

Serological tests such as dot blot and enzyme-linked immunosorbent assays were used to detect antibodies to *Listeria* species antigens in the milk [61].

6. Mastitis prevention and control strategies

Five-point core plan for mastitis prevention and control include the following—a) disinfection of teats, b) adherence to hygienic practices in the milking procedures, c) removal of cows with chronic mastitis, d) dry cow therapy with antibiotics, and e) treatment of clinical mastitis [62, 63]. Additionally, in cow herds the measures used in prevention and control of mastitis are as follows—a) reduce introduction of new infections, b) shorten the duration of existing infections, c) maintenance of the normal udder health [64], d) dry and clean storehouse for fodder, feeders, and mangers, e) supply of palatable water, f) disinfection of milking area, g) shoe dipping disinfecting solutions for visitors to check any entry of the pathogen, h) check spread of infection by sprinkling lime powder [65], and i) keeping dairy animals stress free that encourage the development of healthy immune system [66].

10 point mastitis control program in cow herds recommended by National Mastitis Council is described in brief as follows: 1) establishment of goals for udder health and review those to prioritize changes in management to achieve set goals, 2) maintenance of a clean and comfortable environment by bedding management, keeping areas clean and dry, ensuring proper ventilation and provision of feed soon after milking so that animals remain in standing position, 3) adopting proper milking procedures by washing and drying teats, cleaning of udder with single-use clothes, examining foremilk and palpating glands to facilitate early detection of clinical cases, maintaining clean hands or wear gloves during the process of milking, 4) proper maintenance and use of milking equipment by routinely servicing it

and thoroughly washing and sanitizing, 5) good record keeping of incidence and prevalence of clinical and subclinical mastitis in cow herd, individual examinations, and treatments, 6) appropriate management of clinical mastitis during lactation by selecting appropriate therapeutic regimen and avoid treatment of cases suffering from resistant microbial and non-responsive agents, 7) effective dry cow management by drying cows off abruptly and careful administration of all quarters of all cows with suitable antibiotic, 8) maintenance of biosecurity for contagious pathogens by assessing test reports of cows (SCC, CMT) and also obtaining aseptically collected milk cultures from suspect cows before purchasing are must. Newly purchased cows must be kept and milked separately to ensure the absence of intramammary infection. Cows with a persistently high SCC (greater than 3,00,000) are segregated and observed for response to the treatment. Marketing of chronically infected cows with *Staphylococcus aureus* and antimicrobial-resistant microbial agents (*Mycoplasma*, *Nocardia*, *Pseudomonas*, *Arcanobacterium pyogenes*) that are unresponsive to treatment is carried out, and 9) regular monitoring of udder health status by enrolling cows for SCC program and monitoring rates and distributions of cows with high SCC. These cows with high SCC and cows with clinical infection are used for cultural examination. Inflammation is monitored by cow-side CMT. Reports from the regulatory agencies and marketing organizations are used for monitoring udder health for the herd, and 10) periodic review of the mastitis control program is evaluated by representatives from veterinary, industry and extension fields.

Bedding materials are the primary source of environmental mastitis. Hence, reducing the bacterial count in bedding generally decreases the risk of environmental mastitis. Teat cleaning (which includes wet cleaning followed by manual drying)/pre-milking teat dipping in antiseptic solution is important to reduce bacterial counts on the skin. In cows kept indoors, it reduces the incidence of new intramammary infection [67]. Teat dipping after milking dairy animals results in a reduction in the rate of new infection significantly. It is most effective against contagious mastitis pathogens such as *Staphylococcus aureus* and *Streptococcus agalactiae*. During the dry period, administration of, broad-spectrum antibiotics in each quarter of the udder at the last milking of lactation reduces the incidence of new intramammary infections. Dry cow therapy is the best way to cure chronic and subclinical mastitis. Effective and timely treatment of active cases is helpful to prevent new cases. Antibacterial therapy through the intramammary route is not always successful for treatment against mastitis. Particular antibiotics must be selected based on nature of the pathogen, results of antibiogram assay, and drug characteristics [68–70].

Farmer's awareness about a) what causes mastitis and b) how to prevent it, is essential to minimize economic losses not only in the state of Madhya Pradesh but also in India [71, 72]. Feeding micronutrients such as selenium and vitamin E are also very important to boost the immune response of the animals. Besides, recommending improvements in managerial practices, there must be proper attention made to provide better nutrition to animals for increasing immunity and reducing stress, and also encouraging farmers to participate in various awareness programs [73]. Knowledge of risk factors and characteristics of pathogens causing mastitis are essential to control the disease at farm level [74–77].

7. Summary

Mastitis affects the bovine population in India, it is also of major health concern to the bovine population of the state of Madhya Pradesh. Madhya Pradesh ranks fifth in milk production in India. Activities related to milk production and sale

fetch many livelihoods in the state. Bovine cases of clinical and subclinical mastitis caused by various pathogens are reported from the state and have economic implications for the dairy animal owners in terms of direct and indirect losses. There are various mastitis diagnostic tests routinely carried out by the institutes in the state for early and timely diagnosis of mastitis. It is aiding the authorities involved in livestock health for the better treatment of mastitis. Farmer awareness campaigns are undergoing but more impetus is needed to spread the word among them for better implementation of mastitis prevention and control strategies in bovines recommended by the National Mastitis Council.

Author details

S.D. Audarya^{1*}, D. Chhabra², R. Sharda², R. Gangil², R. Sikrodia², J. Jogi² and N. Shrivastava¹

¹ Department of Veterinary Microbiology, College of Veterinary Science and Animal Husbandry, Nanaji Deshmukh Veterinary Science University, Rewa, Madhya Pradesh, India

² Department of Veterinary Microbiology, College of Veterinary Science and Animal Husbandry, Nanaji Deshmukh Veterinary Science University, Dr. Ambedkar Nagar-Mhow, Madhya Pradesh, India

*Address all correspondence to: asd_vet@yahoo.com

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Radostits OM, Gay CC, Blood DC, Hinchcliff KW. *Veterinary Medicine: A Textbook of Diseases of Cattle, Sheep, Pigs, Goats and Horses*. 9th ed. London, United Kingdom: Book Power; 2003. 1877 p.
- [2] Padmanaban VD. Diseases caused by bacteria and fungi. In: *Handbook of Animal Husbandry*. New Delhi, India: Indian Council of Agricultural Research, Directorate of Information and Publications of Agriculture; 2002. p. 489-551.
- [3] Patil MP, Nagvekar AS, Ingole SD, Bharucha SV, Palve VT. Somatic cell count and alkaline phosphatase activity in milk for evaluation of mastitis in buffalo. *Veterinary World*. 2015;**8**:363-366.
- [4] Pankaj, Sharma A, Chhabra R, Sindhu N. Prevalence of sub clinical mastitis in cows: its etiology and antibiogram. *Indian Journal of Animal Research*. 2012;**46**(4):348-353.
- [5] Lakshmi R. A review on mastitis. *International Journal of Medicine Research*. 2016;**1**:118-123.
- [6] Singh B, Rao MLV, Gupta DK, Jain NK, Shukla PC. Epidemiological studies of sub-clinical mastitis in different organized dairy farms. *JNKVV Research Journal*. 2010;**44**(2):221-222.
- [7] Anonymous [Internet]. 2020. Available from: <https://mp.gov.in/state-profile> [Accessed: 04 December 2020]
- [8] Anonymous [Internet]. 2020. Available from: <http://www.mplivestock.com/> [Accessed: 04 December 2020]
- [9] Anonymous [Internet]. 2020. Available from: <http://www.mpcdf.nic.in/> [Accessed: 04 December 2020]
- [10] Anonymous. Key statistics provisional [Internet]. 2020. Available from: <http://164.100.117.97/WriteReadData/userfiles/key%20results.pdf> [Accessed: 03 December 2020]
- [11] Singh K, Mishra KK, Shrivastava N, Jha AK, Ranjan R. Prevalence of sub-clinical mastitis in dairy cow of Rewa district of Madhya Pradesh. *Journal of Animal Research*. 2021;**11**:89-95.
- [12] Anonymous [Internet]. 2020. Available from: <https://www.financialexpress.com/market/commodities/bovine-mastitis-resulting-in-70-per-cent-loss-in-milk-production-iit-roorkee-study/2062540/> [Accessed: 03 December 2020]
- [13] Sharma A, Sindhu N. Occurrence of clinical and subclinical mastitis in buffaloes in the state of Haryana (India). *Italian Journal of Animal Science*. 2007;**6**(Suppl. 2):965-967.
- [14] Ghose B. Studies on certain aspects of bacterial mastitis in cows [thesis]. Jabalpur: Jawaharlal Nehru Krishi Vishwavidyalaya (JNKVV); 2000.
- [15] Bhat AM, Soodan JS, Singh R, Dhobi IA, Hussain T, Dar MY, et al. Incidence of bovine clinical mastitis in Jammu region and antibiogram of isolated pathogens. *Veterinary World*. 2017;**10**:984-989.
- [16] Dua K. Incidence, etiology and estimated economic losses due to mastitis in Punjab and in India- An update. *Indian Dairyman*. 2001;**53**: 41-48.
- [17] Mohit K, Rao MLV, Shukla PC, Qadri MA. Incidence of latent mastitis in buffaloes of Jabalpur region of Madhya Pradesh. *International Journal of Livestock Research*. 2012;**2**:138-139.
- [18] Maheshwari P, Shukla PC, Rao MLV, Shukla SN. Occurrence of subclinical

- mastitis in cattle in and around Jabalpur, Madhya Pradesh. Haryana Veterinarian. 2016;**55**:160-162.
- [19] Mourya A, Shukla PC, Gupta DK, Sharma RK, Nayak A, Singh B, et al. Prevalence of subclinical mastitis in cows in and around Jabalpur, Madhya Pradesh. Journal of Entomology and Zoology Studies. 2020;**8**:40-44.
- [20] Kaur H, Chawla R. Importance of vitamin-E in animal health. Indian Dairyman. 2002;**54**:47-50.
- [21] Erskine RJ. MSD and the MSD Veterinary Manual. Merck Sharp and Dohme Corporation, a subsidiary of Merck and Co., Inc., Kenilworth, New Jersey, U.S.A. 2021. Available from: <https://www.Msdvetmanual.com/reproductive-system/mastitis-in-large-animals/mastitis-in-cattle> [Accessed: 14 October 2021]
- [22] Swami SV, Patil RA, Gadekar SD. Studies on prevalence of subclinical mastitis in dairy animals. Journal of Entomology and Zoology Studies. 2017;**5**:1297-1300.
- [23] Kumari T, Bhakat C, Choudhary RK. A review on sub clinical mastitis in dairy cattle. International Journal of Pure and Applied Biosciences. 2018;**6**:1291-1299.
- [24] Devi BK, Shukla PC, Bagherwal RK. Incidence of subclinical mastitis in cows. Indian Journal of Dairy Science. 1997;**50**:477-478.
- [25] Ramachandraiah K, Kumar KS, Sreemannarayana O. Survey of mastitis in a pure Jersey herd. Indian Veterinary Journal. 1990;**67**:105-106.
- [26] Quinn PJ, Markey BK, Leonard FC, FitzPatrick ES, Fanning S, Hartigan PJ. Veterinary Microbiology and Microbial Disease. 2nd ed. West Sussex, United Kingdom: Wiley-Blackwell; 2011. 912 p.
- [27] Ghose B, Sharda R, Chhabra D, Sharma V. Subclinical bacterial mastitis in cows of Malwa region of Madhya Pradesh. Indian Veterinary Journal. 2003;**80**:499-501.
- [28] Shrivastava N, Sharma V, Shrivastav A, Nayak A, Rai AK. Prevalence and characterization of panton-valentine leucocidin positive *Staphylococcus aureus* in bovine milk in Jabalpur district of Madhya Pradesh, India. Veterinary World. 2018;**11**: 316-320. DOI: 10.14202/vetworld.2018.316-320
- [29] Swarnkar R, Chhabra D, Barbudhe SB. Prevalence and molecular characterization of *Listeria* spp. in raw milk samples in Mhow and Indore region of Madhya Pradesh. The Indian Journal of Veterinary Sciences and Biotechnology. 2015;**10**:82-84.
- [30] Baghel A, Chhabra D, Sharda R, Shukla S, Audarya S, Sikrodia R, et al. Isolation of *Staphylococcus* from bovine mastitis and their antibiotic sensitivity pattern. The Indian Journal of Veterinary Sciences and Biotechnology. 2018;**13**:49-52.
- [31] Singh A, Chhabra D, Sikrodia R, Shukla S, Sharda R, Audarya S. Isolation of *Escherichia coli* from bovine mastitis and their antibiotic sensitivity pattern. International Journal of Current Microbiology and Applied Sciences. 2018;**7**:11-18.
- [32] Oliveira SS, Pova DC, Nascimento JS, Pereira MSV, Siqueira JP, Bastos MCF. Antimicrobial substances produced by *Staphylococcus aureus* strains isolated from cattle in Brazil. Letters in Applied Microbiology. 1998;**27**:229-234.
- [33] Shrivastava N, Sharma V, Nayak A, Shrivastav AB, Sarkhel BC, Shukla PC, et al. Prevalence and characterization of methicillin resistant *Staphylococcus aureus* (MRSA) mastitis in dairy cattle in

Jabalpur, Madhya Pradesh. Journal of Animal Research. 2017;7:77-84. DOI: 10.5958/2277-940X.2017.00011.0

[34] Schukken YH, Tikofsky L, Zadoks RN. Environmental control for mastitis prevention, milk quality and food safety. In: Hogeveen H, editor. Mastitis in Dairy Production Current Knowledge and Future Solutions. The Netherlands: Wageningen Academic Publishers; 2005. p. 109-114.

[35] Bradley AJ, Green MJ. Adaptation of *Escherichia coli* to the bovine mammary gland. Journal of Clinical Microbiology. 2001;39:1845-1849.

[36] Fagiolo A, Lai O. Mastitis in buffalo. Italian Journal of Animal Science. 2007;6:200-206.

[37] Anonymous. Mastitis. In: Laboratory Manual for Systematic Veterinary Bacteriology and Mycology. Mhow, India: College of Veterinary Science and Animal Husbandry; 2013. p. 20-23.

[38] Lago A, Godden SM, Bey R, Ruegg PL, Leslie K. The selective treatment of clinical mastitis based on on-farm culture results: I. Effects on antibiotic use, milk withholding time, and short-term clinical and bacteriological outcomes. Journal of Dairy Science. 2011;94:4441-4456.

[39] Barreiro JR, Ferriera CR, Sanvido GB, Kostrzewa M, Maier T, Wegemann B, et al. Short communication: identification of subclinical cow mastitis pathogens in milk by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. Journal of Dairy Science. 2010;93:5661-5667.

[40] Raemy A, Meylan M, Casati S, Gaia V, Berchtold B, Boss R, et al. Phenotypic and genotypic identification of streptococci and related bacteria isolated from bovine intramammary

infections. Acta Veterinaria Scandinavica. 2013;55:53.

[41] Mellmann A, Bimet F, Bizet C, Borovskaya AD, Drake RR, Eigner U, et al. High interlaboratory reproducibility of matrix-assisted laser desorption ionization-time of flight mass spectrometry-based species identification of nonfermenting bacteria. Journal of Clinical Microbiology. 2009;47:3732-3734.

[42] Bizzini A, Greub G. Matrix-assisted laser desorption ionization time-of-flight mass spectrometry, a revolution in clinical microbial identification. Clinical Microbiology and Infection. 2010;16:1614-1619.

[43] Zadoks RN, Watts JL. Species identification of coagulase-negative staphylococci: genotyping is superior to phenotyping. Veterinary Microbiology. 2009;134:20-28.

[44] Taponen S, Koort J, Bjorkroth J, Saloniemi H, Pyorala S. Bovine intramammary infections caused by coagulase-negative staphylococci may persist throughout lactation according to amplified fragment length polymorphism based analysis. Journal of Dairy Science. 2007;90:3301-3307.

[45] Saei HD, Ahmadi M, Mardani K, Batavani RA. Genotyping of *Staphylococcus aureus* isolated from bovine mastitis based on PCR-RFLP analysis of the *aroA* gene. Comparative Clinical Pathology. 2010;19:163-168.

[46] Pinho L, Thompson G, Rosenbusch R, Carvalheira J. Genotyping of *Mycoplasma bovis* isolates using multiple-locus variable-number tandem-repeat analysis. Journal of Microbiological Methods. 2012;88:377-385.

[47] Carretto E, Barbarini D, Couto I, De Vitis D, Marone P, Verhoef J, et al. Identification of coagulase-negative

staphylococci other than *Staphylococcus epidermidis* by automated ribotyping. *Clinical Microbiology and Infection*. 2005;**11**:177-184.

[48] Heir E, Sundheim G, Holck AL. Identification and characterization of quaternary ammonium compound resistant staphylococci from the food industry. *International Journal of Food Microbiology*. 1999;**48**:211-219.

[49] Stepanović S, Dakic I, Martel A, Vaneechoutte M, Morrison D, Shittu A, et al. A comparative evaluation of phenotypic and molecular methods in the identification of members of the *Staphylococcus sciuri* group. *Systematic Applied Microbiology*. 2005;**28**:353-357.

[50] Douglas VL, Fenwick SG, Pfeiffer DU, Williamson NB, Holmes CW. Genomic typing of *Streptococcus uberis* isolates from cases of mastitis, in New Zealand dairy cows, using pulsed-field gel electrophoresis. *Veterinary Microbiology*. 2000;**75**:27-41.

[51] Heikens E, Fleer A, Paauw A, Florijn A, Fluit AC. Comparison of genotypic and phenotypic methods for species-level identification of clinical isolates of coagulase-negative staphylococci. *Journal of Clinical Microbiology*. 2005;**43**:2286-2290.

[52] Kitchen BJ. Review of the progress of dairy science: bovine mastitis: milk compositional changes and related diagnostic tests. *Journal of Dairy Research*. 1981;**48**:167-188.

[53] Hovinen M, Aisla AM, Pyorala S. Accuracy and reliability of mastitis detection with electrical conductivity and milk colour measurement in automatic milking. *Acta Agriculturae Scandinavica Section-A Animal Science*. 2006;**56**:121-127.

[54] Korhonen H, Kaartinen L. Changes in the composition of milk induced by mastitis. In: Sandholm M, Honkanen

Buzalski T, Kaartinen L, Pyorala S, editors. *The Bovine Udder and Mastitis*. Helsinki, Finland: University of Helsinki, Faculty of Veterinary Medicine. 1995:76-82.

[55] Pilla R, Schwarz D, König S, Piccinini R. Microscopic differential cell counting to identify inflammatory reactions in dairy cow quarter milk samples. *Journal of Dairy Science*. 2012;**95**:4410-4420.

[56] Anonymous [Internet]. 2021. Available from: <https://www.nmconline.org/wp-content/uploads/2016/09/Guidelines-on-Normal-and-Abnormal-Raw-Milk.pdf> [Accessed: 06 July 2021]

[57] Davis JG. *Milk Testing (the Laboratory Control of Milk)*. Bikaner, India: Agro Botanical; 1999. 240 p.

[58] Mattila T, Saari S, Vartiala H, Sandholm M. Milk antitrypsin as a marker of bovine mastitis-correlation with bacteriology. *Journal of Dairy Science*. 1985;**68**:114-122.

[59] Pyorala S. Indicators of inflammation in the diagnosis of mastitis. *Veterinary Research*. 2003;**34**:565-578.

[60] Chouhan D, Aich R, Jain RK, Chhabra D. Acute phase protein as biomarker for diagnosis of subclinical mastitis in cross-bred cows. *Indian Journal of Animal Research*. 2020; **B-3943**:193-198. DOI: 10.18805/ijar.B-3943

[61] Baetz AL, Wesley IV. Detection of anti-listeriolysin O in dairy cattle experimentally infected with *Listeria monocytogenes*. *Journal of Veterinary Diagnostic Investigation*. 1995;**7**:82-86.

[62] Vlieghe SD, Ohnstad I, Piepers S. Management and prevention of mastitis: A multifactorial approach with a focus on milking, bedding and data-management. *Journal of Integrative Agriculture*. 2018;**17**(6):1214-1233.

- [63] Giesecke WH, Preez JHD, Petzer IM. Practical Mastitis Control in Dairy Herds. Durban: Butterworth Publishers South Africa; 1994. p. 317.
- [64] Philpot WN, Nickerson SC. Mastitis: Counter Attack. Illinois, USA: Westfalia Surge LLC; 1999. 150 p.
- [65] Pieterse R, Todorov SD. Bacteriocins – exploring alternatives to antibiotics in mastitis treatment. Brazilian Journal of Microbiology. 2010;**41**:542-562.
- [66] Kumar C, Kamboj ML, Chandra S, Kumar A. Dairy cattle welfare in India: A review. Asian Journal of Dairy and Food Research. 2017;**36**:85-92.
- [67] Gleeson D, Flynn J, Brien BO. Effect of pre-milking teat disinfection on new mastitis infection rates of dairy cows. Irish Veterinary Journal. 2018;**71**:11.
- [68] Chhabra D, Moghe MN, Tanwani SK, Sharda R. Studies on mycotic mastitis in cattle. Indian Veterinary Journal. 1999;**76**:1043-1045.
- [69] Sikrodia R, Chhabra D, Gangil R, Audarya SD, Sharda R, Mahor SS. Comparative *in-vitro* antibacterial efficacy of the different common herbs and antibiotics against *E. coli* and *S. aureus* isolated from bovine mastitis. International Journal of Current Microbiology and Applied Sciences. 2020;**9**:1069-1075.
- [70] Baghel A, Chhabra D, Sharda R, Audarya SD, Sikrodia R. Detection of biofilm forming ability of staphylococci causing bovine mastitis using phenotypic and genotypic assays. Journal of Immunology and Immunopathology. 2019;**21**(2):98-102. DOI: 10.5958/0973-9149.2019.00014.5
- [71] Kavitha KL, Rajesh K, Suresh K, Satheesh K, Sundar NS. Buffalo mastitis-risk factors. Buffalo Bulletin. 2009;**28**:135-137.
- [72] Mooventhana P, Kadian KS, Kumar RS, Kumaresan A, Manimaran A, Selvan AS. Effectiveness of mastitis management instructional video on knowledge of tribal farmers in northern hills zone of Chhattisgarh. Indian Journal of Animal Sciences. 2017;**87**:391-395.
- [73] Green MJ, Bradley AJ. Should you be using sand for dairy cow cubicles? In: Proceedings of the British Mastitis Conference, Garstang and Lancashire. Compton: Institute of Animal Health; 2001. p. 91-92.
- [74] Food and Agriculture Organization (FAO). Impact of mastitis in small scale dairy production systems. Animal Production and Health Working Paper No. 13. Rome, Italy: Food and Agriculture Organization of the United Nations; 2014. 34 p.
- [75] Singh M, Sharma A, Singh A. Isolation and antibiogram of beta haemolytic *Staphylococcus aureus* associated with bovine clinical mastitis. Haryana Veterinarian. 2013;**52**:54-56.
- [76] Johri A, Singh PK, Diwakar RP, Singh AP. *In vitro* herbal sensitivity test against mastitis milk of large animals. International Journal of Current Microbiology and Applied Sciences. 2018;**7**:81-89.
- [77] Tiwari A, Sisodia RS, Sharma RK, Misraulia KS, Garg UK. Incidence of sub-clinical mastitis in cows of Malwa region of Madhya Pradesh. Indian Journal of Dairy Science. 2000;**53**(4): 328-331.