We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Laparoscopic Findings of Rare Pediatric Inguinal Hernias

*Michinobu Ohno, Yasushi Fuchimoto, Akihiro Fujino,
Toshihiko Watanabe and Yutaka Kanamori*

Abstract

Pediatric inguinal hernias are caused to the patency of the processus vaginalis (PPV). The principle for the repair of indirect inguinal hernias in children consists of complete ligation of the PPV. Laparoscopic percutaneous extraperitoneal closure (LPEC) has spread rapidly since it was reported by some groups from around 1998, and the number of institutions adopting this method as a standard procedure for pediatric inguinal hernia is increasing in Japan. Since the closure of PPV by laparoscopic surgery is popular, rare hernias in children can be observed from the abdominal cavity. We present the laparoscopic findings of rare pediatric inguinal hernias and report their experience.

Keywords: indirect inguinal hernia, direct hernia, femoral hernia, child, adult

1. Introduction

Most inguinal hernias in children are classified as indirect inguinal hernias. Laparoscopic percutaneous extraperitoneal closure (LPEC) repair for pediatric indirect inguinal hernia is a standard technique in our facilities [1, 2]. We experienced two rare cases of pediatric inguinal hernias and reported their laparoscopic findings.

2. Case reports

2.1 Case 1

A one-year-old boy was admitted to our hospital for the incarceration of right inguinal hernia several times. He underwent LPEC procedure as a right indirect inguinal hernia. Laparoscopic findings demonstrated that the hernia orifice was present in the medial inguinal fossa (**Figure 1**). We converted the open approach and performed to reinforce the posterior wall of the inguinal canal. After the hernia sac was opened and excised, the transversalis fascia and Cooper's ligament were closed (McVay repair).

Figure 1.
Laparoscopic finding of right direct inguinal hernia. Hernia orifice presented medially to the external iliac vessels (arrowhead).

Figure 2.
Laparoscopic finding of femoral hernia at first operation. Hernia orifice presented medially to the epigastric vessels and below the inguinal ligament (arrowhead).

2.2 Case 2

The patient was a seven-year-old girl. She was diagnosed with the right external inguinal hernia and planned for LPEC procedure. Based on laparoscopic findings, we misdiagnosed an internal inguinal hernia (**Figure 2**). Iliopubic tract repair was performed with an open approach. One month after the operation, recurrence of inguinal protrusion was observed, and ultrasonography demonstrated femoral hernia (**Figure 3**). Therefore, the laparoscopic examination was scheduled. Intra-abdominal findings showed that the hernia orifice appeared to be closed, but traction with forceps confirmed the presence of a femoral hernia sac (**Figure 4**).

Figure 3.
 Ultrasonography of femoral hernia at recurrence. Herniation was recognized caudally to inguinal region and medially to the epigastric vessels (circle).

Figure 4.
 Laparoscopic finding of femoral hernia at second operation. A hernia sac with fat was identified (arrowhead).

We expected that the second open approach would have strong adhesion, therefore, we chose laparoscopic surgery. The hernia sac with the adipose tissue was resected. The iliopubic tract and Cooper's ligament were exposed and the femoral ring was closed (modified McVay repair).

3. Discussion

Direct inguinal hernia in children is quite rare. The incidence of direct inguinal hernias is as low as 0.2–1.2% of all pediatric inguinal hernias [3, 4]. Direct inguinal hernias seem to occur in males, and the affected side is on the right [4]. Fonkalsrud or other presumed the two or three etiologies of direct inguinal hernia: (1) attenuation of transversalis fascia, (2) increased abdominal pressure, and (3) weakness of the internal inguinal ring due to the previous surgery for indirect inguinal hernia. Wright divides the direct hernia into five based on the findings from the inguinal region [5, 6].

The correct diagnosis rate of indirect inguinal hernia is 38% preoperatively, and diagnosis is often missed even during surgery [5]. Recently, the laparoscopic diagnosis seems to be a feasible choice in this rare condition. In our case, preoperative diagnosis is an indirect inguinal hernia, however laparoscopic findings demonstrated direct hernia during surgery.

The treatment of direct hernia is different in each facility. Some authors recommend Bassini repair [4, 5], and some recommend McVay repair [6]. Laparoscopic closure of indirect hernia is described as an excellent technique for rare hernia. They use the vesical or umbilical ligament to close the direct defect [3, 7, 8]. However, laparoscopic treatment of pediatric inguinal indirect hernia is still controversial, and we performed McVay repair rather than laparoscopic closure to ensure posterior wall reinforcement.

Femoral hernia in children is less than 1% of all groin hernia [9–11]. Previous statements that femoral hernia is equally in males and females [12], and some authors indicated more frequently in boys than girls [13, 14]. A predominance of affected sides reported the right side, however, the cause is not understood [14, 15]. Regarding the etiology of femoral hernia, the most supported hypothesis was described by McVay and Savage [16]. They proposed that a congenital narrow insertion of the posterior inguinal wall onto Cooper's ligament caused an enlargement of the femoral ring.

A femoral hernia is also often misdiagnosed as a direct hernia. The correct diagnosis rate of femoral hernia is 53% preoperatively [17]. Several authors recommended a meticulous physical examination in the inguinal area, and ultrasonography is especially useful to distinguish femoral hernia from indirect inguinal hernia [14, 18]. We misunderstood our case as an indirect inguinal hernia in the first operation. Reconfirming the physical examination, the orifice of hernia was slightly caudal position and ultrasonography demonstrated femoral hernia. Pediatric hernias required a more accurate examination at the first visit.

The advantage of the laparoscopic technique in pediatric hernia includes the accurate diagnosis, minimal pain, and cosmesis. Although laparoscopic repair of femoral hernia was established in adults [19], most pediatric surgeons chose the open approach. In recent years, laparoscopic modified McVay technique reported in children [17, 20]. The laparoscopic approach was performed because of recurrent cases and the possibility of severe adhesion in our case. Laparoscopic closure required more operative time than an open approach. We consider the key to femoral repair is the closure of the hernia orifice without tension, following the resection of lipoma with the hernia sac.

4. Conclusions

Numerous pediatric surgeons have never seen rare hernias, however, in the laparoscopic era, intraabdominal findings revealed more higher rate of rare hernias. Rare pediatric inguinal hernias are challenging to diagnose and treat. In our both cases, the laparoscope approach was useful for diagnosis, however, simple or modified laparoscopic closure is still controversial.

Acknowledgements

I am grateful to Yuko Nakamura and Yuka Nomura for cooperating and supporting me as a medical team.

Conflict of interest

The authors declare no conflict of interest.

Notes/thanks/other declarations

Thanks to my colleagues.

Acronyms and abbreviations

PPV	patency of the processus vaginalis
LPEC	laparoscopic percutaneous extraperitoneal closure

Author details

Michinobu Ohno^{1*}, Yasushi Fuchimoto², Akihiro Fujino³, Toshihiko Watanabe⁴
and Yutaka Kanamori³

1 Department of Pediatric Surgery, Saitama City Hospital, Saitama, Japan

2 Department of Pediatric Surgery, International University of Health and Welfare,
Narita, Japan

3 Division of Surgery, National Center for Child Health and Development,
Tokyo, Japan

4 Department of Pediatric Surgery, Tokai University, Isehara, Japan

*Address all correspondence to: mohnomohno2000@hotmail.co.jp

IntechOpen

© 2022 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Endo M, Ukiyama E. Laparoscopic closure of patent processus vaginalis in girls with inguinal hernia using a specially devised suture needle. *Pediatric Endosurgery & Innovative Techniques*. 2001;**5**:187-191
- [2] Takehara H, Yakabe H, Kameoka K. Laparoscopic percutaneous extracorporeal closure for inguinal hernia in children: Clinical outcome of 972 repairs done in 3 pediatric surgical institutions. *Journal of Pediatric Surgery*. 2006;**41**:1999-2003
- [3] Schier F. Direct inguinal hernias in children: Laparoscopic aspects. *Pediatric Surgery International*. 2000;**16**(8):562-564
- [4] Wright JE. Direct inguinal hernia in infancy and childhood. *Pediatric Surgery International*. 1994;**9**:161-163
- [5] Fonkalsrud EW, Delorimier AA, Clatworthy HW Jr. Femoral and direct inguinal hernias in infants and children. *JAMA*. 1965;**192**:597-599
- [6] Viidik T, Marshall DG. Direct inguinal hernias in infancy and early childhood. *Journal of Pediatric Surgery*. 1980;**15**(5):646-647
- [7] Esposito C, Montinaro L, Alicchio F, Scermino S, Basile A, Armenise T, et al. Technical standardization of laparoscopic herniorrhaphy in pediatric patients. *World Journal of Surgery*. 2009;**33**(9):1846-1850
- [8] Lima M, Ruggeri G, Domini M, Lauro V, Libri M, Bertozzi M. Laparoscopic treatment of bilateral direct inguinal hernia by using the vesical ligament as an autologous patch. *Pediatric Endosurgery & Innovative Techniques*. 2002;**4**:277-280
- [9] Fosburg RG, Marlin HP. Femoral hernia in children. *American Journal of Surgery*. 1965;**109**:470-475
- [10] Marshall DG. Femoral hernia in children. *Journal of Pediatric Surgery*. 1983;**18**:160-162
- [11] Schier R, Klizaite J. Rare inguinal hernia forms in children. *Pediatric Surgery International*. 2004;**20**:748-752
- [12] Donnellan WL. Inguinal and femoral hernias. In: Donnellan WL, Burrington JD, Kimura K, editors. *Abdominal Surgery of Infancy and Childhood*. Luxemburg: Harwood; 1996. pp. 52/16-52/18
- [13] Temiz A, Akcora B, Temiz M, et al. A rare and frequently unrecognised pathology in children: Femoral hernia. *Hernia*. 2008;**12**:553-556
- [14] Aneiros Castro B, Cano Novillo I, García Vázquez A, López Díaz M, Benavent Gordo MI, Gómez Fraile A. Pediatric femoral hernia in the laparoscopic era. *Asian Journal of Endoscopic Surgery*. 2018;**11**(3):233-237
- [15] Wright MF, Scollay JM, McCabe AJ, et al. Paediatric femoral hernia: The diagnostic challenge. *International Journal of Surgery*. 2011;**9**:472-474
- [16] McVay CB, Savage LE. Etiology of femoral hernia. *Annals of Surgery*. 1961;**154**:25-32
- [17] De Caluwe D, Chertin B, Puri P. Childhood femoral hernia: A commonly misdiagnosed condition. *Pediatric Surgery International*. 2003;**19**:608-609
- [18] Erez I, Rathause V, Vacian I, et al. Preoperative ultrasound and

intraoperative findings of inguinal hernias in children: A prospective study of 642 children. *Journal of Pediatric Surgery*. 2002;**37**:865-868

[19] Peitsch WK. A modified laparoscopic hernioplasty (TAPP) is the standard procedure for inguinal and femoral hernias: A retrospective 17-year analysis with 1123 hernia repairs. *Surgical Endoscopy*. 2014;**28**:671-682

[20] Tainaka T, Uchida H, Ono Y, Tanano A, Shirota C, Yokota K, et al. A new modification of laparoscopic percutaneous extraperitoneal closure procedure for repairing pediatric femoral hernias involving a special needle and a wire loop. *Nagoya Journal of Medical Science*. 2015;**77**(3):531-535