We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Chapter

Immunoinformatics and Computer-Aided Drug Design as New Approaches against Emerging and Re-Emerging Infectious Diseases

Mohammed Othman Aljahdali,

Mohammad Habibur Rahman Molla and Foysal Ahammad

Abstract

Infectious diseases are initiated by small pathogenic living germs that are transferred from person to person by direct or indirect contact. Recently, different newly emerging and reemerging infectious viral diseases have become greater threats to human health and global stability. Investigators can anticipate epidemics through the advent of numerous mathematical tools that can predict specific pathogens and identify potential targets for vaccine and drug design and will help to fight against these challenges. Currently, computational approaches that include mathematical and essential tools have unfolded the way for a better understanding of newly originated emerging and re-emerging infectious disease, pathogenesis, diagnosis, and treatment option of specific diseases more easily, where immunoinformatics plays a crucial role in the discovery of novel peptides and vaccine candidates against the different viruses within a short time. Computational approaches include immunoinformatics, and computer-aided drug design (CADD)-based model trained biomolecules that offered reasonable and quick implementation approaches for the modern discovery of effective viral therapies. The essence of this review is to give insight into the multiple approaches not only for the detection of infectious diseases but also profound how people can pick appropriate models for the detection of viral therapeutics through computational approaches.

Keywords: immunoinformatic, pharmacophore modeling, molecular docking, quantum mechanism, peptide and vaccine design

1. Introduction

Infectious diseases are types of transmissible or communicable diseases mainly caused by pathogenic living microorganisms. The disease not only can transmit from animal to animal but also transfer from animal to human through the parasite, virus,

and bacteria [1]. The sudden invasion of infectious disease is a critical objective for the quietly alive of people in the ground [2]. However, it is not always an easy task to identify and anticipate the small pathogenic like small particles that are responsible for sudden chronic situations increasing incidence in geographic range [3]. Therefore, the situation demands to identify the unpredictable appearance of new infectious diseases as soon as possible that can utilize for further development of new therapeutic agents. Before identifying the causal infectious agents, the inevitable, but unpredictable, the appearance of newly emerging and re-emerging infectious diseases should be understood. The term emerging infectious diseases (EID) refers to infections that are newly arrived or evolved in a certain population, whose incidence can rapidly increase worldwide and threaten for future. On the other hand, infectious diseases that were previously appeared in the population and gradually decrease the incident but currently expanding into new geographical, host, or vector immensely that called re-emerging viral disease [4]. The virus disease has been recognized mostly from zoonotic by infecting a human. The primary source of zoonotic virus from farm and wild animal causes disease in human. Approximately 60 to 75% of known human pathogens ascend from animals [5]. Animal reservoirs have been involved in numerous virus families such as Filoviridae, Arenaviridae, Flaviviridae, and Bunyaviridae that were responsible to transmitted virus animals to humans *vice versa*. Most of the viruses abovementioned can be occurred emerging or re-emerging diseases, which do not have any specific therapeutic agents. Therefore, the development of vaccines and antimicrobial drugs candidates is an urgent issue that can control or prevent emerging or re-emerging diseases as soon as possible.

With the advancement of computational biology and immunoinformatics, rapid detection of pathogens and related proteins responsible for the disease has been developed. The technology helps to determine pathogen types within a short time and is utilized for therapeutic development. For example, immunoinformatic approaches can predict epitopes and their target protein is the recent advancement in vaccine design and development process that controlled the uses of antigen variation as well as hitting conserved epitopes [6]. It mainly designs immunogens by the protective responses of the target-specific receptor [7]. In the consequences of DNA virus infection and replication, it first attaches to the outer cell of the host through the protein receptor and replicates DNA by using host cell enzymes [8]. Finally, DNA goes to messenger RNA and translates it into a viral protein. Complete viral particles have converted by the replicate DNA and viral protein when new viruses were released from the host cell [9]. Additionally, RNA viruses are operated precisely as messenger RNA to make viral proteins. This mechanism of viral infections can be predicted through computational tools and immunoinformatics can identify desired epitopes for designing vaccines against the infections. On the other hand, computer-aided drug design (CADD) consists of a computer-assisted *de novo* design that can predict several models based on drug-target interaction network and help to early stage of drug design through molecular docking, similarity search methods, and deep learning-based model. The confirmation of ligand and the target protein can be initially predicted through molecular docking [10]. However, the performance of docking is quietly dependent on various types of receptors as well as acting best for the hydrophobic vs. hydrophilic pockets, which can also be determined by the CADD approaches.

Furthermore, consistent advancement of medical and pharmaceutical research has been playing an important role in the proper solution of several diseases but remained some problems with viral disease that may suffer or burden to public and

animal health [11]. Therefore, computational techniques have opened a new avenue for minimizing the problems of drug discovery. The pharmaceutical industry has been starting to take up help from computational methods for drug design and development, drug repurposing, enlightening medicinal efficiency, and clinical trial [12]. The increasing data digitalization in a pharmaceutical company can be solved clinical problems that can also be done through computational methods. They have maintained a large volume of data for solving infectious diseases by their automation. The extremely widespread diseases remain virus diseases that cause infection by a type of microorganisms [13]. The most prominent type of virus diseases is common cold, while another one is an infection of virus that affected the upper respiratory tract such as nose and throat [14]. Furthermore, antibiotics have a magical role in preventing bacterial disease and infections but have no effects on viral disease. The most significant challenge to the researcher is finding out the epidemiology, vaccine design, and eradication in a worthwhile manner [15].

2. Infectious disease

Infectious disease is a kind of disorder that is caused by one or more organisms. Many organisms consist of our body where some are normally helpful or infectious with parasites and viruses as well as bacteria and fungi [16]. SARS, recently emerged SARS-CoV-2, tuberculosis (TB), HIV/AIDS, influenza, Chickenpox, the common cold, and Hepatitis A and B are some examples of infectious diseases that can easily transmit from human to human or animal to human under adverse conditions and help to cause diseases. However, the animals or insects are directly or indirectly responsible for transmission to these organisms. Contaminated food items have exposed some infectious to the new environment and move apart from diseases [17]. There are various signs and symptoms of infectious diseases such as fever, fatigue, coughing, muscle aches, and diarrhea, and so on. Infectious diseases can be remedied or controlled by vaccines, but it takes a long time for application. Most infectious diseases controlled by consciousness of diseases such as handwashing can help to protect from many infectious diseases [18].

2.1 Global burden of infectious diseases

Over the last few centuries, millions of people have lost their lives to infectious diseases throughout the world. The risk of public health has been reduced and controlled through the improvement of the sanitation system, the progress of antibiotics and vaccines, living condition, and food quality, which is linked to the socioeconomic modernization of these societies [16]. Despite the improvement of health care facilities and surveillance systems, there have been some uncertainties that still now lead to an increase the human mortality from infectious diseases. The current threat to global human health is zoonoses [19]. The mutually transmitted 200 diseases occurred between humans and animals. The zoonotic disease has been increasing due to the overpopulation, wars, and food scarcity that are associated with humans who contract face to face [20]. However, the global death increased in 2010 due to infectious disease from HIV/AIDS that constitutes 1.5 million, and hereby malaria raised to 1.17 million [21]. At the same time, about 152,000 people died by the neglected tropical disease, and 1.2 million people died of tuberculosis [22]. Long-term illness, disability, and social stigma are occurring from poverty

and are renamed by infectious diseases of poverty (IDoP). Moreover, coronavirus disease 2019 (COVID-19) has the main worldwide burden in this era expressing itself as a pandemic and killed almost 4,825,433 people from the beginning and has been infecting till now [23]. The global health system has been affected by the pandemic due to the unintentional interruption of service delivery. Until now, a total of 369 infectious diseases have been identified based-on mortality and the estimation of life expectancy according to global burden diseases in 2020. There are several factors responsible for diseases that have compiled during the COVID-19 pandemic by analyzing 286 causes of death, about 369 injuries and infections, and 87 threat issues from the 204 nations [24].

2.2 Emerging and re-emerging infectious disease

Several infectious pathogens have been halting their activities from the initial discovery and reappearance after a long and short period in several places [25]. Among them are Emerging and Re-Emerging Infectious Diseases (EIDs). They are types of infections that have newly originated or previously existed in a population that is rapidly increasing in incidence or geographic range. The newly emerging and re-emerging viral disease has threatened public and animal health, which is increasing due to human activities [26]. There are many associated factors related to the spread of infectious viral disease from one place to another place such as population migration, urbanization, public gathering, poverty, malnutrition, increased domestic and global connectivity and environmental changes, and so on [4]. The alteration of genetic phenomena has also been responsible for to spread of disease to a greater extent. Most of the infectious diseases have been estimated in zoonotic that constituted 60 to 70% of total diseases [27]. Hence, the animal virus also mutated with a human virus that accelerates infectious viral disease and generates the chronic problem. As a result, thousands of people die without medicine or even without knowing about the threat [28]. Novel pathogens have occurred due to unplanned urbanization of habitat destruction that enhances the contact or susceptibility to infections between human and animal vectors of viral diseases herewith the lack of immunity of these communities [29]. However, many pathogens re-emerging again after many years such as Chikungunya and Zika virus.

2.3 History of emerging and re-emerging virus

Initially, there was an outbreak of chikungunya between 1963 and 1973 with serious arboviral illness and it re-emerged in 2006, which firstly was observed in the East African strain of chikungunya [30]. Similarly, the Zika virus was initially associated with a serious illness that has conducted by serological studies in 1960 and after a long time, it has identified and reported from Brazil in 2015 [31]. The discovery of novel pathogens in the world has not been clogged and they are associated with economic cost reflect the burden for many developed and underdeveloped countries (Table 1). The high cost of medical and intensive care has been banned for all kinds of work from the affected region.

2.4 Factor influences emerging and re-emerging viral bond diseases

Viral diseases have chorionic effects on public health worldwide. The annual death has been attributed to about 20 million by infectious diseases [34]. Most of the death

Family	Pandemic Year	Viruses	Probable/mode of transmission	Outbreak potential
Bunyaviridae	1917	Ganjam virus or Nairobi sheep disease virus (NSDV)	Tick-borne	Yes*
	1954	Bhanja virus	Tick-borne	Yes*
	2009	SFTS virus	Tick-borne	Yes
	1719	Chobar Gorge virus	Tick-borne	No
	2008	EEV	Arthropod-borne	No
	2004	Cat Que. virus	Arthropod-borne	Yes*
	1957	Kaisodi virus	Tick-borne	Yes*
	1955	Umbre virus	Arthropod-borne	Yes*
	1965	Ingwavuma virus	Arthropod-borne	No
	1957	Chittoor virus	Tick-borne	Yes*
	1964	Thottapalayam virus	Rodent-borne	No
Nairoviridae	1944	CCHFs virus	Tick-borne, human to human	Yes
Flaviviridae	1955	Yellow fever	Arthropod-borne	Yes
	1947	Zika virus	Arthropod-borne, mother to child, sexual route	Yes
	1957	KFD	Tick-borne	Yes
	1952	JE	Arthropod-borne	Yes
	1992	Dengue	Arthropod-borne	Yes
	1966	Bagaza virus	Arthropod-borne	Yes*
Paramyxoviridae	1968	Influenza - (H3N2) v alias	Air-borne	Yes
	2004	Avian Influenza	Air-borne	Yes
	2006	Influenza -Avian (H5N1)	Air-borne	
	1956	RSV	Air-borne	Yes
	1953	Quaranfil virus	Tick-borne	Yes*
	late 1950s	Parainfluenza 1–4	Air-borne	Yes*
	2009	Influenza H1N1	Air-borne	
	1962	Enterovirus-D68	Air-borne	Yes
Paramyxoviridae	2001	Nipah virus	Human to human Direct contact/consumption of infected bat/fruit infected with bat	Yes
Picornaviridae	1953	Human rhinovirus A, B and C	Air-borne	Yes
	1957	Hand, foot and mouth disease	Direct contact, feco-oral route	Yes
	1948–49	Coxsackie-A21 virus	Feco-oral route	Yes
	1948–1949	Coxsackie-A10 virus	Feco-oral route	Yes
	1977	Sapoviruses	Feco-oral route	Yes
	1973	Rota	Feco-oral route	Yes
	1997	Polio and non-polio flaccid paralysis	Feco-oral route	Yes

Family	Pandemic Year	Viruses	Probable/mode of transmission	Outbreak potential
Caliciviridae	1936	Noroviruses	Feco-oral route	Yes
Hepadnaviridae	2007	Hepatitis KIs virus new and vaccine escape mutants of HBV	Blood-borne	Yes
Togaviridae	1740	Rubella virus	Air-borne	Yes
	2005	Chikungunya virus	Arthropod-borne	Yes
Poxviridae	1934	Buffalopox virus (orthopoxvirus)	Direct contact	Yes
	1958	Human monkey pox	Air-borne	
Parvoviridae	1975	Human parvovirus 4	Parenteral transmission?	Yes
Arenaviridae	1934	LCMV	Rodent-borne	Yes*
Herpesviridae	1934	CMV	Direct contact	Yes
	1974	Chickenpox (varicella) VZV	Air-borne, direct contact	Yes
Rhabdoviridae	1965	Chandipura virus	Arthropod-borne	Yes
Reoviridae	1963	Kammavanpettai virus (orbiviruses)	Tick-borne	No
Coronaviridae	2003	SARS Coronavirus	Air-borne	Yes
	2019	Coronavirus disease-2019 (COVID-19)	Air-borne	Yes
Polyomaviridae	1953	Polyoma-like virus	Human to human	Yes
Phenuiviridae	1931	Rift valley fever	Blood-borne	Yes

Table 1.
Emerging and re-emerging virus figure out with the inauguration periods [32, 33].

not only occurred by acute respiratory tract infection and gastrointestinal infections but also come out through tuberculosis and malaria that remained unchanged till now [35]. Many factors have been leading to emerging and re-emerging infectious diseases such as demographic factors, population distribution, sexual behavior, childcare, food-borne and water-borne diseases, ecological alteration and land use, chronic manifestations, enhanced pathogen detection, microbial evolution, and failure of community health scheme and bioterrorism shown in **Figure 1** [36]. The variations of the global population have been contributing to the growth and density in per capita, resettlement to the city area, international tourism, migration, and retaining density that directly or indirectly leads to the infectious virus disease [37]. However, there are some intentional reasons too responsible for the emerging and re-emerging virus such as the deliverance of sexual practice, enhanced childcare beyond the family, alcohol and drug abuse, food supply, transportation, and immunization practices [32]. Moreover, the environmental and land use causes of global warming, deforestation, and natural disasters such as floods, drought, and *El Nino* effect will be the lead to current and future infectious diseases [38]. The invention of modern technology can manifest the infectious diseases that are prolonging the life of people in the world. The use of new molecular techniques has enhanced to detection of fastidious and uncultivable organisms [39]. As a result, a variety of pathogens were discovered in a short time. Nowadays, microbiomes naturally adapt to their environment for survival causing a wide range of microorganisms that

Figure 1.
Representing different factors that influence the activities of emerging and re-emerging viral bond diseases.

have become resistant to diseases [40]. Consequently, it can alter the present situation and convey the hazardous situation for human life.

2.5 Preventive measures of emerging and re-emerging disease

Due to the degradation of the environment, many contagious infections spread out the world for a long time. However, some diseases have been treated through the antibiotic and controlled with the vaccine but most of the disease's remedies remained unchanged still now [41]. Scientists and researchers have been searching for a new link in an infectious chain through their internal activities. The consumption of raw food is one of the triggers of infectious diseases. Raw food usually has lots of harmful microorganisms that can hamper your normal life within a short time [41]. Several factors have been contributing to the vast range of emerging and non-emerging diseases such as changes in human behaviors, enhancement of technology, change in the land pattern and economic progress, tourism, microbial change, and collapse of public health measures [41]. However, the prevention and control have been taken by the Pan American Health Organization (PAHO) that regulates some measures of emerging and re-emerging diseases [42]. First, there is need to increase and strengthen regional surveillance networks that control infectious diseases from one place to another place. Later, the rapid responses of infectious diseases thereby enhance the laboratory equipment and training. Moreover, the development of applied research fields can prevent the disease through the rapid diagnosis of epidemiology as well as prevention [43]. Finally, the enhancement of regional capacity and strength can control and prevent emerging and re-emerging diseases through effective implementation [44].

2.6 Traditional methods of preventing emerging and re-emerging disease

Emerging viral diseases have occurred previously and resultant pandemics where plant derived was the first choice for treatment. Around 1500 BC, Egyptians were started herbal drug preparation by the medicinal plant where later Greek and Roman

have improved it [45]. Emerging infectious diseases have been treated traditionally by the medicinal plant before inventing a drug or vaccine [46, 47]. In China, it was widely used to deal with infectious diseases. They believe that it boosts the immune system and protects the virus to enter the respiratory tract [48]. Many medicinal herbs have a unique antiviral effect and are used as a substitute for an antimicrobial drug against infectious diseases. They have used the whole body of herb or part of the plant such as leaves, roots, bark, fruit, seeds, flowers, and so on, for preventing and healing respiratory tract infections [49]. In the modern time, it has been perceived that the traditional drug safer and healthier than synthetic drug those comes from plant-based traditional medicine. Plant-derived drug discovery has been renowned for the last decade [50]. It is considered that plant products will be an indispensable source of a new drug in the future.

3. Computer-aided drug design approaches

3.1 Pharmacophore modeling

Pharmacophore is one of the most promising *in silico* concepts, which is utilized to screen large compound libraries. The process includes combining medicinal chemistry and computational chemistry that can screen and optimize lead compounds for the development of the final drug candidate [50]. The pharmacophore model can function in two ways such as ligand-based modeling and structure-based modeling (**Figure 2**). The ligand-based model is fully based on computerized for simplifying drug discovery in the macromolecular target structure [51]. It generates three-dimensional (3D) structures for interacting ligand and macromolecules. Software such as Schrödinger (<https://www.schrodinger.com/>) and so on are used for pharmacophore drug design. Furthermore, the direct interaction of 3D structure with macromolecule ligand complex is arranged by structure-based pharmacophore modeling. The investigation of the complementary chemical feature of the active site and their spatial relationships and assembly with the selected feature will be generated using the structure-based pharmacophore modeling [52]. Discovery studio is usually practiced for the implementation of the structure-based pharmacophore method. The catalytic feature of pharmacophoric is H-bond acceptor, H-bond donor, and hydrophobic. Virtual-based pharmacophore screening has been reducing the possibility of arising problems about inadequate consideration of protein selectivity with the optimizing

Figure 2.

Pharmacophore model-based virtual screening process for identification of small molecular candidates against a specific disease.

score and insufficient design by leading a tolerance radius for each feature [53]. The main purpose of pharmacophore model-based virtual screening is to count the highest molecules hit, which has similar or nearest the target.

3.2 Virtual screening

3.2.1 Molecular docking

In modern drug design and discovery, molecular docking is one of the foremost strategies to find the best molecules for a specific target. It assists to predict the interaction of protein and a ligand at the bond stage. Molecular docking is used to calculate the scoring function by sampling algorithms shown in **Figure 3** [54]. Furthermore, it is reasonable, time demanding, and effective more than any conventional technology. The computer-based drug designing method predicts the function of any individual protein of interest that conducts a very comprehensive approach [55]. It has various applications among them protein-ligand docking is one of them. It is primarily designed to predict the binding of small drug molecules from medicinal herbs or plants, invertebrates, and so on to target proteins [56]. The number of diseases is caused by the harmful receptors; hence, docking is designed to inhibit or induce the target protein. However, the discovery of drugs in the traditional way is very expensive because of the need to search a large volume of compounds to select against a particular protein for the proper binding interaction and targeting diseases [57]. Therefore, computerized docking can screen virtually thousands of compounds in a short time by experimental high-throughput screening and confirmed the ligand and receptor for stable binding. It also provides the knowledge-based scoring functions that interact with atom pairs between protein and ligand complexes along with three-dimensional structure. There are several purposes of molecular drug design [58]. Among them, three are main purposes such as predicting the active ligand, binding affinities, and identifying new ligand. Based on molecular

Figure 3.
Molecular docking-based screening approaches of small molecules against a specific target.

docking, it has numerous applications such as documenting the lowest free energy structure for the receptor-ligand complex, estimating the differential binding of a ligand of two separate macromolecular receptors, the geometry of the specific ligand-receptor complex, lead generation and optimization for future drug candidate, and so on [59].

3.2.2 Molecular dynamic simulation

Molecular dynamics (MD) simulation is a computer-based simulation technique that mainly analyzed the physical motion of atoms and molecules across a specific conformational space [60]. MD simulation is the unique way to drug discovery in the present era that led to work in a wet lab after computer simulation. The MD simulation ensures the detail of the dynamic properties of selected protein and plays a key role in the modeling and characterization of a protein [61]. The MD simulation also helps to determine the stability of a ligand to the active site of the targeted macromolecules. Moreover, MD simulation not only provides the information of the ligand optimization process at the qualitative level but also estimates accurately in ligand binding affinities. The tiny, microscopic event (protein-ligand interaction and molecular motion) occurs with a micro- or nano-second time scale that is not possible to determine without the technique [62]. Finally, dynamic simulation is a faster, reasonable, widely accessible, and perfect method for drug design *in silico* techniques. Moreover, it provides information of atomic position with respect to time. It usually explains the atomic and molecular properties of the protein, drug-target interaction, solvation of compound, and conformational changes that a receptor expresses in various conditions [63]. It works on the base of Newtonian mechanics (NM), which is related to the motion of large particles. The force fields are an important factor for MD simulation, which are mainly a set of the potential energy function for employing the relation between structure and potential energy. The particles are coordinated by the system of mathematical expression [64]. The computerized technique (force and energy) is used for building blocks that combined bonded and non-bonded interaction. Force acting on each atom was counted by using the force field of molecular machines that were developed through the four principal such as Born-Oppenheimer method, bond length, and bond angle, and potential energy of the surface molecule and atom type [65].

3.2.3 Quantum mechanics

Quantum mechanics has been leading a valuable method for drug discovery through characterizing the structure, dynamics, and energies of protein-ligand interaction [66]. In the medicinal industry and academic research, it is an inevitable part of drug design that fixes the problem by calculating chemical reactivity and helps to optimize structure [67]. The new drug candidate has been chosen by molecular mechanics (MM) but the quantum mechanics (QM)-based approach provides accurate accuracy and efficiency in the complexity of protein-ligand interactions [68]. Furthermore, it does not provide only the estimation of being affinities, determining ligand energies and bioactive conformations, refinement of molecular geometries but also added scoring docked ligand poses, describing molecular similarity, structure-activity, relationship analysis, and ADMET prediction in the activities of drug design [69]. QM is getting popular day by day in drug discovery because of the improving power or speed of the computer that led to advanced QM algorithm progress as well as a new application to address the shortcoming [70]. The popular method of fragment molecular orbital (FMO) offers a wide range of solutions where there are combined

accuracy, speed, and the ability to characterize important interaction such as strength in kcal/mol as well as hydrophobic, electrostatic, and so on. In summary, FMO analyzed the length such as polarization, desolation, and interaction with the illustrated for a water dimer and protein-ligand complex [71].

3.2.4 Pharmacokinetic properties

Pharmacokinetics is the unique part of the drug design where the time course study is done through the mathematical characterization for absorption of drug, distribution, metabolism as well as excretion (ADME) [72]. Therefore, before the clinical trial, it is the advanced procedure to select a drug for the development and decision-maker by AMDE test [73]. As a result, it is safe and effective for a specific patient by decreasing toxicity and increasing efficacy in drug therapy. However, the central compartment of the human body was interconnected by a series of compartments reversibly on the base of pharmacokinetic [74]. Therefore, drugs enter each compartment and are distributed homogenously to display consistent kinetics. Hence, it is separated by a mathematical model where AMDE can differ in each compartment and help to predict drug metabolisms or actions [75]. Moreover, the central compartment received drug and entry on the body called absorption. The drug is distributed to the peripheral compartment after absorbing into the body and beginning the metabolic processes. Finally, the excretion of the drug from the body is in three ways such as hydrophilic molecules by the renal system, hydrophobic molecules by the biliary system, and volatile substances by the pulmonary system [76].

4. Immunoinformatic

4.1 Peptide design

In the drug discovery, peptides are not new, but it makes a choice drug candidate in the challenging situation with the collaboration of immunoinformatics. Immunoinformatics can be developed as natural endogenous scaffolds with familiar biological activities for solving challenging medical problems by it distinguishing characteristics such as the ability to act firstly, more specific, the minimum range of toxicity, and so on [77]. Approximately 55 therapeutic peptides have been approved for clinical trial through the regulatory agency [78]. Currently, many researchers have been working to solve various diseases through antimicrobial peptide design. The target of antimicrobial peptides has accelerated through the deep generative models and molecular dynamics. By utilizing the immunoinformatic approach, the researcher has developed 20 novel peptides that were validated through deep learning and high-throughput molecular simulation [79]. The peptides are not only used in broad-spectrum potency but also lead to multi-drug resistant strain and low toxicity. Immunoinformatic approaches help to observe peptide activity, selectivity, toxicity, ease of synthesis stability, etc. During the present situation of COVID-19, immunoinformatics help to identify peptide candidates from diverse viral sequences as shown in **Figure 4**, which can be utilized for vaccine design [80]. It is identified and designed from the motifs and subsequently a peptide library. As a result, strong binding affinities of peptides had happened with the main protease of SARS-CoC-2 as well as maintained stability and physiological condition observed by molecular dynamic simulation [81].

4.2 Vaccine design

Vaccine design is a very powerful tool for saving human life in the world broadly. The advancement of technology has been trigger to develop new strategies by the target and the structures of antibodies, hitting conserved epitopes and controlling the usage of antigenic variation [82]. It designed immunogens for provoking foreign responses because of no single or best solution of immune drugs design till now [83]. However, artificial intelligence and system biology together make an opportunity for avoiding inefficiencies and failures that help to classical vaccine development pipeline [84]. *In silico* vaccine design process has been selecting well fragment of virus protein [85], thereby finally leading to a final vaccine as shown in **Figure 4** [86].

However, in the vaccine design process, DeepVacPerd is an efficient tool that predicts the best vaccine subunit candidates with 30 subunit candidates from the various protein sequences within a second by replacing the predicted and selected with deep neural network architecture [87]. Therefore, it has become a promising of higher efficiencies for the vaccine design and test process. Furthermore, systems biology developed various tools by analyzing large data set through the complex modeling interaction between the individual interaction [88]. The omics disciplines such as genomics, proteomics, metabolisms, and so on produced a simulation of the immune response, identified response-specific signature, and assessed their predictive value through the basilar idea with the integration of high-throughput data [89]. Therefore, several programs and consortiums have performed by developing novel analytic tools that may integrate the information from omics.

4.3 Future perspective of vaccine

Personalized vaccine candidates are developed against a specific “targeted” to maintain an optimized outcome. Failure of the vaccine candidates can increase

Figure 4. Immunoinformatics aided identification of peptide candidates and procedure for designing multi-epitope vaccine by utilizing the peptide.

immunogenicity subsequently reactogenicity and adverse effects [90]. Therefore, the individual level, the gender level, the racial/ethnic level, and the subpopulation level should be considered during the personalized vaccine development process. Haplotype and polymorphism are mainly focused on the individual level selection as they can retard the formation of a protective immune response, where the gender level can determine the antibody titer against a particular vaccine to males and females [91]. On the other hand, different human races or ethnic groups can show higher or lower immune responses to a specific vaccine candidate. Additionally, different interactions between host environmental, genetic, and some other factors may be influencing the vaccine immune responses; therefore, the subpopulation level should be also considered during vaccine design that can trigger an optimum immune response against a specific disease [92].

5. Conclusion

The study describes numerous computational tools that are widely used for drug and vaccine design against infectious viral diseases. However, improving ideas and methodologies can solve problems and provide system-level interconnected data. A promising idea accelerates the rational design of drug candidates that are used in the specific subfield. Here, we focused on computational methods that will help and provide a clear idea of modern drug and vaccine design approaches adequately.

Acknowledgements

The authors are in debt to King Abdulaziz University for their encouragements and unlimited support.

Conflict of interest

The authors declare no conflict of interest.

IntechOpen

Author details

Mohammed Othman Aljahdali*, Mohammad Habibur Rahman Molla
and Foysal Ahammad
Faculty of Science, Department of Biological Sciences, King Abdulaziz University,
Jeddah, Saudi Arabia

*Address all correspondence to: moaljhdali@kau.edu.sa

IntechOpen

© 2022 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] Shaw WR, Catteruccia F. Vector biology meets disease control: Using basic research to fight vector-borne diseases. *Nature Microbiology*. 2019;**4**(1):20-34. DOI: 10.1038/s41564-018-0214-7
- [2] Rogers DJ, Packer MJ. Vector-borne diseases, models, and global change. *Lancet*. 1993;**342**(8882):1282-1284. DOI: 10.1016/0140-6736(93)92367-3
- [3] Sainz Á, Roura X, Estrada-Peña A, Kohn B, Harrus S, Solano-Gallego L. Guideline for veterinary practitioners on canine ehrlichiosis and anaplasmosis in Europe. *Parasites Vectors*. 2015;**8**(1):1-20. DOI: 10.1186/S13071-015-0649-0
- [4] Mourya DT, Yadav PD, Ullas PT, Bhardwaj SD, Sahay RR, Chadha MS, et al. Emerging/re-emerging viral diseases & new viruses on the Indian horizon. *Indian Journal of Medical Research*. 2019;**149**(4):447. DOI: 10.4103/IJMR.IJMR_1239_18
- [5] Kruse H, Kirkemo A-M, Handeland K. Wildlife as source of zoonotic infections. *Emerging Infectious Diseases*. 2004;**10**(12):2067-2072. DOI: 10.3201/EID1012.040707
- [6] Pulendran B, Arunachalam PS, O'Hagan DT. Emerging concepts in the science of vaccine adjuvants. *Nature Reviews Drug Discovery*. 2021;**20**(6):1. DOI: 10.1038/S41573-021-00163-Y
- [7] Terry FE, Moise L, Martin RF, Torres M, Pilotte N, Williams SA, et al. Time for T? Immunoinformatics addresses vaccine design for neglected tropical and emerging infectious diseases. *Expert Review of Vaccines*. 2015;**14**(1):21-35. DOI: 10.1586/14760584.2015.955478
- [8] V'kovski P, Kratzel A, Steiner S, Stalder H, Thiel V. Coronavirus biology and replication: Implications for SARS-CoV-2. *Nature Reviews Microbiology*. 2020;**19**(3):155-170. DOI: 10.1038/s41579-020-00468-6
- [9] Modrow S, Falke D, Truyen U, Schätzl H. Viruses with single-stranded, positive-sense RNA genomes. *Molecular Virology*. 2013:185-349. DOI: 10.1007/978-3-642-20718-1_14
- [10] Khan T, Lawrence AJ, Azad I, Raza S, Joshi S, Khan AR. Computational drug designing and prediction of important parameters using *in silico* methods- A review. *Current Computer-Aided Drug Design*. 2019;**15**(5):384-397. DOI: 10.2174/1573399815666190326120006
- [11] Sweiti H, Wiegand F, Bug C, Vogel M, Lavie F, Winiger-Candolfi I, et al. Physicians in the pharmaceutical industry: Their roles, motivations, and perspectives. *Drug Discovery Today*. 2019;**24**(9):1865-1870. DOI: 10.1016/J.DRUDIS.2019.05.021
- [12] Rifaioglu AS, Atas H, Martin MJ, Cetin-Atalay R, Atalay V, Doğan T. Recent applications of deep learning and machine intelligence on *in silico* drug discovery: Methods, tools and databases. *Briefings in Bioinformatics*. 2019;**20**(5):1878-1912. DOI: 10.1093/BIB/BBY061
- [13] Furuse Y. Analysis of research intensity on infectious disease by disease burden reveals which infectious diseases are neglected by researchers. *Proceedings of the National Academy of Science of the United States of America*. 2019;**116**(2):478-483. DOI: 10.1073/PNAS.1814484116
- [14] Dasaraju PV, Liu C. Infections of the Respiratory System. In: Baron S, editor.

- Medical Microbiology. 4th ed. Galveston (TX): University of Texas Medical Branch at Galveston; 1996
- [15] Relman DA, Lipsitch M. Microbiome as a tool and a target in the effort to address antimicrobial resistance. *Proceedings of the National Academy of Sciences of the United States of America*. 2018;**115**(51):12902-12910. DOI: 10.1073/PNAS.1717163115
- [16] Janeway CA Jr, Travers P, Walport M, Shlomchik MJ. *Infectious Agents and How They cause Disease*. 5th ed. New York: Garland Science; 2001. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK27114/> [Accessed: October 07, 2021]
- [17] Edemekong PF, Huang B. Epidemiology of prevention of communicable diseases. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; Jan 2021
- [18] Funk S, Gilad E, Watkins C, Jansen VA. The spread of awareness and its impact on epidemic outbreaks. *Proceedings of the National Academy of Sciences of the United States of America*. 2009;**106**(16):6872-6877. DOI: 10.1073/pnas.0810762106
- [19] Rohr JR, Barrett CB, Civitello DJ, Craft ME, Delius B, GA DL, et al. Emerging human infectious diseases and the links to global food production. *Nature Sustainability*. 2019;**2**(6):445-456. DOI: 10.1038/s41893-019-0293-3
- [20] Greger M. The human/animal interface: Emergence and resurgence of zoonotic infectious diseases. *Critical Reviews in Microbiology*. 2007;**33**(4):243-299. DOI: 10.1080/10408410701647594
- [21] Smith KF, Goldberg M, Rosenthal S, Carlson L, Chen J, Chen C, et al. Global rise in human infectious disease outbreaks. *Journal of the Royal Society Interface*. 2014;**11**(101):20140950. DOI: 10.1098/RSIF.2014.0950
- [22] Bhutta ZA, Sommerfeld J, Lassi ZS, Salam RA, Das JK. Global burden, distribution, and interventions for infectious diseases of poverty. *Infectious Diseases of Poverty*. 2014;**3**(1):1-7. DOI: 10.1186/2049-9957-3-21
- [23] Alghamdi AA. Impact of the COVID-19 pandemic on the social and educational aspects of Saudi university students' lives. *PLoS One*. 2021;**16**(4):e0250026. DOI: 10.1371/JOURNAL.PONE.0250026
- [24] Mathers CD, Bernard C, Iburg KM, Inoue M, Ma Fat D, Shibuya K, et al. Global Burden of Disease in 2002: Data sources, methods and results [Internet]. 2003. Available from: www.who.int/evidence/bod [Accessed: October 07, 2021]
- [25] Ung L, Chodosh J. Foundational concepts in the biology of bacterial keratitis. *Experimental Eye Research*. 2021;**209**:108647. DOI: 10.1016/J.EXER.2021.108647
- [26] Mourya DT, Yadav PD, Ullas PT, Bhardwaj SD, Sahay RR, Chadha MS, et al. Emerging/re-emerging viral diseases & new viruses on the Indian horizon. *Indian Journal of Medical Research*. 2019;**149**(4):447-467. DOI: 10.4103/IJMR.IJMR_1239_18
- [27] Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, et al. Global trends in emerging infectious diseases. *Nature*. 2008;**451**(7181):990. DOI: 10.1038/NATURE06536
- [28] Li H, Yang Y, Hong W, Huang M, Wu M, Zhao X. Applications of genome editing technology in the targeted therapy of human diseases: Mechanisms, advances and prospects. *Signal Transduction and*

- Targeted Therapy. 2020;5(1):1-23. DOI: 10.1038/s41392-019-0089-y
- [29] Pinter-Wollman N, Jelić A, Wells NM. The impact of the built environment on health behaviours and disease transmission in social systems. *Philosophical Transactions of the Royal Society B Biological Sciences*. 2018;373(1753). DOI: 10.1098/RSTB.2017.0245
- [30] Gomez-Lus R, Clavel A, Castillo J, Seral C, Rubio C. Emerging and reemerging pathogens. *International Journal of Antimicrobial Agents*. 2000;16(3):335-339. DOI: 10.1016/S0924-8579(00)00261-2
- [31] Weaver SC, Costa F, Garcia-Blanco MA, Ko AI, Ribeiro GS, Saade G, et al. Zika virus: History, emergence, biology, and prospects for control. *Antiviral Research*. 2016;130:69. DOI: 10.1016/J.ANTIVIRAL.2016.03.010
- [32] Meyding-Lamadé U, Craemer E, Schnitzler P. Emerging and re-emerging viruses affecting the nervous system. *Neurological Research and Practice*. 2019;1(1):1-9. DOI: 10.1186/S42466-019-0020-6
- [33] Dabhu Kumar J, Jian L, Rong H, Hua Z. Emerging and reemerging human viral diseases. *Annals of Microbiology and Research*. 2018;2(1):31-44. DOI: 10.36959/958/567
- [34] Bloom DE, Cadarette D. Infectious disease threats in the twenty-first century: Strengthening the global response. *Frontiers in Immunology*. 2019;10:549. DOI: 10.3389/FIMMU.2019.00549
- [35] Feldman C, Shaddock E. Epidemiology of lower respiratory tract infections in adults. *Expert Review of Respiratory Medicine*. 2019;13(1):63-77. DOI: 10.1080/17476348.2019.1555040
- [36] Morse SS. Factors in the emergence of infectious diseases. *Emerging Infectious Diseases*. 1995;1(1):7-15. DOI: 10.3201/EID0101.950102
- [37] Schneider UA, Havlik P, Schmid E, Valin H, Mosnier A, Obersteiner M, et al. Impacts of population growth, economic development, and technical change on global food production and consumption. *Agricultural Systems*. 2011;104(2):204-215. DOI: 10.1016/J.AGSY.2010.11.003
- [38] Reed KD. Respiratory tract infections: A clinical approach. *Molecular Medical Microbiology*. 2015;3:1499-1506. DOI: 10.1016/B978-0-12-397169-2.00084-6
- [39] Ogilvie M. Molecular techniques should not now replace cell culture in diagnostic virology laboratories. *Reviews in Medical Virology*. 2001;11(6):351-354. DOI: 10.1002/RMV.335
- [40] Zampieri F. The impact of modern medicine on human evolution. In: Tibayrene M, Ayala FJ, editors. *On Human Nature Biology, Psychology, Ethics, Politics, and Religion*. Academic Press; 2017. pp. 707-727. DOI: 10.1016/B978-0-12-420190-3.00043-0. Available from: <https://www.semanticscholar.org/paper/The-Impact-of-Modern-Medicine-on-Human-Evolution-Zampieri/af39aabd82bbeace3588e98cfb23538e77946e66>
- [41] Morens DM, Folkers GK, Fauci AS. The challenge of emerging and re-emerging infectious diseases. *Nature*. 2004;430(6996):242-249. DOI: 10.1038/nature02759
- [42] Plotkin BJ, Kimball AM. Designing an international policy and legal framework for the control of emerging infectious diseases: First steps. *Emerging Infectious Diseases*. 1997;3(1):1-9. DOI: 10.3201/EID0301.970101

- [43] Blastland M, Freeman ALJ, van der Linden S, Marteau TM, Spiegelhalter D. Five rules for evidence communication. *Nature*. 2020;**587**(7834):362-364. DOI: 10.1038/D41586-020-03189-1
- [44] Araj R, Alqasrawi S, Samy S, Alwahdanee G, Walid J, Mofleh J, et al. Preventing emerging and re-emerging infections in the eastern mediterranean region: Gaps, challenges, and priorities. *JMIR Public Health and Surveillance*. 2019;**5**(4):e14348. DOI: 10.2196/14348
- [45] Hughes JM, Wilson ME, Pike BL, Saylor KE, Fair JN, LeBreton M, et al. The origin and prevention of pandemics. *Clinical Infectious Diseases*. 2010;**50**(12):1636-1640. DOI: 10.1086/652860
- [46] Mahady GB. Medicinal plants for the prevention and treatment of bacterial infections. *Current Pharmaceutical Design*. 2005;**11**(19):2405-2427. DOI: 10.2174/1381612054367481
- [47] Aljahdali MO, Molla MHR, Ahammad F. Compounds identified from marine mangrove plant (*Avicennia alba*) as potential antiviral drug candidates against WDSV, an in-silico approach. *Marine Drugs*. 2021;**19**(5):253. DOI: 10.3390/MD19050253
- [48] Wang L, Wang Y, Jin S, Wu Z, Chin DP, Koplan JP, et al. Emergence and control of infectious diseases in China. *The Lancet*. 2008;**372**(9649):1598-1605. DOI: 10.1016/S0140-6736(08)61365-3
- [49] Mukherjee S. Emerging infectious diseases: Epidemiological perspective. *Indian Journal of Dermatology*. Sep. 2017;**62**(5):459-467. DOI: 10.4103/IJD.IJD_379_17
- [50] Achan J, Talisuna AO, Erhart A, Yeka A, Tibenderana JK, Baliraine FN, et al. Quinine, an old anti-malarial drug in a modern world: Role in the treatment of malaria. *Malaria Journal*. 2011;**10**:144. DOI: 10.1186/1475-2875-10-144
- [51] Kutlushina A, Khakimova A, Madzhidov T, Polishchuk P. Ligand-based pharmacophore modeling using novel 3D pharmacophore signatures. *Molecules*. 2018;**23**(12):3094. DOI: 10.3390/MOLECULES23123094
- [52] Duerr TJ, Comellas E, Jeon EK, Farkas JE, Joetzjer M, Garnier J, et al. 3D visualization of macromolecule synthesis. *Elife*. 2020;**9**:e60354. DOI: 10.1101/2020.06.24.169300
- [53] Yang SY. Pharmacophore modeling and applications in drug discovery: Challenges and recent advances. *Drug Discovery Today*. 2010;**15**(11-12):444-450. DOI: 10.1016/J.DRUDIS.2010.03.013
- [54] Ferreira LG, Dos Santos RN, Oliva G, Andricopulo AD. Molecular docking and structure-based drug design strategies. *Molecules*. 2015;**20**(7):13384-13421. DOI: 10.3390/MOLECULES200713384
- [55] Wang W, Ye Z, Gao H, Ouyang D. Computational pharmaceutics - A new paradigm of drug delivery. *Journal of Controlled Release*. 2021;**338**:119-136. DOI: 10.1016/J.JCONREL.2021.08.030
- [56] Sousa SF, Fernandes PA, Ramos MJ. Protein-ligand docking: Current status and future challenges. *Proteins*. 2006;**65**(1):15-26. DOI: 10.1002/PROT.21082
- [57] Khanam H, Ali A, Asif M, Shamsuzzaman. Neurodegenerative diseases linked to misfolded proteins and their therapeutic approaches: A review. *European Journal of Medicinal Chemistry*. 2016;**124**:1121-1141. DOI: 10.1016/J.EJMECH.2016.08.006
- [58] Sliwoski G, Kothiwale S, Meiler J, Lowe EW Jr. Computational methods

in drug discovery. *Pharmacological Reviews*. 2014;**66**(1):334. DOI: 10.1124/PR.112.007336

[59] Atanasov AG, Waltenberger B, Pferschy-Wenzig E-M, Linder T, Wawrosch C, Uhrin P, et al. Discovery and resupply of pharmacologically active plant-derived natural products: A review. *Biotechnology Advances*. 2015;**33**(8):1582-1614. DOI: 10.1016/j.biotechadv.2015.08.001

[60] Mori T, Miyashita N, Im W, Feig M, Sugita Y. Molecular dynamics simulations of biological membranes and membrane proteins using enhanced conformational sampling algorithms. *Biochimica et Biophysica Acta (BBA) - Biomembranes*. 2016;**1858**(7):1635-1651. DOI: 10.1016/J.BBAMEM.2015.12.032

[61] Pritchett DB, Seeburg PH. The role of molecular biology in drug discovery and design. In: Harris TJR, editor. *Protein Production by Biotechnology*. Boston: Springer; 1990. pp. 181-190. DOI: 10.1007/978-1-4613-1565-0_14

[62] Swegat W, Schlitter J, Krüger P, Wollmer A. MD simulation of protein-ligand interaction: Formation and dissociation of an insulin-phenol complex. *Biophysical Journal*. 2003;**84**(3):1493-1506. DOI: 10.1016/S0006-3495(03)74962-5

[63] Lee S, Bhattacharya S, Tate CG, Grisshammer R, Vaidehi N. Structural dynamics and thermostabilization of neurotensin receptor 1. *The Journal of Physical Chemistry B*. 2015;**119**(15):4917-4928. DOI: 10.1021/JP510735F

[64] González MA. Force fields and molecular dynamics simulations. *Collection SFN*. 2011;**12**:169-200. DOI: 10.1051/sfn/201112009

[65] Lesiuk M. Implementation of the coupled-cluster method with single,

double, and triple excitations using tensor decompositions. *Journal of Chemical Theory and Computation*. 2019;**16**(1):453-467. DOI: 10.1021/ACS.JCTC.9B00985

[66] Arodola OA, Soliman ME. Quantum mechanics implementation in drug-design workflows: Does it really help? *Drug Design, Development and Therapy*. 2017;**11**:2551-2564. DOI: 10.2147/DDDT.S126344

[67] Taylor D. The pharmaceutical industry and the future of drug development. *Issues in Environmental Science and Technology*. 2015;**2016**(41):1-33. DOI: 10.1039/9781782622345-00001

[68] Lonsdale R, Fort RM, Rydberg P, Harvey JN, Mulholland AJ. Quantum mechanics/molecular mechanics modeling of drug metabolism: Mexiletine N-hydroxylation by cytochrome P450 1A2. *Chemical Research in Toxicology*. 2016;**29**(6):963-971. DOI: 10.1021/ACS.CHEMRESTOX.5B00514

[69] Kalaszi A, Imre G, Jakli I, Farkas O. Identification of the bioactive conformation for mucin epitope peptides. *Journal of Molecular Structure: THEOCHEM*. 2007;**823**(1-3):16-27. DOI: 10.1016/J.THEOCHEM.2007.08.016

[70] Salo-Ahen OMH, Alanko I, Bhadane R, Bonvin AMJJ, Honorato RV, Hossain S, et al. Molecular dynamics simulations in drug discovery and pharmaceutical development. *Process*. 2021;**9**(1):71. DOI: 10.3390/PR9010071

[71] Fedorov DG, Nagata T, Kitaura K. Exploring chemistry with the fragment molecular orbital method. *Physical Chemistry Chemical Physics*. 2012;**14**(21):7562-7577. DOI: 10.1039/C2CP23784A

[72] Wu F, Zhou Y, Li L, Shen X, Chen G, Wang X, et al. Computational

approaches in preclinical studies on drug discovery and development. *Frontiers in Chemistry*. 2020;**8**:726. DOI: 10.3389/fchem.2020.00726

[73] Sutton RT, Pincock D, Baumgart DC, Sadowski DC, Fedorak RN, Kroeker KI. An overview of clinical decision support systems: Benefits, risks, and strategies for success. *npj Digital Medicine*. 2020;**3**(1):1-10. DOI: 10.1038/s41746-020-0221-y

[74] Kok-Yong S, Lawrence L. Drug Distribution and Drug Elimination. *Basic Pharmacokinetic Concepts and Some Clinical Applications*. London: IntechOpen; 2015. DOI: 10.5772/59929

[75] Vendel E, Rottschäfer V, de Lange ECM. The need for mathematical modelling of spatial drug distribution within the brain. *Fluids and Barriers of the CNS*. 2019;**16**(1):1-33. DOI: 10.1186/S12987-019-0133-X

[76] Prabu SL, Suriyaprakash TNK, Ruckmani K, Thirumurugan R. Biopharmaceutics and Pharmacokinetics. *Basic Pharmacokinetic Concepts and Some Clinical Applications*. IntechOpen; 2015. DOI: 10.5772/61160. Available from: <https://www.intechopen.com/chapters/48805>

[77] Henninot A, Collins JC, Nuss JM. The current state of peptide drug discovery: Back to the future? *Journal of Medicinal Chemistry*. 2017;**61**(4):1382-1414. DOI: 10.1021/ACS.JMEDCHEM.7B00318

[78] Lau JL, Dunn MK. Therapeutic peptides: Historical perspectives, current development trends, and future directions. *Bioorganic & Medicinal Chemistry*. 2018;**26**(10):2700-2707. DOI: 10.1016/J.BMC.2017.06.052

[79] Zharkova MS, Orlov DS, Golubeva OY, Chakchir OB, Eliseev IE, Grinchuk TM, et al. Application of

antimicrobial peptides of the innate immune system in combination with conventional antibiotics—A novel way to combat antibiotic resistance? *Frontiers in Cellular and Infection Microbiology*. 2019;**9**:128. DOI: 10.3389/FCIMB.2019.00128

[80] Debnath S, Barnaby DP, Coppa K, Makhnevich A, Kim EJ, Chatterjee S, et al. Machine learning to assist clinical decision-making during the COVID-19 pandemic. *Bioelectronic Medicine*. 2020;**6**(1):1-8. DOI: 10.1186/S42234-020-00050-8

[81] Eichmann M, de Ru A, van Veelen PA, Peakman M, Kronenberg-Versteeg D. Identification and characterisation of peptide binding motifs of six autoimmune disease-associated human leukocyte antigen-class I molecules including HLA-B*39:06. *Tissue Antigens*. 2014;**84**(4):378-388. DOI: 10.1111/TAN.12413

[82] Curtiss R. Expert review of vaccines vaccine design: Innovative approaches and novel strategies. *Expert Review Vaccines*. 2011;**10**(10):1385-1387. DOI: 10.1586/erv.11.114

[83] De Groot AS, Moise L, Terry F, Guterrez AH, Hindocha P, Richard G, et al. Better epitope discovery, precision immune engineering, and accelerated vaccine design using immunoinformatics tools. *Frontiers in Immunology*. 2020;**11**:442. DOI: 10.3389/FIMMU.2020.00442

[84] Urrutia-Baca VH, Gomez-Flores R, De La Garza-Ramos MA, Tamez-Guerra P, Lucio-Sauceda DG, Rodríguez-padilla MC. Immunoinformatics approach to design a novel epitope-based oral vaccine against *Helicobacter pylori*. *Journal of Computational Biology*. 2019;**26**(10):1177-1190. DOI: 10.1089/CMB.2019.0062

[85] Ali SA, Almofti YA, Khoubieb KA. Immunoinformatics approach for

- multiepitopes vaccine prediction against glycoprotein B of avian infectious laryngotracheitis virus. *Advances in Bioinformatics*. 2019;**2019**:1270485. DOI: 10.1155/2019/1270485
- [92] Huang T, Shu Y, Cai Y-D. Genetic differences among ethnic groups. *BMC Genomics*. 2015;**16**(1):1-10. DOI: 10.1186/S12864-015-2328-0
- [86] Russo G, Reche P, Pennisi M, Pappalardo F. The combination of artificial intelligence and systems biology for intelligent vaccine design. *Expert Opinion on Drug Discovery*. 2020;**15**:11, 1267-1281. DOI: 10.1080/17460441.2020.1791076
- [87] Arora P, Kumar H, Panigrahi BK. Prediction and analysis of COVID-19 positive cases using deep learning models: A descriptive case study of India. *Chaos, Solitons & Fractals*. 2020;**139**:110017. DOI: 10.1016/J.CHAOS.2020.110017
- [88] Nazem-Bokaee H, Hom EFY, Warden AC, Mathews S, Gueidan C. Towards a systems biology approach to understanding the lichen symbiosis: Opportunities and challenges of implementing network modelling. *Frontiers in Microbiology*. 2021;**12**:1028. DOI: 10.3389/FMICB.2021.667864
- [89] Jerez CA. The use of genomics, proteomics and other OMICS technologies for the global understanding of biomining microorganisms. *Hydrometallurgy*. 2008;**94**(1-4):162-169. DOI: 10.1016/J.HYDROMET.2008.05.032
- [90] Oli AN, Obialor WO, Ifeanyichukwu MO, Odimegwu DC, Okoyeh JN, Emechebe GO, et al. Immunoinformatics and vaccine development: An overview. *Immunotargets and Therapy*. 2020;**9**: 13-30. DOI: 10.2147/ITT.S241064
- [91] Filatov DA, Charlesworth D. DNA polymorphism, haplotype structure and balancing selection in the *Leavenworthia PgiC* locus. *Genetics*. 1999;**153**(3):1423-1434